

Jun Huang

List of Publications by Year in descending order

Source: <https://exaly.com/author-pdf/4636335/publications.pdf>

Version: 2024-02-01

15
papers

119
citations

1306789

7
h-index

1281420

11
g-index

17
all docs

17
docs citations

17
times ranked

137
citing authors

#	ARTICLE	IF	CITATIONS
1	Left Ventricular Systolic Function Changes in Primary Hypertension Patients Detected by the Strain of Different Myocardium Layers. <i>Medicine (United States)</i> , 2016, 95, e2440.	0.4	18
2	Left ventricular short-axis systolic function changes in patients with hypertrophic cardiomyopathy detected by two-dimensional speckle tracking imaging. <i>BMC Cardiovascular Disorders</i> , 2018, 18, 13.	0.7	14
3	Left ventricular systolic function changes in hypertrophic cardiomyopathy patients detected by the strain of different myocardium layers and longitudinal rotation. <i>BMC Cardiovascular Disorders</i> , 2017, 17, 214.	0.7	13
4	Global myocardial work: A new way to detect subclinical myocardial dysfunction with normal left ventricle ejection fraction in essential hypertension patients: Compared with myocardial layer-specific strain analysis. <i>Echocardiography</i> , 2021, 38, 850-860.	0.3	13
5	Assessment of the Rotation Motion at the Papillary Muscle Short-Axis Plane with Normal Subjects by Two-Dimensional Speckle Tracking Imaging: A Basic Clinical Study. <i>PLoS ONE</i> , 2013, 8, e83071.	1.1	9
6	Peak systolic longitudinal rotation: a new tool for detecting left ventricular systolic function in patients with type 2 diabetes mellitus by two-dimensional speckle tracking echocardiography. <i>BMC Cardiovascular Disorders</i> , 2019, 19, 137.	0.7	9
7	Right atrial function assessed by volume-derived values and speckle tracking echocardiography in patients with hypertrophic cardiomyopathy. <i>BMC Cardiovascular Disorders</i> , 2020, 20, 335.	0.7	8
8	Assessment of subclinical left ventricle myocardial dysfunction using global myocardial work in type 2 diabetes mellitus patients with preserved left ventricle ejection fraction. <i>Diabetology and Metabolic Syndrome</i> , 2022, 14, 17.	1.2	8
9	Assessment of subclinical left atrial myocardial dysfunction in essential hypertension patients with normal left ventricle function by two-dimensional strain and volume-derived variables. <i>Journal of Clinical Ultrasound</i> , 2021, 49, 659-666.	0.4	7
10	Left ventricular longitudinal function assessment in rabbits after acute occlusion of left anterior descending coronary artery by two-dimensional speckle tracking imaging. <i>BMC Cardiovascular Disorders</i> , 2017, 17, 219.	0.7	5
11	Layer-specific speckle tracking analysis of left ventricular systolic function and synchrony in maintenance hemodialysis patients. <i>BMC Cardiovascular Disorders</i> , 2020, 20, 126.	0.7	4
12	Assessment of left atrial function in patients with hypertrophic cardiomyopathy using two-dimensional strain: a comparison with volume-derived values. <i>Minerva Cardiology and Angiology</i> , 2017, 66, 26-37.	0.4	3
13	Evaluation of right ventricular systolic and diastolic dysfunctions in patients with type 2 diabetes mellitus with poor glycemic control by layer specific global longitudinal strain and strain rate. <i>Diabetology and Metabolic Syndrome</i> , 2022, 14, 49.	1.2	2
14	Girl With Abdominal Pain. <i>Annals of Emergency Medicine</i> , 2020, 76, e11-e12.	0.3	0
15	Neonate With Vomiting and Jaundice. <i>Annals of Emergency Medicine</i> , 2020, 75, 734-754.	0.3	0