

Shigeto Ueda

List of Publications by Year in descending order

Source: <https://exaly.com/author-pdf/4650361/publications.pdf>

Version: 2024-02-01

21
papers

1,083
citations

623734

14
h-index

794594

19
g-index

21
all docs

21
docs citations

21
times ranked

1726
citing authors

#	ARTICLE	IF	CITATIONS
1	Early Therapeutic Prediction Based on Tumor Hemodynamic Response Imaging: Clinical Studies in Breast Cancer with Time-Resolved Diffuse Optical Spectroscopy. <i>Applied Sciences (Switzerland)</i> , 2019, 9, 3.	2.5	12
2	Recurrent Angiosarcoma of the Breast Treated with Gemcitabine and Paclitaxel. <i>Nihon Rinsho Geka Gakkai Zasshi (Journal of Japan Surgical Association)</i> , 2018, 79, 687-691.	0.0	0
3	Bevacizumab Induces Acute Hypoxia and Cancer Progression in Patients with Refractory Breast Cancer: Multimodal Functional Imaging and Multiplex Cytokine Analysis. <i>Clinical Cancer Research</i> , 2017, 23, 5769-5778.	7.0	70
4	In vivo imaging of eribulin-induced reoxygenation in advanced breast cancer patients: a comparison to bevacizumab. <i>British Journal of Cancer</i> , 2016, 114, 1212-1218.	6.4	82
5	Near-Infrared Diffuse Optical Imaging for Early Prediction of Breast Cancer Response to Neoadjuvant Chemotherapy: A Comparative Study Using ¹⁸ F-FDG PET/CT. <i>Journal of Nuclear Medicine</i> , 2016, 57, 1189-1195.	5.0	18
6	Neoadjuvant triweekly nanoparticle albumin-bound paclitaxel followed by epirubicin and cyclophosphamide for Stage II/III HER2-negative breast cancer: evaluation of efficacy and safety. <i>Japanese Journal of Clinical Oncology</i> , 2015, 45, 642-649.	1.3	15
7	Phase I study of HER3 targeted antibody patritumab in combination with trastuzumab and paclitaxel in patients with HER2-overexpressing metastatic breast cancer (MBC).. <i>Journal of Clinical Oncology</i> , 2015, 33, 584-584.	1.6	4
8	Optical Imaging for Monitoring Tumor Oxygenation Response after Initiation of Single-Agent Bevacizumab followed by Cytotoxic Chemotherapy in Breast Cancer Patients. <i>PLoS ONE</i> , 2014, 9, e98715.	2.5	20
9	Clinicopathological and prognostic impact of imaging of breast cancer angiogenesis and hypoxia using diffuse optical spectroscopy. <i>Cancer Science</i> , 2014, 105, 833-839.	3.9	15
10	Near-infrared optical imaging of cancer vascular remodeling after antiangiogenic therapy. <i>Breast Cancer</i> , 2014, 21, 776-779.	2.9	2
11	Optical imaging for monitoring tumor oxygenation response after initiation of single-agent bevacizumab followed by neoadjuvant chemotherapy in breast cancer patients.. <i>Journal of Clinical Oncology</i> , 2014, 32, e13503-e13503.	1.6	0
12	Optical imaging of tumor vascularity associated with proliferation and glucose metabolism in early breast cancer: clinical application of total hemoglobin measurements in the breast. <i>BMC Cancer</i> , 2013, 13, 514.	2.6	27
13	Genomic Profiling Shows Increased Glucose Metabolism in Luminal B Breast Cancer. <i>Journal of Breast Cancer</i> , 2013, 16, 342.	1.9	3
14	Early metabolic response to neoadjuvant letrozole, measured by FDG PET/CT, is correlated with a decrease in the Ki67 labeling index in patients with hormone receptor-positive primary breast cancer: a pilot study. <i>Breast Cancer</i> , 2011, 18, 299-308.	2.9	41
15	Optical imaging of breast cancer oxyhemoglobin flare correlates with neoadjuvant chemotherapy response one day after starting treatment. <i>Proceedings of the National Academy of Sciences of the United States of America</i> , 2011, 108, 14626-14631.	7.1	130
16	Utility of 18F-fluoro-deoxyglucose emission tomography/computed tomography fusion imaging (18F-FDG PET/CT) in combination with ultrasonography for axillary staging in primary breast cancer. <i>BMC Cancer</i> , 2008, 8, 165.	2.6	104
17	Expression of centromere protein F (CENP-F) associated with higher FDG uptake on PET/CT, detected by cDNA microarray, predicts high-risk patients with primary breast cancer. <i>BMC Cancer</i> , 2008, 8, 384.	2.6	32
18	Clinicopathological and Prognostic Relevance of Uptake Level using 18F-fluorodeoxyglucose Positron Emission Tomography/Computed Tomography Fusion Imaging (18F-FDG PET/CT) in Primary Breast Cancer. <i>Japanese Journal of Clinical Oncology</i> , 2008, 38, 250-258.	1.3	202

#	ARTICLE	IF	CITATIONS
19	Alternative tyrosine phosphorylation of signaling kinases according to hormone receptor status in breast cancer overexpressing the insulin-like growth factor receptor type 1. <i>Cancer Science</i> , 2006, 97, 597-604.	3.9	38
20	Potential crosstalk between insulin-like growth factor receptor type 1 and epidermal growth factor receptor in progression and metastasis of pancreatic cancer. <i>Modern Pathology</i> , 2006, 19, 788-796.	5.5	52
21	The Correlation Between Cytoplasmic Overexpression of Epidermal Growth Factor Receptor and Tumor Aggressiveness. <i>Pancreas</i> , 2004, 29, e1-e8.	1.1	216