

# Dohee Kwon

## List of Publications by Year in descending order

Source: <https://exaly.com/author-pdf/4657768/publications.pdf>

Version: 2024-02-01

15  
papers

664  
citations

840119

11  
h-index

996533

15  
g-index

15  
all docs

15  
docs citations

15  
times ranked

1525  
citing authors

#	ARTICLE	IF	CITATIONS
1	Clinicopathological analysis of programmed cell death 1 and programmed cell death ligand 1 expression in the tumour microenvironments of diffuse large B cell lymphomas. <i>Histopathology</i> , 2016, 68, 1079-1089.	1.6	135
2	PD-L1 expression is associated with epithelial-to-mesenchymal transition in adenocarcinoma of the lung. <i>Human Pathology</i> , 2016, 58, 7-14.	1.1	135
3	Comparative analysis of PD-L1 expression between primary and metastatic pulmonary adenocarcinomas. <i>European Journal of Cancer</i> , 2017, 75, 141-149.	1.3	84
4	NTRK and RET fusion-directed therapy in pediatric thyroid cancer yields a tumor response and radioiodine uptake. <i>Journal of Clinical Investigation</i> , 2021, 131, .	3.9	62
5	Overexpression of endoplasmic reticulum stress-related proteins, XBP1s and GRP78, predicts poor prognosis in pulmonary adenocarcinoma. <i>Lung Cancer</i> , 2018, 122, 131-137.	0.9	44
6	MYC and BCL2 overexpression is associated with a higher class of Memorial Sloan-Kettering Cancer Center prognostic model and poor clinical outcome in primary diffuse large B-cell lymphoma of the central nervous system. <i>BMC Cancer</i> , 2016, 16, 363.	1.1	37
7	Prognostic implications of tumor-infiltrating macrophages, M2 macrophages, regulatory T-cells, and indoleamine 2,3-dioxygenase-positive cells in primary diffuse large B-cell lymphoma of the central nervous system. <i>Oncolmmunology</i> , 2018, 7, e1442164.	2.1	34
8	MET exon 14 skipping mutation in triple-negative pulmonary adenocarcinomas and pleomorphic carcinomas: An analysis of intratumoral MET status heterogeneity and clinicopathological characteristics. <i>Lung Cancer</i> , 2017, 106, 131-137.	0.9	30
9	High tumoral PD-L1 expression and low PD-1 or CD8 tumor-infiltrating lymphocytes are predictive of a poor prognosis in primary diffuse large B-cell lymphoma of the central nervous system. <i>Oncolmmunology</i> , 2019, 8, e1626653.	2.1	30
10	Programmed death ligand-1 expression and its prognostic role in esophageal squamous cell carcinoma. <i>World Journal of Gastroenterology</i> , 2016, 22, 8389.	1.4	22
11	Clinicopathological features of programmed cell death-1 and programmed cell death-ligand-1 expression in the tumor cells and tumor microenvironment of angioimmunoblastic T cell lymphoma and peripheral T cell lymphoma not otherwise specified. <i>Virchows Archiv Fur Pathologische Anatomie Und Physiologie Und Fur Klinische Medizin</i> , 2020, 477, 131-142.	1.4	14
12	Prognostic implications of FGFR1 and MYC status in esophageal squamous cell carcinoma. <i>World Journal of Gastroenterology</i> , 2016, 22, 9803.	1.4	11
13	Bronchovascular injury associated with clinically significant hemoptysis after CT-guided core biopsy of the lung: Radiologic and histopathologic analysis. <i>PLoS ONE</i> , 2018, 13, e0204064.	1.1	11
14	Blastic plasmacytoid dendritic cell neoplasm with unusual extracutaneous manifestation. <i>Medicine (United States)</i> , 2019, 98, e14344.	0.4	10
15	Pediatric Case Report on an Interstitial Lung Disease with a Novel Mutation of SFTPC Successfully Treated with Lung Transplantation. <i>Journal of Korean Medical Science</i> , 2018, 33, e159.	1.1	5