

Youhua Liu

List of Publications by Year in descending order

Source: <https://exaly.com/author-pdf/4665244/publications.pdf>

Version: 2024-02-01

187
papers

19,050
citations

9234

74
h-index

12233

133
g-index

191
all docs

191
docs citations

191
times ranked

14689
citing authors

#	ARTICLE	IF	CITATIONS
1	CXCR4 induces podocyte injury and proteinuria by activating β -catenin signaling. <i>Theranostics</i> , 2022, 12, 767-781.	4.6	20
2	A Klotho-derived peptide protects against kidney fibrosis by targeting TGF- β signaling. <i>Nature Communications</i> , 2022, 13, 438.	5.8	53
3	Ageing, cellular senescence and chronic kidney disease: experimental evidence. <i>Current Opinion in Nephrology and Hypertension</i> , 2022, 31, 235-243.	1.0	14
4	Matrix Metalloproteinase-10 in Kidney Injury Repair and Disease. <i>International Journal of Molecular Sciences</i> , 2022, 23, 2131.	1.8	5
5	β -catenin-controlled tubular cell-derived exosomes play a key role in fibroblast activation via the OPN-CD44 axis. <i>Journal of Extracellular Vesicles</i> , 2022, 11, e12203.	5.5	31
6	Non-canonical Wnt/calcium signaling is protective against podocyte injury and glomerulosclerosis. <i>Kidney International</i> , 2022, 102, 96-107.	2.6	7
7	High Fat Diet Induces Kidney Injury via Stimulating Wnt/ β -Catenin Signaling. <i>Frontiers in Medicine</i> , 2022, 9, 851618.	1.2	7
8	Follistatin-like 1 (FSTL1) interacts with Wnt ligands and Frizzled receptors to enhance Wnt/ β -catenin signaling in obstructed kidneys in vivo. <i>Journal of Biological Chemistry</i> , 2022, 298, 102010.	1.6	13
9	Klotho-derived peptide 6 ameliorates diabetic kidney disease by targeting Wnt/ β -catenin signaling. <i>Kidney International</i> , 2022, 102, 506-520.	2.6	26
10	B7-1 mediates podocyte injury and glomerulosclerosis through communication with Hsp90ab1-LRP5- β -catenin pathway. <i>Cell Death and Differentiation</i> , 2022, 29, 2399-2416.	5.0	7
11	The fibrogenic niche in kidney fibrosis: components and mechanisms. <i>Nature Reviews Nephrology</i> , 2022, 18, 545-557.	4.1	89
12	Cannabinoid receptor type 2 promotes kidney fibrosis through orchestrating β -catenin signaling. <i>Kidney International</i> , 2021, 99, 364-381.	2.6	32
13	Fibrillin-1-enriched microenvironment drives endothelial injury and vascular rarefaction in chronic kidney disease. <i>Science Advances</i> , 2021, 7, .	4.7	25
14	Matrix metalloproteinase-10 protects against acute kidney injury by augmenting epidermal growth factor receptor signaling. <i>Cell Death and Disease</i> , 2021, 12, 70.	2.7	10
15	LRP5 and LRP6 in Wnt Signaling: Similarity and Divergence. <i>Frontiers in Cell and Developmental Biology</i> , 2021, 9, 670960.	1.8	61
16	MicroRNA-10 negatively regulates inflammation in diabetic kidney via targeting activation of the NLRP3 inflammasome. <i>Molecular Therapy</i> , 2021, 29, 2308-2320.	3.7	35
17	Cannabinoid receptor 2 plays a central role in renal tubular mitochondrial dysfunction and kidney ageing. <i>Journal of Cellular and Molecular Medicine</i> , 2021, 25, 8957-8972.	1.6	14
18	The hepatocyte growth factor/c-met pathway is a key determinant of the fibrotic kidney local microenvironment. <i>IScience</i> , 2021, 24, 103112.	1.9	5

#	ARTICLE	IF	CITATIONS
19	Identification of matrix metalloproteinase-10 as a key mediator of podocyte injury and proteinuria. <i>Kidney International</i> , 2021, 100, 837-849.	2.6	15
20	Exogenous Wnt1 Prevents Acute Kidney Injury and Its Subsequent Progression to Chronic Kidney Disease. <i>Frontiers in Physiology</i> , 2021, 12, 745816.	1.3	3
21	Intensity of Macrophage Infiltration in Glomeruli Predicts Response to Immunosuppressive Therapy in Patients with IgA Nephropathy. <i>Journal of the American Society of Nephrology: JASN</i> , 2021, 32, 3187-3196.	3.0	28
22	Role of miRNA-671-5p in Mediating Wnt/ β 2-Catenin-Triggered Podocyte Injury. <i>Frontiers in Pharmacology</i> , 2021, 12, 784489.	1.6	7
23	Urinary Matrix Metalloproteinase 7 and Prediction of IgA Nephropathy Progression. <i>American Journal of Kidney Diseases</i> , 2020, 75, 384-393.	2.1	29
24	Tubule-derived exosomes play a central role in fibroblast activation and kidney fibrosis. <i>Kidney International</i> , 2020, 97, 1181-1195.	2.6	82
25	Molecular basis of kidney disease. , 2020, , 425-440.		0
26	MicroRNA-4660 mediates β 2-catenin-induced podocyte injury by targeting Wilms tumor 1. <i>FASEB Journal</i> , 2020, 34, 14424-14439.	0.2	8
27	Inhibition of Estrogen Sulfotransferase (SULT1E1/EST) Ameliorates Ischemic Acute Kidney Injury in Mice. <i>Journal of the American Society of Nephrology: JASN</i> , 2020, 31, 1496-1508.	3.0	12
28	CXCR4 motif chemokine receptor 4 aggravates renal fibrosis through activating JAK/STAT/GSK3 β / β 2-catenin pathway. <i>Journal of Cellular and Molecular Medicine</i> , 2020, 24, 3837-3855.	1.6	30
29	The Many Faces of Matrix Metalloproteinase-7 in Kidney Diseases. <i>Biomolecules</i> , 2020, 10, 960.	1.8	48
30	Sequential Wnt Agonist Then Antagonist Treatment Accelerates Tissue Repair and Minimizes Fibrosis. <i>IScience</i> , 2020, 23, 101047.	1.9	9
31	Sympathetic Overactivity in CKD Disrupts Buffering of Neurotransmission by Endothelium-Derived Hyperpolarizing Factor and Enhances Vasoconstriction. <i>Journal of the American Society of Nephrology: JASN</i> , 2020, 31, 2312-2325.	3.0	7
32	Tenascin-C promotes acute kidney injury to chronic kidney disease progression by impairing tubular integrity via α 2 β 1 integrin signaling. <i>Kidney International</i> , 2020, 97, 1017-1031.	2.6	41
33	Cellular Senescence in Kidney Fibrosis: Pathologic Significance and Therapeutic Strategies. <i>Frontiers in Pharmacology</i> , 2020, 11, 601325.	1.6	40
34	Myofibroblast in Kidney Fibrosis: Origin, Activation, and Regulation. <i>Advances in Experimental Medicine and Biology</i> , 2019, 1165, 253-283.	0.8	118
35	Sonic hedgehog selectively promotes lymphangiogenesis after kidney injury through noncanonical pathway. <i>American Journal of Physiology - Renal Physiology</i> , 2019, 317, F1022-F1033.	1.3	13
36	Wnt/ β 2-catenin/RAS signaling mediates age-related renal fibrosis and is associated with mitochondrial dysfunction. <i>Aging Cell</i> , 2019, 18, e13004.	3.0	155

#	ARTICLE	IF	CITATIONS
37	Early activation of fibroblasts is required for kidney repair and regeneration after injury. <i>FASEB Journal</i> , 2019, 33, 12576-12587.	0.2	27
38	Adipocytes initiate an adipose-cerebral-peripheral sympathetic reflex to induce insulin resistance during high-fat feeding. <i>Clinical Science</i> , 2019, 133, 1883-1899.	1.8	15
39	Wnt/ β -catenin regulates blood pressure and kidney injury in rats. <i>Biochimica Et Biophysica Acta - Molecular Basis of Disease</i> , 2019, 1865, 1313-1322.	1.8	29
40	A stimuli-responsive drug release nanoplatform for kidney-specific anti-fibrosis treatment. <i>Biomaterials Science</i> , 2019, 7, 1554-1564.	2.6	19
41	Matrix metalloproteinase-7 protects against acute kidney injury by priming renal tubules for survival and regeneration. <i>Kidney International</i> , 2019, 95, 1167-1180.	2.6	51
42	Wnt/ β -catenin links oxidative stress to podocyte injury and proteinuria. <i>Kidney International</i> , 2019, 95, 830-845.	2.6	105
43	Wnt/ β -catenin signaling mediates both heart and kidney injury in type 2 cardiorenal syndrome. <i>Kidney International</i> , 2019, 95, 815-829.	2.6	66
44	Tenascin-C protects against acute kidney injury by recruiting Wnt ligands. <i>Kidney International</i> , 2019, 95, 62-74.	2.6	34
45	Tubular injury triggers podocyte dysfunction by β -catenin-driven release of MMP-7. <i>JCI Insight</i> , 2019, 4, .	2.3	39
46	Sonic hedgehog connects podocyte injury to mesangial activation and glomerulosclerosis. <i>JCI Insight</i> , 2019, 4, .	2.3	14
47	A new model of diabetic nephropathy in C57BL/6 mice challenged with advanced oxidation protein products. <i>Free Radical Biology and Medicine</i> , 2018, 118, 71-84.	1.3	15
48	Wnt9a Promotes Renal Fibrosis by Accelerating Cellular Senescence in Tubular Epithelial Cells. <i>Journal of the American Society of Nephrology: JASN</i> , 2018, 29, 1238-1256.	3.0	163
49	Fibroblast-Specific β -Catenin Signaling Dictates the Outcome of AKI. <i>Journal of the American Society of Nephrology: JASN</i> , 2018, 29, 1257-1271.	3.0	55
50	Activation of Constitutive Androstane Receptor Ameliorates Renal Ischemia-Reperfusion-Induced Kidney and Liver Injury. <i>Molecular Pharmacology</i> , 2018, 93, 239-250.	1.0	14
51	A renal-cerebral-peripheral sympathetic reflex mediates insulin resistance in chronic kidney disease. <i>EBioMedicine</i> , 2018, 37, 281-293.	2.7	18
52	New insights into the role and mechanism of Wnt/ β -catenin signalling in kidney fibrosis. <i>Nephrology</i> , 2018, 23, 38-43.	0.7	69
53	Long noncoding RNA <i>lnc-TSI</i> inhibits renal fibrogenesis by negatively regulating the TGF- β /Smad3 pathway. <i>Science Translational Medicine</i> , 2018, 10, .	5.8	129
54	IL-17 Receptor Signaling Negatively Regulates the Development of Tubulointerstitial Fibrosis in the Kidney. <i>Mediators of Inflammation</i> , 2018, 2018, 1-14.	1.4	22

#	ARTICLE	IF	CITATIONS
55	Wnt Signaling in Kidney Development and Disease. <i>Progress in Molecular Biology and Translational Science</i> , 2018, 153, 181-207.	0.9	93
56	A New Criterion for Pediatric AKI Based on the Reference Change Value of Serum Creatinine. <i>Journal of the American Society of Nephrology: JASN</i> , 2018, 29, 2432-2442.	3.0	52
57	Targeted inhibition of the type 2 cannabinoid receptor is a novel approach to reduce renal fibrosis. <i>Kidney International</i> , 2018, 94, 756-772.	2.6	48
58	Molecular Basis of Kidney Disease. , 2018, , 531-553.		3
59	An essential role for Wnt/ β 2-catenin signaling in mediating hypertensive heart disease. <i>Scientific Reports</i> , 2018, 8, 8996.	1.6	68
60	(Pro)renin Receptor Is an Amplifier of Wnt/ β 2-Catenin Signaling in Kidney Injury and Fibrosis. <i>Journal of the American Society of Nephrology: JASN</i> , 2017, 28, 2393-2408.	3.0	86
61	Tubule-Derived Wnts Are Required for Fibroblast Activation and Kidney Fibrosis. <i>Journal of the American Society of Nephrology: JASN</i> , 2017, 28, 2322-2336.	3.0	95
62	Reno-Cerebral Reflex Activates the Renin-Angiotensin System, Promoting Oxidative Stress and Renal Damage After Ischemia-Reperfusion Injury. <i>Antioxidants and Redox Signaling</i> , 2017, 27, 415-432.	2.5	53
63	C-X-C Chemokine Receptor Type 4 Plays a Crucial Role in Mediating Oxidative Stress-Induced Podocyte Injury. <i>Antioxidants and Redox Signaling</i> , 2017, 27, 345-362.	2.5	37
64	Contrast-Enhanced Ultrasound for Assessing Renal Perfusion Impairment and Predicting Acute Kidney Injury to Chronic Kidney Disease Progression. <i>Antioxidants and Redox Signaling</i> , 2017, 27, 1397-1411.	2.5	40
65	Urinary Matrix Metalloproteinase-7 Predicts Severe AKI and Poor Outcomes after Cardiac Surgery. <i>Journal of the American Society of Nephrology: JASN</i> , 2017, 28, 3373-3382.	3.0	52
66	Tenascin-C Is a Major Component of the Fibrogenic Niche in Kidney Fibrosis. <i>Journal of the American Society of Nephrology: JASN</i> , 2017, 28, 785-801.	3.0	87
67	Matrix Metalloproteinase-7 Is a Urinary Biomarker and Pathogenic Mediator of Kidney Fibrosis. <i>Journal of the American Society of Nephrology: JASN</i> , 2017, 28, 598-611.	3.0	118
68	Numb contributes to renal fibrosis by promoting tubular epithelial cell cycle arrest at G2/M. <i>Oncotarget</i> , 2016, 7, 25604-25619.	0.8	21
69	Sonic hedgehog signaling in kidney fibrosis: a master communicator. <i>Science China Life Sciences</i> , 2016, 59, 920-929.	2.3	43
70	Keap1 hypomorphism protects against ischemic and obstructive kidney disease. <i>Scientific Reports</i> , 2016, 6, 36185.	1.6	32
71	Signaling Crosstalk between Tubular Epithelial Cells and Interstitial Fibroblasts after Kidney Injury. <i>Kidney Diseases (Basel, Switzerland)</i> , 2016, 2, 136-144.	1.2	90
72	Wnt/ β 2-catenin signaling and renin-angiotensin system in chronic kidney disease. <i>Current Opinion in Nephrology and Hypertension</i> , 2016, 25, 100-106.	1.0	61

#	ARTICLE	IF	CITATIONS
73	Therapy for kidney fibrosis: is the Src kinase a potential target?. <i>Kidney International</i> , 2016, 89, 12-14.	2.6	18
74	Understanding the mechanisms of kidney fibrosis. <i>Nature Reviews Nephrology</i> , 2016, 12, 68-70.	4.1	156
75	Wnt/ β -catenin signaling in kidney injury and repair: a double-edged sword. <i>Laboratory Investigation</i> , 2016, 96, 156-167.	1.7	146
76	Sustained Activation of Wnt/ β -Catenin Signaling Drives AKI to CKD Progression. <i>Journal of the American Society of Nephrology: JASN</i> , 2016, 27, 1727-1740.	3.0	189
77	Wnt/ β -catenin signalling and podocyte dysfunction in proteinuric kidney disease. <i>Nature Reviews Nephrology</i> , 2015, 11, 535-545.	4.1	167
78	Klotho Ameliorates Kidney Injury and Fibrosis and Normalizes Blood Pressure by Targeting the Renin-Angiotensin System. <i>American Journal of Pathology</i> , 2015, 185, 3211-3223.	1.9	124
79	Extracellular Superoxide Dismutase Protects against Proteinuric Kidney Disease. <i>Journal of the American Society of Nephrology: JASN</i> , 2015, 26, 2447-2459.	3.0	54
80	Loss of Klotho in CKD Breaks One's Heart. <i>Journal of the American Society of Nephrology: JASN</i> , 2015, 26, 2305-2307.	3.0	11
81	Oestrogen sulfotransferase ablation sensitizes mice to sepsis. <i>Nature Communications</i> , 2015, 6, 7979.	5.8	33
82	Mutual Antagonism of Wilms' Tumor 1 and β -Catenin Dictates Podocyte Health and Disease. <i>Journal of the American Society of Nephrology: JASN</i> , 2015, 26, 677-691.	3.0	55
83	Multiple Genes of the Renin-Angiotensin System Are Novel Targets of Wnt/ β -Catenin Signaling. <i>Journal of the American Society of Nephrology: JASN</i> , 2015, 26, 107-120.	3.0	184
84	Wnt/ β -catenin signaling and kidney fibrosis. <i>Kidney International Supplements</i> , 2014, 4, 84-90.	4.6	221
85	Renal expression of advanced oxidative protein products predicts progression of renal fibrosis in patients with IgA nephropathy. <i>Laboratory Investigation</i> , 2014, 94, 966-977.	1.7	13
86	New insights into the pathogenesis and therapeutics of kidney fibrosis. <i>Kidney International Supplements</i> , 2014, 4, 1.	4.6	8
87	Klotho suppresses renal tubulointerstitial fibrosis by controlling basic fibroblast growth factor signaling. <i>Journal of Pathology</i> , 2014, 234, 560-572.	2.1	77
88	Arrestin(g) Podocyte Injury with Endothelin Antagonism. <i>Journal of the American Society of Nephrology: JASN</i> , 2014, 25, 423-425.	3.0	4
89	RANK- and c-Met-mediated signal network promotes prostate cancer metastatic colonization. <i>Endocrine-Related Cancer</i> , 2014, 21, 311-326.	1.6	74
90	Complement Component C5a Permits the Coexistence of Pathogenic Th17 Cells and Type I IFN in Lupus. <i>Journal of Immunology</i> , 2014, 193, 3288-3295.	0.4	21

#	ARTICLE	IF	CITATIONS
91	Sonic Hedgehog Is a Novel Tubule-Derived Growth Factor for Interstitial Fibroblasts after Kidney Injury. <i>Journal of the American Society of Nephrology: JASN</i> , 2014, 25, 2187-2200.	3.0	116
92	Loss of Klotho Contributes to Kidney Injury by Derepression of Wnt/ β -Catenin Signaling. <i>Journal of the American Society of Nephrology: JASN</i> , 2013, 24, 771-785.	3.0	309
93	Activation of hepatocyte growth factor receptor, c-met, in renal tubules is required for renoprotection after acute kidney injury. <i>Kidney International</i> , 2013, 84, 509-520.	2.6	108
94	Kindlin-2 Mediates Activation of TGF- β /Smad Signaling and Renal Fibrosis. <i>Journal of the American Society of Nephrology: JASN</i> , 2013, 24, 1387-1398.	3.0	83
95	Macrophage-derived TGF- β in renal fibrosis: not a macro-impact after all. <i>American Journal of Physiology - Renal Physiology</i> , 2013, 305, F821-F822.	1.3	7
96	Fibrosis and anaemia in CKD—two beasts, one ancestor. <i>Nature Reviews Nephrology</i> , 2013, 9, 563-565.	4.1	5
97	Kidney tubular β -catenin signaling controls interstitial fibroblast fate via epithelial-mesenchymal communication. <i>Scientific Reports</i> , 2013, 3, 1878.	1.6	64
98	Endothelin Receptor A Blockade Is an Ineffective Treatment for Adriamycin Nephropathy. <i>PLoS ONE</i> , 2013, 8, e79963.	1.1	13
99	AGE-LDL Activates Toll Like Receptor 4 Pathway and Promotes Inflammatory Cytokines Production in Renal Tubular Epithelial Cells. <i>International Journal of Biological Sciences</i> , 2013, 9, 94-107.	2.6	36
100	Sonic Hedgehog Signaling Mediates Epithelial-Mesenchymal Communication and Promotes Renal Fibrosis. <i>Journal of the American Society of Nephrology: JASN</i> , 2012, 23, 801-813.	3.0	166
101	MiR-382 targeting of kallikrein 5 contributes to renal inner medullary interstitial fibrosis. <i>Physiological Genomics</i> , 2012, 44, 259-267.	1.0	71
102	Tubule-specific ablation of endogenous β -catenin aggravates acute kidney injury in mice. <i>Kidney International</i> , 2012, 82, 537-547.	2.6	181
103	The receptor of advanced glycation end products plays a central role in advanced oxidation protein products-induced podocyte apoptosis. <i>Kidney International</i> , 2012, 82, 759-770.	2.6	104
104	Matrix Metalloproteinase-7 as a Surrogate Marker Predicts Renal Wnt/ β -Catenin Activity in CKD. <i>Journal of the American Society of Nephrology: JASN</i> , 2012, 23, 294-304.	3.0	131
105	Matrix metalloproteinases in kidney homeostasis and diseases. <i>American Journal of Physiology - Renal Physiology</i> , 2012, 302, F1351-F1361.	1.3	204
106	Loss of vitamin D receptor in chronic kidney disease: a potential mechanism linking inflammation to epithelial-to-mesenchymal transition. <i>American Journal of Physiology - Renal Physiology</i> , 2012, 303, F1107-F1115.	1.3	50
107	Tubular cell dedifferentiation and peritubular inflammation are coupled by the transcription regulator Id1 in renal fibrogenesis. <i>Kidney International</i> , 2012, 81, 880-891.	2.6	24
108	Cellular and molecular mechanisms of renal fibrosis. <i>Nature Reviews Nephrology</i> , 2011, 7, 684-696.	4.1	1,067

#	ARTICLE	IF	CITATIONS
109	PINCH1 Is Transcriptional Regulator in Podocytes That Interacts with WT1 and Represses Podocalyxin Expression. PLoS ONE, 2011, 6, e17048.	1.1	20
110	Blockade of Wnt/ β -Catenin Signaling by Paricalcitol Ameliorates Proteinuria and Kidney Injury. Journal of the American Society of Nephrology: JASN, 2011, 22, 90-103.	3.0	242
111	Targeted Inhibition of β -Catenin/CBP Signaling Ameliorates Renal Interstitial Fibrosis. Journal of the American Society of Nephrology: JASN, 2011, 22, 1642-1653.	3.0	210
112	Albumin overload activates intrarenal renin-angiotensin system through protein kinase C and NADPH oxidase-dependent pathway. Journal of Hypertension, 2011, 29, 1411-1421.	0.3	54
113	Inhibition of Proinflammatory RANTES Expression by TGF- β 1 Is Mediated by Glycogen Synthase Kinase-3-dependent β -Catenin Signaling. Journal of Biological Chemistry, 2011, 286, 7052-7059.	1.6	28
114	Canonical Wnt/ β -catenin signaling mediates transforming growth factor- β 1-driven podocyte injury and proteinuria. Kidney International, 2011, 80, 1159-1169.	2.6	131
115	Plasminogen Activator Inhibitor-1 Is a Transcriptional Target of the Canonical Pathway of Wnt/ β -Catenin Signaling. Journal of Biological Chemistry, 2010, 285, 24665-24675.	1.6	97
116	New Insights into Epithelial-Mesenchymal Transition in Kidney Fibrosis. Journal of the American Society of Nephrology: JASN, 2010, 21, 212-222.	3.0	753
117	Opposite Action of Peroxisome Proliferator-activated Receptor- β 3 in Regulating Renal Inflammation. Journal of Biological Chemistry, 2010, 285, 29981-29988.	1.6	27
118	Hepatocyte growth factor signaling ameliorates podocyte injury and proteinuria. Kidney International, 2010, 77, 962-973.	2.6	87
119	Inhibition of integrin-linked kinase blocks podocyte epithelial-mesenchymal transition and ameliorates proteinuria. Kidney International, 2010, 78, 363-373.	2.6	134
120	tPA Is a Potent Mitogen for Renal Interstitial Fibroblasts. American Journal of Pathology, 2010, 177, 1164-1175.	1.9	22
121	Wnt/ β -Catenin Signaling Promotes Renal Interstitial Fibrosis. Journal of the American Society of Nephrology: JASN, 2009, 20, 765-776.	3.0	510
122	Inhibition of Integrin-Linked Kinase Attenuates Renal Interstitial Fibrosis. Journal of the American Society of Nephrology: JASN, 2009, 20, 1907-1918.	3.0	108
123	Advanced oxidation protein products: a causative link between oxidative stress and podocyte depletion. Kidney International, 2009, 76, 1125-1127.	2.6	18
124	Wnt/ β -Catenin Signaling Promotes Podocyte Dysfunction and Albuminuria. Journal of the American Society of Nephrology: JASN, 2009, 20, 1997-2008.	3.0	356
125	Combination therapy with paricalcitol and trandolapril reduces renal fibrosis in obstructive nephropathy. Kidney International, 2009, 76, 1248-1257.	2.6	65
126	A role of Wnt/ β -catenin signaling in the pathogenesis of renal interstitial fibrosis. FASEB Journal, 2009, 23, 359.3.	0.2	0

#	ARTICLE	IF	CITATIONS
127	Epithelial-to-Mesenchymal Transition Is a Potential Pathway Leading to Podocyte Dysfunction and Proteinuria. <i>American Journal of Pathology</i> , 2008, 172, 299-308.	1.9	300
128	Hepatocyte Growth Factor Exerts Its Anti-Inflammatory Action by Disrupting Nuclear Factor- κ B Signaling. <i>American Journal of Pathology</i> , 2008, 173, 30-41.	1.9	111
129	tPA Protects Renal Interstitial Fibroblasts and Myofibroblasts from Apoptosis. <i>Journal of the American Society of Nephrology: JASN</i> , 2008, 19, 503-514.	3.0	64
130	Smad ubiquitination regulatory factor-2 in the fibrotic kidney: regulation, target specificity, and functional implication. <i>American Journal of Physiology - Renal Physiology</i> , 2008, 294, F1076-F1083.	1.3	68
131	Paricalcitol Inhibits Renal Inflammation by Promoting Vitamin D Receptor-mediated Sequestration of NF- κ B Signaling. <i>Journal of the American Society of Nephrology: JASN</i> , 2008, 19, 1741-1752.	3.0	238
132	Animal Models of Kidney Diseases. , 2008, , 657-664.		7
133	Novel actions of tissue-type plasminogen activator in chronic kidney disease. <i>Frontiers in Bioscience - Landmark</i> , 2008, Volume, 5174.	3.0	35
134	Role of Bcl-xL induction in HGF-mediated renal epithelial cell survival after oxidant stress. <i>International Journal of Clinical and Experimental Pathology</i> , 2008, 1, 242-53.	0.5	19
135	Cell Phenotype-specific Down-regulation of Smad3 Involves Decreased Gene Activation as Well as Protein Degradation. <i>Journal of Biological Chemistry</i> , 2007, 282, 15534-15540.	1.6	43
136	Molecular Basis for the Cell Type-specific Induction of SnoN Expression by Hepatocyte Growth Factor. <i>Journal of the American Society of Nephrology: JASN</i> , 2007, 18, 2340-2349.	3.0	31
137	Tubular Epithelial Cell Dedifferentiation Is Driven by the Helix-Loop-Helix Transcriptional Inhibitor Id1. <i>Journal of the American Society of Nephrology: JASN</i> , 2007, 18, 449-460.	3.0	80
138	PINCH-1 Promotes Tubular Epithelial-to-Mesenchymal Transition by Interacting with Integrin-Linked Kinase. <i>Journal of the American Society of Nephrology: JASN</i> , 2007, 18, 2534-2543.	3.0	58
139	Therapeutic role and potential mechanisms of active Vitamin D in renal interstitial fibrosis. <i>Journal of Steroid Biochemistry and Molecular Biology</i> , 2007, 103, 491-496.	1.2	62
140	Tissue-type plasminogen activator promotes murine myofibroblast activation through LDL receptor-related protein 1-mediated integrin signaling. <i>Journal of Clinical Investigation</i> , 2007, 117, 3821-32.	3.9	91
141	Hepatocyte Growth Factor Attenuates Liver Fibrosis Induced by Bile Duct Ligation. <i>American Journal of Pathology</i> , 2006, 168, 1500-1512.	1.9	186
142	Renal fibrosis: New insights into the pathogenesis and therapeutics. <i>Kidney International</i> , 2006, 69, 213-217.	2.6	909
143	Rapamycin and chronic kidney disease: beyond the inhibition of inflammation. <i>Kidney International</i> , 2006, 69, 1925-1927.	2.6	23
144	Hepatocyte growth factor: New arsenal in the fights against renal fibrosis?. <i>Kidney International</i> , 2006, 70, 238-240.	2.6	59

#	ARTICLE	IF	CITATIONS
145	Potential role of active vitamin D in retarding the progression of chronic kidney disease. <i>Nephrology Dialysis Transplantation</i> , 2006, 22, 321-328.	0.4	60
146	Distinctive role of Stat3 and Erk-1/2 activation in mediating interferon- β inhibition of TGF- β 1 action. <i>American Journal of Physiology - Renal Physiology</i> , 2006, 290, F1234-F1240.	1.3	17
147	Downregulation of SnoN Expression in Obstructive Nephropathy Is Mediated by an Enhanced Ubiquitin-Dependent Degradation. <i>Journal of the American Society of Nephrology: JASN</i> , 2006, 17, 2781-2791.	3.0	63
148	Tissue-type Plasminogen Activator Acts as a Cytokine That Triggers Intracellular Signal Transduction and Induces Matrix Metalloproteinase-9 Gene Expression. <i>Journal of Biological Chemistry</i> , 2006, 281, 2120-2127.	1.6	177
149	Essential Role of Integrin-Linked Kinase in Podocyte Biology: Bridging the Integrin and Slit Diaphragm Signaling. <i>Journal of the American Society of Nephrology: JASN</i> , 2006, 17, 2164-2175.	3.0	123
150	Paricalcitol Attenuates Renal Interstitial Fibrosis in Obstructive Nephropathy. <i>Journal of the American Society of Nephrology: JASN</i> , 2006, 17, 3382-3393.	3.0	250
151	Hepatocyte Growth Factor Is a Downstream Effector that Mediates the Antifibrotic Action of Peroxisome Proliferator-Activated Receptor- β Agonists. <i>Journal of the American Society of Nephrology: JASN</i> , 2006, 17, 54-65.	3.0	129
152	β -Catenin: A Principal In Vivo Mediator of HGF-induced Liver Growth. <i>FASEB Journal</i> , 2006, 20, A1079.	0.2	0
153	1,25-dihydroxyvitamin D3 inhibits renal interstitial myofibroblast activation by inducing hepatocyte growth factor expression. <i>Kidney International</i> , 2005, 68, 1500-1510.	2.6	87
154	Both Sp1 and Smad participate in mediating TGF- β 1-induced HGF receptor expression in renal epithelial cells. <i>American Journal of Physiology - Renal Physiology</i> , 2005, 288, F16-F26.	1.3	52
155	A Novel Mechanism by which Hepatocyte Growth Factor Blocks Tubular Epithelial to Mesenchymal Transition. <i>Journal of the American Society of Nephrology: JASN</i> , 2005, 16, 68-78.	3.0	169
156	Hepatocyte Growth Factor Receptor Signaling Mediates the Anti-Fibrotic Action of 9-cis-Retinoic Acid in Glomerular Mesangial Cells. <i>American Journal of Pathology</i> , 2005, 167, 947-957.	1.9	42
157	β -Cell-Specific Ablation of the Hepatocyte Growth Factor Receptor Results in Reduced Islet Size, Impaired Insulin Secretion, and Glucose Intolerance. <i>American Journal of Pathology</i> , 2005, 167, 429-436.	1.9	70
158	Epithelial to Mesenchymal Transition in Renal Fibrogenesis: Pathologic Significance, Molecular Mechanism, and Therapeutic Intervention. <i>Journal of the American Society of Nephrology: JASN</i> , 2004, 15, 1-12.	3.0	1,005
159	Hepatocyte Growth Factor Antagonizes the Profibrotic Action of TGF- β 1 in Mesangial Cells by Stabilizing Smad Transcriptional Corepressor TGIF. <i>Journal of the American Society of Nephrology: JASN</i> , 2004, 15, 1402-1412.	3.0	134
160	Intravenous Administration of Hepatocyte Growth Factor Gene Ameliorates Diabetic Nephropathy in Mice. <i>Journal of the American Society of Nephrology: JASN</i> , 2004, 15, 2637-2647.	3.0	108
161	Hepatocyte growth factor in kidney fibrosis: therapeutic potential and mechanisms of action. <i>American Journal of Physiology - Renal Physiology</i> , 2004, 287, F7-F16.	1.3	234
162	Hepatocyte Growth Factor Suppresses Renal Interstitial Myofibroblast Activation and Intercepts Smad Signal Transduction. <i>American Journal of Pathology</i> , 2003, 163, 621-632.	1.9	142

#	ARTICLE	IF	CITATIONS
163	Transforming Growth Factor- α 1 Potentiates Renal Tubular Epithelial Cell Death by a Mechanism Independent of Smad Signaling. <i>Journal of Biological Chemistry</i> , 2003, 278, 12537-12545.	1.6	135
164	Downregulation of Smad Transcriptional Corepressors SnoN and Ski in the Fibrotic Kidney: An Amplification Mechanism for TGF- β 1 Signaling. <i>Journal of the American Society of Nephrology: JASN</i> , 2003, 14, 3167-3177.	3.0	85
165	Urokinase Receptor Deficiency Accelerates Renal Fibrosis in Obstructive Nephropathy. <i>Journal of the American Society of Nephrology: JASN</i> , 2003, 14, 1254-1271.	3.0	111
166	Hepatocyte Growth Factor Preserves Beta Cell Mass and Mitigates Hyperglycemia in Streptozotocin-induced Diabetic Mice. <i>Journal of Biological Chemistry</i> , 2003, 278, 27080-27087.	1.6	74
167	Suppression of HGF receptor gene expression by oxidative stress is mediated through the interplay between Sp1 and Egr-1. <i>American Journal of Physiology - Renal Physiology</i> , 2003, 284, F1216-F1225.	1.3	44
168	Sp1 and Sp3 transcription factors synergistically regulate HGF receptor gene expression in kidney. <i>American Journal of Physiology - Renal Physiology</i> , 2003, 284, F82-F94.	1.3	64
169	Delayed administration of hepatocyte growth factor reduces renal fibrosis in obstructive nephropathy. <i>American Journal of Physiology - Renal Physiology</i> , 2003, 284, F349-F357.	1.3	77
170	Role for integrin-linked kinase in mediating tubular epithelial to mesenchymal transition and renal interstitial fibrogenesis. <i>Journal of Clinical Investigation</i> , 2003, 112, 503-516.	3.9	155
171	Role for integrin-linked kinase in mediating tubular epithelial to mesenchymal transition and renal interstitial fibrogenesis. <i>Journal of Clinical Investigation</i> , 2003, 112, 503-516.	3.9	314
172	Hepatocyte Growth Factor Gene Therapy and Angiotensin II Blockade Synergistically Attenuate Renal Interstitial Fibrosis in Mice. <i>Journal of the American Society of Nephrology: JASN</i> , 2002, 13, 2464-2477.	3.0	145
173	Hepatocyte growth factor and the kidney. <i>Current Opinion in Nephrology and Hypertension</i> , 2002, 11, 23-30.	1.0	108
174	Disruption of tissue-type plasminogen activator gene in mice reduces renal interstitial fibrosis in obstructive nephropathy. <i>Journal of Clinical Investigation</i> , 2002, 110, 1525-1538.	3.9	180
175	Disruption of tissue-type plasminogen activator gene in mice reduces renal interstitial fibrosis in obstructive nephropathy. <i>Journal of Clinical Investigation</i> , 2002, 110, 1525-1538.	3.9	137
176	Blockage of Tubular Epithelial to Myofibroblast Transition by Hepatocyte Growth Factor Prevents Renal Interstitial Fibrosis. <i>Journal of the American Society of Nephrology: JASN</i> , 2002, 13, 96-107.	3.0	295
177	Single Injection of Naked Plasmid Encoding Hepatocyte Growth Factor Prevents Cell Death and Ameliorates Acute Renal Failure in Mice. <i>Journal of the American Society of Nephrology: JASN</i> , 2002, 13, 411-422.	3.0	107
178	Dissection of Key Events in Tubular Epithelial to Myofibroblast Transition and Its Implications in Renal Interstitial Fibrosis. <i>American Journal of Pathology</i> , 2001, 159, 1465-1475.	1.9	773
179	Sustained expression of naked plasmid DNA encoding hepatocyte growth factor in mice promotes liver and overall body growth. <i>Hepatology</i> , 2001, 33, 848-859.	3.6	92
180	Endogenous hepatocyte growth factor ameliorates chronic renal injury by activating matrix degradation pathways. <i>Kidney International</i> , 2000, 58, 2028-2043.	2.6	124

#	ARTICLE	IF	CITATIONS
181	Hepatocyte growth factor promotes renal epithelial cell survival by dual mechanisms. American Journal of Physiology - Renal Physiology, 1999, 277, F624-F633.	1.3	70
182	Up-regulation of hepatocyte growth factor receptor: An amplification and targeting mechanism for hepatocyte growth factor action in acute renal failure. Kidney International, 1999, 55, 442-453.	2.6	113
183	The human hepatocyte growth factor receptor gene: complete structural organization and promoter characterization. Gene, 1998, 215, 159-169.	1.0	90
184	Hepatocyte Growth Factor Protects Renal Epithelial Cells from Apoptotic Cell Death. Biochemical and Biophysical Research Communications, 1998, 246, 821-826.	1.0	85
185	Constitutive Expression of HGF Modulates Renal Epithelial Cell Phenotype and Induces c-met and Fibronectin Expression. Experimental Cell Research, 1998, 242, 174-185.	1.2	35
186	The Upstream Regulatory Regions of the Hepatocyte Growth Factor Gene Promoter Are Essential for Its Expression in Transgenic Mice. Journal of Biological Chemistry, 1998, 273, 6900-6908.	1.6	16
187	Modulation of hepatocyte growth factor gene expression by estrogen in mouse ovary. Molecular and Cellular Endocrinology, 1994, 104, 173-181.	1.6	67