

Kai Cai

List of Publications by Year in Descending Order

Source: <https://exaly.com/author-pdf/4680010/kai-cai-publications-by-year.pdf>

Version: 2024-04-27

This document has been generated based on the publications and citations recorded by exaly.com. For the latest version of this publication list, visit the link given above.

The third column is the impact factor (IF) of the journal, and the fourth column is the number of citations of the article.

22
papers

675
citations

15
h-index

25
g-index

28
ext. papers

881
ext. citations

6.6
avg, IF

3.9
L-index

#	Paper	IF	Citations
22	Structural organization of the C1b projection within the ciliary central apparatus. <i>Journal of Cell Science</i> , 2021 , 134,	5.3	2
21	ISCU interacts with NFU1, and ISCU[4Fe-4S] transfers its Fe-S cluster to NFU1 leading to the production of holo-NFU1. <i>Journal of Structural Biology</i> , 2020 , 210, 107491	3.4	10
20	Mitochondrial metabolism promotes adaptation to proteotoxic stress. <i>Nature Chemical Biology</i> , 2019 , 15, 681-689	11.7	62
19	Conformational flexibility in the enterovirus RNA replication platform. <i>Rna</i> , 2019 , 25, 376-387	5.8	5
18	ISCU(M108I) and ISCU(D39V) Differ from Wild-Type ISCU in Their Failure To Form Cysteine Desulfurase Complexes Containing Both Frataxin and Ferredoxin. <i>Biochemistry</i> , 2018 , 57, 1491-1500	3.2	12
17	Interactions of iron-bound frataxin with ISCU and ferredoxin on the cysteine desulfurase complex leading to Fe-S cluster assembly. <i>Journal of Inorganic Biochemistry</i> , 2018 , 183, 107-116	4.2	35
16	Architectural Features of Human Mitochondrial Cysteine Desulfurase Complexes from Crosslinking Mass Spectrometry and Small-Angle X-Ray Scattering. <i>Structure</i> , 2018 , 26, 1127-1136.e4	5.2	15
15	NMR as a Tool to Investigate the Processes of Mitochondrial and Cytosolic Iron-Sulfur Cluster Biosynthesis. <i>Molecules</i> , 2018 , 23,	4.8	5
14	Mitochondrial Cysteine Desulfurase and ISD11 Coexpressed in Escherichia coli Yield Complex Containing Acyl Carrier Protein. <i>ACS Chemical Biology</i> , 2017 , 12, 918-921	4.9	23
13	Human Mitochondrial Ferredoxin 1 (FDX1) and Ferredoxin 2 (FDX2) Both Bind Cysteine Desulfurase and Donate Electrons for Iron-Sulfur Cluster Biosynthesis. <i>Biochemistry</i> , 2017 , 56, 487-499	3.2	46
12	Structural/Functional Properties of Human NFU1, an Intermediate [4Fe-4S] Carrier in Human Mitochondrial Iron-Sulfur Cluster Biogenesis. <i>Structure</i> , 2016 , 24, 2080-2091	5.2	37
11	The specialized Hsp70 (HscA) interdomain linker binds to its nucleotide-binding domain and stimulates ATP hydrolysis in both cis and trans configurations. <i>Biochemistry</i> , 2014 , 53, 7148-59	3.2	20
10	Metamorphic protein IscU alternates conformations in the course of its role as the scaffold protein for iron-sulfur cluster biosynthesis and delivery. <i>FEBS Letters</i> , 2013 , 587, 1172-9	3.8	57
9	Human mitochondrial chaperone (mtHSP70) and cysteine desulfurase (NFS1) bind preferentially to the disordered conformation, whereas co-chaperone (HSC20) binds to the structured conformation of the iron-sulfur cluster scaffold protein (ISCU). <i>Journal of Biological Chemistry</i> , 2013 , 288, 28755-70	5.4	44
8	Electron transfer mechanism of the Rieske protein from <i>Thermus thermophilus</i> from solution nuclear magnetic resonance investigations. <i>Biochemistry</i> , 2013 , 52, 2862-73	3.2	5
7	Cadmium(II) complexes with 3,5-di(1H-imidazol-1-yl)benzoate: topological and structural diversity tuned by counteranions. <i>CrystEngComm</i> , 2010 , 12, 100-108	3.3	65
6	Synthesis, Structures, and Properties of Zinc(II) and Cadmium(II) Complexes with 1,2,4,5-Tetrakis(imidazol-1-ylmethyl)benzene and Multicarboxylate Ligands. <i>Crystal Growth and Design</i> , 2010 , 10, 2553-2562	3.5	75

- | | | | |
|---|---|-----|----|
| 5 | Copper(II) and zinc(II) complexes with macrocyclic ligand: Structure variation via counteranion and co-ligand. <i>Journal of Molecular Structure</i> , 2010 , 973, 104-115 | 3.4 | 7 |
| 4 | Imidazolate-bridged dinuclear copper(II) complex with new macrocyclic ligand bearing two 1H-imidazol-4-yl-pendants. <i>Inorganic Chemistry Communication</i> , 2010 , 13, 847-851 | 3.1 | 16 |
| 3 | Zinc(II) Complexes with 1H-Imidazol-4-yl-Containing Polyamine Ligand. <i>Zeitschrift Fur Anorganische Und Allgemeine Chemie</i> , 2010 , 636, 2009-2015 | 1.3 | 4 |
| 2 | Imidazolate-bridged dicopper(II) and copper(II)-zinc(II) complexes of macrocyclic ligand with methylimidazol pendants: Model study of copper(II)-zinc(II) superoxide dismutase. <i>Journal of Inorganic Biochemistry</i> , 2009 , 103, 1156-61 | 4.2 | 21 |
| 1 | pH-dependent self-assembly of copper(II) complexes with a new imidazole-containing polyamine ligand: Synthesis, structure and magnetic property. <i>Polyhedron</i> , 2008 , 27, 2672-2680 | 2.7 | 27 |