

Kazuhiro Komura

List of Publications by Year in descending order

Source: <https://exaly.com/author-pdf/4681968/publications.pdf>

Version: 2024-02-01

72
papers

4,743
citations

94269

37
h-index

95083

68
g-index

74
all docs

74
docs citations

74
times ranked

5088
citing authors

#	ARTICLE	IF	CITATIONS
1	Augmented interferon I signaling in a patient with COVID toes. <i>Journal of Dermatology</i> , 2021, 48, e242-e243.	0.6	0
2	A synthetic PPAR- β agonist triterpenoid ameliorates experimental fibrosis: PPAR- β -independent suppression of fibrotic responses. <i>Annals of the Rheumatic Diseases</i> , 2014, 73, 446-454.	0.5	62
3	Common and Distinct Clinical Features in Adult Patients with Anti-Aminoacyl-tRNA Synthetase Antibodies: Heterogeneity within the Syndrome. <i>PLoS ONE</i> , 2013, 8, e60442.	1.1	306
4	The roles of P- and E-selectins and P-selectin glycoprotein ligand-1 in primary and metastatic mouse melanomas. <i>Journal of Dermatological Science</i> , 2011, 64, 99-107.	1.0	9
5	Canonical Wnt signaling induces skin fibrosis and subcutaneous lipoatrophy: A novel mouse model for scleroderma?. <i>Arthritis and Rheumatism</i> , 2011, 63, 1707-1717.	6.7	178
6	Clinical Correlations With Dermatomyositis-Specific Autoantibodies in Adult Japanese Patients With Dermatomyositis. <i>Archives of Dermatology</i> , 2011, 147, 391.	1.7	293
7	Elevated serum interleukin-27 levels in patients with systemic sclerosis: association with T cell, B cell and fibroblast activation. <i>Annals of the Rheumatic Diseases</i> , 2011, 70, 194-200.	0.5	54
8	Autoantibody against caspase-3, an executioner of apoptosis, in patients with systemic sclerosis. <i>Rheumatology International</i> , 2010, 30, 871-878.	1.5	10
9	Autoantibody against one of the antioxidant repair enzymes, methionine sulfoxide reductase A, in systemic sclerosis: association with pulmonary fibrosis and vascular damage. <i>Archives of Dermatological Research</i> , 2010, 302, 27-35.	1.1	33
10	Treatment with rapamycin prevents fibrosis in tight skin and bleomycin-induced mouse models of systemic sclerosis. <i>Arthritis and Rheumatism</i> , 2010, 62, 2476-2487.	6.7	118
11	PPAR- β Downregulation by TGF- β in Fibroblast and Impaired Expression and Function in Systemic Sclerosis: A Novel Mechanism for Progressive Fibrogenesis. <i>PLoS ONE</i> , 2010, 5, e13778.	1.1	158
12	Elevated Serum Concentrations of Triggering Receptor Expressed on Myeloid Cells-1 in Diffuse Cutaneous Systemic Sclerosis: Association with Severity of Pulmonary Fibrosis. <i>Journal of Rheumatology</i> , 2010, 37, 787-791.	1.0	6
13	Usefulness of anti-cyclic citrullinated peptide antibody and rheumatoid factor to detect rheumatoid arthritis in patients with systemic sclerosis. <i>Rheumatology</i> , 2010, 49, 2135-2139.	0.9	32
14	Cell Adhesion Molecules Regulate Fibrotic Process via Th1/Th2/Th17 Cell Balance in a Bleomycin-Induced Scleroderma Model. <i>Journal of Immunology</i> , 2010, 185, 2502-2515.	0.4	126
15	Platelets Control Leukocyte Recruitment in a Murine Model of Cutaneous Arthus Reaction. <i>American Journal of Pathology</i> , 2010, 176, 259-269.	1.9	57
16	Increased Serum Pentraxin 3 in Patients with Systemic Sclerosis. <i>Journal of Rheumatology</i> , 2009, 36, 976-983.	1.0	41
17	Reduced red blood cell velocity in nail-fold capillaries as a sensitive and specific indicator of microcirculation injury in systemic sclerosis. <i>Rheumatology</i> , 2009, 48, 696-703.	0.9	47
18	Clinical Significance of Serum HMGB-1 and sRAGE Levels in Systemic Sclerosis: Association with Disease Severity. <i>Journal of Clinical Immunology</i> , 2009, 29, 180-189.	2.0	96

#	ARTICLE	IF	CITATIONS
19	P-Selectin Glycoprotein Ligand-1 Contributes to Wound Healing Predominantly as a P-Selectin Ligand and Partly as an E-Selectin Ligand. <i>Journal of Investigative Dermatology</i> , 2009, 129, 2059-2067.	0.3	11
20	Low Zone Tolerance Requires ICAM-1 Expression to Limit Contact Hypersensitivity Elicitation. <i>Journal of Investigative Dermatology</i> , 2009, 129, 2661-2667.	0.3	2
21	Clinical association of serum CD137 (4-1BB) levels in patients with systemic sclerosis. <i>Journal of Dermatological Science</i> , 2009, 53, 159-161.	1.0	3
22	Decreased levels of autoantibody against histone deacetylase 3 in patients with systemic sclerosis. <i>Autoimmunity</i> , 2009, 42, 120-125.	1.2	14
23	CD19, a Response Regulator of B Lymphocytes, Regulates Wound Healing through Hyaluronan-Induced TLR4 Signaling. <i>American Journal of Pathology</i> , 2009, 175, 649-660.	1.9	84
24	Elevated Serum Concentrations of Polymorphonuclear Neutrophilic Leukocyte Elastase in Systemic Sclerosis: Association with Pulmonary Fibrosis. <i>Journal of Rheumatology</i> , 2009, 36, 99-105.	1.0	15
25	Autoantibody-mediated regulation of B cell responses by functional anti-CD22 autoantibodies in patients with systemic sclerosis. <i>Clinical and Experimental Immunology</i> , 2009, 159, 176-184.	1.1	28
26	CD19 regulates the development of bleomycin-induced pulmonary fibrosis in a mouse model. <i>Arthritis and Rheumatism</i> , 2008, 58, 3574-3584.	6.7	73
27	The clinical relevance of serum antinuclear antibodies in Japanese patients with systemic sclerosis. <i>British Journal of Dermatology</i> , 2008, 158, 487-495.	1.4	108
28	Elevation of IgG levels is a serological indicator for pulmonary fibrosis in systemic sclerosis with anti-topoisomerase I antibodies and those with anticentromere antibody. <i>Clinical and Experimental Dermatology</i> , 2008, 33, 329-332.	0.6	13
29	Involvement of gaseous low molecular monoxides in the cutaneous reverse passive Arthus reaction: cytoprotective action of carbon monoxide. <i>Clinical and Experimental Immunology</i> , 2008, 153, 245-257.	1.1	7
30	Clinical association of serum interleukin-17 levels in systemic sclerosis: Is systemic sclerosis a Th17 disease?. <i>Journal of Dermatological Science</i> , 2008, 50, 240-242.	1.0	110
31	Autoantibodies against matrix metalloproteinase-1 in patients with localized scleroderma. <i>Journal of Dermatological Science</i> , 2008, 52, 47-54.	1.0	38
32	CD19 Regulates Skin and Lung Fibrosis via Toll-Like Receptor Signaling in a Model of Bleomycin-Induced Scleroderma. <i>American Journal of Pathology</i> , 2008, 172, 1650-1663.	1.9	192
33	Increased Serum Levels of N ^ε -(hexanoyl)lysine, A New Marker of Oxidative Stress, in Systemic Sclerosis. <i>Journal of Rheumatology</i> , 2008, 35, 2214-2219.	1.0	23
34	Blockade of CD40/CD40 ligand interactions attenuates skin fibrosis and autoimmunity in the tight-skin mouse. <i>Annals of the Rheumatic Diseases</i> , 2008, 67, 867-872.	0.5	50
35	Increased Serum Soluble OX40 in Patients with Systemic Sclerosis. <i>Journal of Rheumatology</i> , 2008, 35, 2359-2362.	1.0	68
36	Correlation of IgE Autoantibody to BP180 With a Severe Form of Bullous Pemphigoid. <i>Archives of Dermatology</i> , 2008, 144, 41-8.	1.7	121

#	ARTICLE	IF	CITATIONS
37	Three Cases of Juvenile Systemic Sclerosis. <i>Nishinihon Journal of Dermatology</i> , 2008, 70, 371-376.	0.0	1
38	A Case of Child-Onset Eosinophilic Fasciitis. <i>Nishinihon Journal of Dermatology</i> , 2008, 70, 614-617.	0.0	1
39	A Case of Squamous Cell Carcinoma Arising from Discoid Lupus Erythematosus on the Sole. <i>Nishinihon Journal of Dermatology</i> , 2008, 70, 23-26.	0.0	0
40	Identification of a novel autoantibody reactive with 155 and 140 kDa nuclear proteins in patients with dermatomyositis: an association with malignancy. <i>Rheumatology</i> , 2007, 46, 25-28.	0.9	277
41	Autoantibody against peroxiredoxin I, an antioxidant enzyme, in patients with systemic sclerosis: possible association with oxidative stress. <i>Rheumatology</i> , 2007, 46, 790-795.	0.9	37
42	Endothelial selectins regulate skin wound healing in cooperation with L-selectin and ICAM-1. <i>Journal of Leukocyte Biology</i> , 2007, 82, 519-531.	1.5	39
43	BAFF Antagonist Attenuates the Development of Skin Fibrosis in Tight-Skin Mice. <i>Journal of Investigative Dermatology</i> , 2007, 127, 2772-2780.	0.3	69
44	Antigen specificity of antihistone antibodies in connective tissue disease patients with anti-U1RNP antibodies. <i>Rheumatology International</i> , 2007, 28, 113-119.	1.5	4
45	Clinical evaluation of anti-aminoacyl tRNA synthetase antibodies in Japanese patients with dermatomyositis. <i>Journal of Rheumatology</i> , 2007, 34, 1012-8.	1.0	62
46	Inhibitory Role of CD19 in the Progression of Experimental Autoimmune Encephalomyelitis by Regulating Cytokine Response. <i>American Journal of Pathology</i> , 2006, 168, 812-821.	1.9	109
47	B-Lymphocyte Depletion Reduces Skin Fibrosis and Autoimmunity in the Tight-Skin Mouse Model for Systemic Sclerosis. <i>American Journal of Pathology</i> , 2006, 169, 954-966.	1.9	195
48	Antibody isotype-specific engagement of Fc γ 3 receptors regulates B lymphocyte depletion during CD20 immunotherapy. <i>Journal of Experimental Medicine</i> , 2006, 203, 743-753.	4.2	238
49	Regulation of local and metastatic host-mediated anti-tumour mechanisms by l-selectin and intercellular adhesion molecule-1. <i>Clinical and Experimental Immunology</i> , 2006, 143, 216-227.	1.1	34
50	A Clue for Telangiectasis in Systemic Sclerosis: Elevated Serum Soluble Endoglin Levels in Patients with the Limited Cutaneous Form of the Disease. <i>Dermatology</i> , 2006, 213, 88-92.	0.9	38
51	Serum levels of monocyte chemotactic protein-3/CCL7 are raised in patients with systemic sclerosis: association with extent of skin sclerosis and severity of pulmonary fibrosis. <i>Annals of the Rheumatic Diseases</i> , 2006, 65, 124-126.	0.5	37
52	Drug-Induced Hypersensitivity Syndrome Associated with Human Herpesvirus 6 and Cytomegalovirus Reactivation. <i>Journal of Dermatology</i> , 2005, 32, 976-981.	0.6	14
53	Abnormal Natural Killer Cell Function in Systemic Sclerosis: Altered Cytokine Production and Defective Killing Activity. <i>Journal of Investigative Dermatology</i> , 2005, 125, 731-737.	0.3	55
54	Serum pulmonary and activation-regulated chemokine/CCL18 levels in patients with systemic sclerosis: A sensitive indicator of active pulmonary fibrosis. <i>Arthritis and Rheumatism</i> , 2005, 52, 2889-2896.	6.7	107

#	ARTICLE	IF	CITATIONS
55	Increased cutaneous T-cell-attracting chemokine levels in sera from patients with systemic sclerosis. <i>Rheumatology</i> , 2005, 44, 873-878.	0.9	29
56	Prevalence and clinical characteristics of anti-Mi-2 antibodies in Japanese patients with dermatomyositis. <i>Journal of Dermatological Science</i> , 2005, 40, 215-217.	1.0	51
57	Anti-lipoprotein lipase antibody in systemic sclerosis: association with elevated serum triglyceride concentrations. <i>Journal of Rheumatology</i> , 2005, 32, 629-36.	1.0	29
58	Serum soluble CTLA-4 levels are increased in diffuse cutaneous systemic sclerosis. <i>British Journal of Rheumatology</i> , 2004, 43, 1261-1266.	2.5	73
59	Autoantibodies against phosphatidylserine-prothrombin complex in patients with systemic sclerosis. <i>Annals of the Rheumatic Diseases</i> , 2004, 63, 1514-1517.	0.5	17
60	P-selectin glycoprotein ligand-1 is required for the development of cutaneous vasculitis induced by immune complex deposition. <i>Journal of Leukocyte Biology</i> , 2004, 76, 374-382.	1.5	15
61	Autoantibody against matrix metalloproteinase-3 in patients with systemic sclerosis. <i>Clinical and Experimental Immunology</i> , 2004, 138, 357-363.	1.1	76
62	Autoantibodies to a collagen-specific molecular chaperone, heat-shock protein 47, in systemic sclerosis. <i>Clinical and Experimental Immunology</i> , 2004, 138, 534-539.	1.1	18
63	Management of Squamous Cell Carcinoma in a Patient with Recessive-Type Epidermolysis Bullosa Dystrophica. <i>Dermatologic Surgery</i> , 2004, 30, 1424-1429.	0.4	32
64	Elevated circulating CD40L concentrations in patients with systemic sclerosis. <i>Journal of Rheumatology</i> , 2004, 31, 514-9.	1.0	37
65	L-Selectin or ICAM-1 Deficiency Reduces an Immediate-Type Hypersensitivity Response by Preventing Mast Cell Recruitment in Repeated Elicitation of Contact Hypersensitivity. <i>Journal of Immunology</i> , 2003, 170, 4325-4334.	0.4	54
66	Ultraviolet Light Exposure Suppresses Contact Hypersensitivity by Abrogating Endothelial Intercellular Adhesion Molecule-1 Up-Regulation at the Elicitation Site. <i>Journal of Immunology</i> , 2003, 171, 2855-2862.	0.4	9
67	Elevated levels of circulating CD44 in patients with systemic sclerosis: association with a milder subset. <i>British Journal of Rheumatology</i> , 2002, 41, 1149-1154.	2.5	11
68	The Cutaneous Reverse Arthus Reaction Requires Intercellular Adhesion Molecule 1 and L-Selectin Expression. <i>Journal of Immunology</i> , 2002, 168, 2970-2978.	0.4	42
69	Intercellular Adhesion Molecule-1 and L-Selectin Regulate Bleomycin-Induced Lung Fibrosis. <i>American Journal of Pathology</i> , 2002, 161, 1607-1618.	1.9	55
70	Idiopathic Intracranial Hypertension with Elevated Cerebrospinal Fluid Level of Interleukin-6 in a Patient with Systemic Lupus Erythematosus. <i>Clinical Rheumatology</i> , 2002, 21, 267-268.	1.0	11
71	CD19-dependent B lymphocyte signaling thresholds influence skin fibrosis and autoimmunity in the tight-skin mouse. <i>Journal of Clinical Investigation</i> , 2002, 109, 1453-1462.	3.9	188
72	CD19-dependent B lymphocyte signaling thresholds influence skin fibrosis and autoimmunity in the tight-skin mouse. <i>Journal of Clinical Investigation</i> , 2002, 109, 1453-1462.	3.9	93