

Li Yan

List of Publications by Year in descending order

Source: <https://exaly.com/author-pdf/4690311/publications.pdf>

Version: 2024-02-01

70
papers

2,567
citations

126708

33
h-index

205818

48
g-index

70
all docs

70
docs citations

70
times ranked

1961
citing authors

#	ARTICLE	IF	CITATIONS
1	Tumor Heterogeneity Correlates with Less Immune Response and Worse Survival in Breast Cancer Patients. <i>Annals of Surgical Oncology</i> , 2019, 26, 2191-2199.	0.7	127
2	CD8 T Cell Score as a Prognostic Biomarker for Triple Negative Breast Cancer. <i>International Journal of Molecular Sciences</i> , 2020, 21, 6968.	1.8	118
3	M1 Macrophage and M1/M2 ratio defined by transcriptomic signatures resemble only part of their conventional clinical characteristics in breast cancer. <i>Scientific Reports</i> , 2020, 10, 16554.	1.6	109
4	Cytolytic Activity Score to Assess Anticancer Immunity in Colorectal Cancer. <i>Annals of Surgical Oncology</i> , 2018, 25, 2323-2331.	0.7	107
5	G2M Cell Cycle Pathway Score as a Prognostic Biomarker of Metastasis in Estrogen Receptor (ER)-Positive Breast Cancer. <i>International Journal of Molecular Sciences</i> , 2020, 21, 2921.	1.8	100
6	Triple-Negative Breast Cancer with High Levels of Annexin A1 Expression Is Associated with Mast Cell Infiltration, Inflammation, and Angiogenesis. <i>International Journal of Molecular Sciences</i> , 2019, 20, 4197.	1.8	81
7	Tumor Infiltrating Lymphocytes and Macrophages Improve Survival in Microsatellite Unstable Colorectal Cancer. <i>Scientific Reports</i> , 2019, 9, 13455.	1.6	80
8	Validation of olfactory deficit as a biomarker of Alzheimer disease. <i>Neurology: Clinical Practice</i> , 2017, 7, 5-14.	0.8	78
9	Pancreatic adenocarcinomas with mature blood vessels have better overall survival. <i>Scientific Reports</i> , 2019, 9, 1310.	1.6	77
10	The E2F Pathway Score as a Predictive Biomarker of Response to Neoadjuvant Therapy in ER+/HER2 ⁺ Breast Cancer. <i>Cells</i> , 2020, 9, 1643.	1.8	76
11	Abundance of Regulatory T Cell (Treg) as a Predictive Biomarker for Neoadjuvant Chemotherapy in Triple-Negative Breast Cancer. <i>Cancers</i> , 2020, 12, 3038.	1.7	66
12	Biologically Aggressive Phenotype and Anti-cancer Immunity Counterbalance in Breast Cancer with High Mutation Rate. <i>Scientific Reports</i> , 2020, 10, 1852.	1.6	65
13	Overexpression of suppressive microRNAs, miR-30a and miR-200c are associated with improved survival of breast cancer patients. <i>Scientific Reports</i> , 2017, 7, 15945.	1.6	62
14	Plasmacytoid Dendritic Cell (pDC) Infiltration Correlate with Tumor Infiltrating Lymphocytes, Cancer Immunity, and Better Survival in Triple Negative Breast Cancer (TNBC) More Strongly than Conventional Dendritic Cell (cDC). <i>Cancers</i> , 2020, 12, 3342.	1.7	62
15	Estrogen Receptor Positive Breast Cancer with High Expression of Androgen Receptor has Less Cytolytic Activity and Worse Response to Neoadjuvant Chemotherapy but Better Survival. <i>International Journal of Molecular Sciences</i> , 2019, 20, 2655.	1.8	59
16	ABCC1-Exported Sphingosine-1-phosphate, Produced by Sphingosine Kinase 1, Shortens Survival of Mice and Patients with Breast Cancer. <i>Molecular Cancer Research</i> , 2018, 16, 1059-1070.	1.5	58
17	Intra-Tumoral Angiogenesis Is Associated with Inflammation, Immune Reaction and Metastatic Recurrence in Breast Cancer. <i>International Journal of Molecular Sciences</i> , 2020, 21, 6708.	1.8	56
18	KRAS signaling enriched triple negative breast cancer is associated with favorable tumor immune microenvironment and better survival. <i>American Journal of Cancer Research</i> , 2020, 10, 897-907.	1.4	54

#	ARTICLE	IF	CITATIONS
19	Clinical Relevance of microRNA Expressions in Breast Cancer Validated Using the Cancer Genome Atlas (TCGA). <i>Annals of Surgical Oncology</i> , 2017, 24, 2943-2949.	0.7	51
20	Doxorubicin effect is enhanced by sphingosine-1-phosphate signaling antagonist in breast cancer. <i>Journal of Surgical Research</i> , 2017, 219, 202-213.	0.8	46
21	Immune Cytolytic Activity for Comprehensive Understanding of Immune Landscape in Hepatocellular Carcinoma. <i>Cancers</i> , 2020, 12, 1221.	1.7	46
22	A Novel 4-gene Score to Predict Survival, Distant Metastasis and Response to Neoadjuvant Therapy in Breast Cancer. <i>Cancers</i> , 2020, 12, 1148.	1.7	46
23	Late recurrence of breast cancer is associated with pro-cancerous immune microenvironment in the primary tumor. <i>Scientific Reports</i> , 2019, 9, 16942.	1.6	44
24	High G2M Pathway Score Pancreatic Cancer is Associated with Worse Survival, Particularly after Margin-Positive (R1 or R2) Resection. <i>Cancers</i> , 2020, 12, 2871.	1.7	41
25	Degree of Early Estrogen Response Predict Survival after Endocrine Therapy in Primary and Metastatic ER-Positive Breast Cancer. <i>Cancers</i> , 2020, 12, 3557.	1.7	41
26	Intratumoral Adipocyte-High Breast Cancer Enrich for Metastatic and Inflammation-Related Pathways but Associated with Less Cancer Cell Proliferation. <i>International Journal of Molecular Sciences</i> , 2020, 21, 5744.	1.8	39
27	High expression of bone morphogenetic protein (BMP) 6 and BMP7 are associated with higher immune cell infiltration and better survival in estrogen receptor-positive breast cancer. <i>Oncology Reports</i> , 2019, 42, 1413-1421.	1.2	38
28	High Expression of microRNA-143 is Associated with Favorable Tumor Immune Microenvironment and Better Survival in Estrogen Receptor Positive Breast Cancer. <i>International Journal of Molecular Sciences</i> , 2020, 21, 3213.	1.8	38
29	Tamoxifen sensitivity-related microRNA-342 is a useful biomarker for breast cancer survival. <i>Oncotarget</i> , 2017, 8, 99978-99989.	0.8	38
30	Enhanced DNA Repair Pathway is Associated with Cell Proliferation and Worse Survival in Hepatocellular Carcinoma (HCC). <i>Cancers</i> , 2021, 13, 323.	1.7	36
31	High Expression of miR-34a Associated with Less Aggressive Cancer Biology but Not with Survival in Breast Cancer. <i>International Journal of Molecular Sciences</i> , 2020, 21, 3045.	1.8	35
32	High expression of SLCO2B1 is associated with prostate cancer recurrence after radical prostatectomy. <i>Oncotarget</i> , 2018, 9, 14207-14218.	0.8	35
33	Novel MicroRNA-Based Risk Score Identified by Integrated Analyses to Predict Metastasis and Poor Prognosis in Breast Cancer. <i>Annals of Surgical Oncology</i> , 2018, 25, 4037-4046.	0.7	34
34	Inflammation Is Associated with Worse Outcome in the Whole Cohort but with Better Outcome in Triple-Negative Subtype of Breast Cancer Patients. <i>Journal of Immunology Research</i> , 2020, 2020, 1-17.	0.9	34
35	Molecular Biological Features of Nottingham Histological Grade 3 Breast Cancers. <i>Annals of Surgical Oncology</i> , 2020, 27, 4475-4485.	0.7	34
36	Expression of MicroRNA-9 is Associated With Overall Survival in Breast Cancer Patients. <i>Journal of Surgical Research</i> , 2019, 233, 426-435.	0.8	33

#	ARTICLE	IF	CITATIONS
37	High MYC mRNA Expression Is More Clinically Relevant than MYC DNA Amplification in Triple-Negative Breast Cancer. <i>International Journal of Molecular Sciences</i> , 2020, 21, 217.	1.8	33
38	ITPKC as a Prognostic and Predictive Biomarker of Neoadjuvant Chemotherapy for Triple Negative Breast Cancer. <i>Cancers</i> , 2020, 12, 2758.	1.7	33
39	High Expression of NRF2 Is Associated with Increased Tumor-Infiltrating Lymphocytes and Cancer Immunity in ER-Positive/HER2-Negative Breast Cancer. <i>Cancers</i> , 2020, 12, 3856.	1.7	32
40	High expression of Annexin A2 is associated with DNA repair, metabolic alteration, and worse survival in pancreatic ductal adenocarcinoma. <i>Surgery</i> , 2019, 166, 150-156.	1.0	29
41	Annexin A1 Expression Is Associated with Epithelialâ€“Mesenchymal Transition (EMT), Cell Proliferation, Prognosis, and Drug Response in Pancreatic Cancer. <i>Cells</i> , 2021, 10, 653.	1.8	27
42	Adipogenesis in triple-negative breast cancer is associated with unfavorable tumor immune microenvironment and with worse survival. <i>Scientific Reports</i> , 2021, 11, 12541.	1.6	25
43	Transcriptomic Profile of Lymphovascular Invasion, a Known Risk Factor of Pancreatic Ductal Adenocarcinoma Metastasis. <i>Cancers</i> , 2020, 12, 2033.	1.7	24
44	Th2 cell infiltrations predict neoadjuvant chemotherapy response of estrogen receptor-positive breast cancer. <i>Gland Surgery</i> , 2021, 10, 154-165.	0.5	24
45	High expression of polo-like kinase 1 is associated with TP53 inactivation, DNA repair deficiency, and worse prognosis in ER positive Her2 negative breast cancer. <i>American Journal of Translational Research (discontinued)</i> , 2019, 11, 6507-6521.	0.0	24
46	A Novel Four-Gene Score to Predict Pathologically Complete (R0) Resection and Survival in Pancreatic Cancer. <i>Cancers</i> , 2020, 12, 3635.	1.7	20
47	Abundance of reactive oxygen species (ROS) is associated with tumor aggressiveness, immune response, and worse survival in breast cancer. <i>Breast Cancer Research and Treatment</i> , 2022, 194, 231-241.	1.1	20
48	Abundance of Microvascular Endothelial Cells Is Associated with Response to Chemotherapy and Prognosis in Colorectal Cancer. <i>Cancers</i> , 2021, 13, 1477.	1.7	19
49	Low DMT1 Expression Associates With Increasedâ€“Oxidative Phosphorylation and Earlyâ€“Recurrenceâ€“ in Hepatocellular Carcinoma. <i>Journal of Surgical Research</i> , 2019, 234, 343-352.	0.8	17
50	A Novel Three-Gene Score as a Predictive Biomarker for Pathologically Complete Response after Neoadjuvant Chemotherapy in Triple-Negative Breast Cancer. <i>Cancers</i> , 2021, 13, 2401.	1.7	16
51	The Unfolded Protein Response Is Associated with Cancer Proliferation and Worse Survival in Hepatocellular Carcinoma. <i>Cancers</i> , 2021, 13, 4443.	1.7	12
52	Organoids Are Limited in Modeling the Colon Adenomaâ€“Carcinoma Sequence. <i>Cells</i> , 2021, 10, 488.	1.8	11
53	Low expression of miR-29a is associated with aggressive biology and worse survival in gastric cancer. <i>Scientific Reports</i> , 2021, 11, 14134.	1.6	10
54	NR2F1, a Tumor Dormancy Marker, Is Expressed Predominantly in Cancer-Associated Fibroblasts and Is Associated with Suppressed Breast Cancer Cell Proliferation. <i>Cancers</i> , 2022, 14, 2962.	1.7	10

#	ARTICLE	IF	CITATIONS
55	Angiogenesis is associated with an attenuated tumor microenvironment, aggressive biology, and worse survival in gastric cancer patients. <i>American Journal of Cancer Research</i> , 2021, 11, 1659-1671.	1.4	7
56	Octogenariansâ€™ Breast Cancer Is Associated with an Unfavorable Tumor Immune Microenvironment and Worse Disease-Free Survival. <i>Cancers</i> , 2021, 13, 2933.	1.7	4
57	A novel five-gene score to predict complete pathological response to neoadjuvant chemotherapy in ER-positive/HER2-negative breast cancer. <i>American Journal of Cancer Research</i> , 2021, 11, 3611-3627.	1.4	4
58	Low RUFY3 expression level is associated with lymph node metastasis in older women with invasive breast cancer. <i>Breast Cancer Research and Treatment</i> , 2022, 192, 19-32.	1.1	4
59	G2M checkpoint pathway alone is associated with drug response and survival among cell proliferation-related pathways in pancreatic cancer. <i>American Journal of Cancer Research</i> , 2021, 11, 3070-3084.	1.4	3
60	Immune cytolytic activity is associated with reduced intra-tumoral genetic heterogeneity and with better clinical outcomes in triple negative breast cancer. <i>American Journal of Cancer Research</i> , 2021, 11, 3628-3644.	1.4	3
61	Confidence interval estimation of the common mean of several gamma populations. <i>PLoS ONE</i> , 2022, 17, e0269971.	1.1	3
62	A prognostic score based on long-term survivor unique transcriptomic signatures predicts patient survival in pancreatic ductal adenocarcinoma. <i>American Journal of Cancer Research</i> , 2021, 11, 4294-4307.	1.4	2
63	MELK expression in breast cancer is associated with infiltration of immune cell and pathological complete response (pCR) after neoadjuvant chemotherapy. <i>American Journal of Cancer Research</i> , 2021, 11, 4421-4437.	1.4	2
64	Conflicting roles of expression by subtypes in breast cancer. <i>American Journal of Cancer Research</i> , 2021, 11, 5094-5110.	1.4	2
65	Low expression of miR-195 is associated with cell proliferation, glycolysis and poor survival in estrogen receptor (ER)-positive but not in triple negative breast cancer. <i>American Journal of Cancer Research</i> , 2021, 11, 3320-3334.	1.4	1
66	Low intratumoral genetic neutrophil-to-lymphocyte ratio (NLR) is associated with favorable tumor immune microenvironment and with survival in triple negative breast cancer (TNBC). <i>American Journal of Cancer Research</i> , 2021, 11, 5743-5755.	1.4	1
67	Intratumoral PDGFB gene predominantly expressed in endothelial cells is associated with angiogenesis and lymphangiogenesis, but not with metastasis in breast cancer. <i>Breast Cancer Research and Treatment</i> , 2022, 195, 17-31.	1.1	1
68	Development of KAM score to predict metastasis and worse survival in breast cancer. <i>American Journal of Cancer Research</i> , 2021, 11, 5388-5401.	1.4	0
69	Intratumoral density of regulatory T cells is a predictor of host immune response and chemotherapy response in colorectal cancer.. <i>American Journal of Cancer Research</i> , 2022, 12, 490-503.	1.4	0
70	APOBEC3F expression in triple-negative breast cancer is associated with tumor microenvironment infiltration and activation of cancer immunity and improved survival.. <i>American Journal of Cancer Research</i> , 2022, 12, 744-762.	1.4	0