

Li Yan

List of Publications by Year in descending order

Source: <https://exaly.com/author-pdf/4690311/publications.pdf>

Version: 2024-02-01

70
papers

2,567
citations

126708

33
h-index

205818

48
g-index

70
all docs

70
docs citations

70
times ranked

1961
citing authors

| # | ARTICLE | IF | CITATIONS |
|----|---|-----|-----------|
| 1 | Low RUFY3 expression level is associated with lymph node metastasis in older women with invasive breast cancer. <i>Breast Cancer Research and Treatment</i> , 2022, 192, 19-32. | 1.1 | 4 |
| 2 | Intratumoral density of regulatory T cells is a predictor of host immune response and chemotherapy response in colorectal cancer. <i>American Journal of Cancer Research</i> , 2022, 12, 490-503. | 1.4 | 0 |
| 3 | APOBEC3F expression in triple-negative breast cancer is associated with tumor microenvironment infiltration and activation of cancer immunity and improved survival. <i>American Journal of Cancer Research</i> , 2022, 12, 744-762. | 1.4 | 0 |
| 4 | Abundance of reactive oxygen species (ROS) is associated with tumor aggressiveness, immune response, and worse survival in breast cancer. <i>Breast Cancer Research and Treatment</i> , 2022, 194, 231-241. | 1.1 | 20 |
| 5 | Confidence interval estimation of the common mean of several gamma populations. <i>PLoS ONE</i> , 2022, 17, e0269971. | 1.1 | 3 |
| 6 | NR2F1, a Tumor Dormancy Marker, Is Expressed Predominantly in Cancer-Associated Fibroblasts and Is Associated with Suppressed Breast Cancer Cell Proliferation. <i>Cancers</i> , 2022, 14, 2962. | 1.7 | 10 |
| 7 | Intratumoral PDGFB gene predominantly expressed in endothelial cells is associated with angiogenesis and lymphangiogenesis, but not with metastasis in breast cancer. <i>Breast Cancer Research and Treatment</i> , 2022, 195, 17-31. | 1.1 | 1 |
| 8 | Enhanced DNA Repair Pathway is Associated with Cell Proliferation and Worse Survival in Hepatocellular Carcinoma (HCC). <i>Cancers</i> , 2021, 13, 323. | 1.7 | 36 |
| 9 | Th2 cell infiltrations predict neoadjuvant chemotherapy response of estrogen receptor-positive breast cancer. <i>Gland Surgery</i> , 2021, 10, 154-165. | 0.5 | 24 |
| 10 | Organoids Are Limited in Modeling the Colon Adenomaâ€“Carcinoma Sequence. <i>Cells</i> , 2021, 10, 488. | 1.8 | 11 |
| 11 | Abundance of Microvascular Endothelial Cells Is Associated with Response to Chemotherapy and Prognosis in Colorectal Cancer. <i>Cancers</i> , 2021, 13, 1477. | 1.7 | 19 |
| 12 | Annexin A1 Expression Is Associated with Epithelialâ€“Mesenchymal Transition (EMT), Cell Proliferation, Prognosis, and Drug Response in Pancreatic Cancer. <i>Cells</i> , 2021, 10, 653. | 1.8 | 27 |
| 13 | A Novel Three-Gene Score as a Predictive Biomarker for Pathologically Complete Response after Neoadjuvant Chemotherapy in Triple-Negative Breast Cancer. <i>Cancers</i> , 2021, 13, 2401. | 1.7 | 16 |
| 14 | Adipogenesis in triple-negative breast cancer is associated with unfavorable tumor immune microenvironment and with worse survival. <i>Scientific Reports</i> , 2021, 11, 12541. | 1.6 | 25 |
| 15 | Octogenariansâ€™ Breast Cancer Is Associated with an Unfavorable Tumor Immune Microenvironment and Worse Disease-Free Survival. <i>Cancers</i> , 2021, 13, 2933. | 1.7 | 4 |
| 16 | Low expression of miR-29a is associated with aggressive biology and worse survival in gastric cancer. <i>Scientific Reports</i> , 2021, 11, 14134. | 1.6 | 10 |
| 17 | The Unfolded Protein Response Is Associated with Cancer Proliferation and Worse Survival in Hepatocellular Carcinoma. <i>Cancers</i> , 2021, 13, 4443. | 1.7 | 12 |
| 18 | Angiogenesis is associated with an attenuated tumor microenvironment, aggressive biology, and worse survival in gastric cancer patients. <i>American Journal of Cancer Research</i> , 2021, 11, 1659-1671. | 1.4 | 7 |

| # | ARTICLE | IF | CITATIONS |
|----|---|-----|-----------|
| 19 | G2M checkpoint pathway alone is associated with drug response and survival among cell proliferation-related pathways in pancreatic cancer. American Journal of Cancer Research, 2021, 11, 3070-3084. | 1.4 | 3 |
| 20 | Low expression of miR-195 is associated with cell proliferation, glycolysis and poor survival in estrogen receptor (ER)-positive but not in triple negative breast cancer. American Journal of Cancer Research, 2021, 11, 3320-3334. | 1.4 | 1 |
| 21 | A novel five-gene score to predict complete pathological response to neoadjuvant chemotherapy in ER-positive/HER2-negative breast cancer. American Journal of Cancer Research, 2021, 11, 3611-3627. | 1.4 | 4 |
| 22 | Immune cytolytic activity is associated with reduced intra-tumoral genetic heterogeneity and with better clinical outcomes in triple negative breast cancer. American Journal of Cancer Research, 2021, 11, 3628-3644. | 1.4 | 3 |
| 23 | A prognostic score based on long-term survivor unique transcriptomic signatures predicts patient survival in pancreatic ductal adenocarcinoma. American Journal of Cancer Research, 2021, 11, 4294-4307. | 1.4 | 2 |
| 24 | MELK expression in breast cancer is associated with infiltration of immune cell and pathological complete response (pCR) after neoadjuvant chemotherapy. American Journal of Cancer Research, 2021, 11, 4421-4437. | 1.4 | 2 |
| 25 | Conflicting roles of expression by subtypes in breast cancer. American Journal of Cancer Research, 2021, 11, 5094-5110. | 1.4 | 2 |
| 26 | Development of KAM score to predict metastasis and worse survival in breast cancer. American Journal of Cancer Research, 2021, 11, 5388-5401. | 1.4 | 0 |
| 27 | Low intratumoral genetic neutrophil-to-lymphocyte ratio (NLR) is associated with favorable tumor immune microenvironment and with survival in triple negative breast cancer (TNBC). American Journal of Cancer Research, 2021, 11, 5743-5755. | 1.4 | 1 |
| 28 | High MYC mRNA Expression Is More Clinically Relevant than MYC DNA Amplification in Triple-Negative Breast Cancer. International Journal of Molecular Sciences, 2020, 21, 217. | 1.8 | 33 |
| 29 | CD8 T Cell Score as a Prognostic Biomarker for Triple Negative Breast Cancer. International Journal of Molecular Sciences, 2020, 21, 6968. | 1.8 | 118 |
| 30 | Abundance of Regulatory T Cell (Treg) as a Predictive Biomarker for Neoadjuvant Chemotherapy in Triple-Negative Breast Cancer. Cancers, 2020, 12, 3038. | 1.7 | 66 |
| 31 | High G2M Pathway Score Pancreatic Cancer is Associated with Worse Survival, Particularly after Margin-Positive (R1 or R2) Resection. Cancers, 2020, 12, 2871. | 1.7 | 41 |
| 32 | M1 Macrophage and M1/M2 ratio defined by transcriptomic signatures resemble only part of their conventional clinical characteristics in breast cancer. Scientific Reports, 2020, 10, 16554. | 1.6 | 109 |
| 33 | ITPKC as a Prognostic and Predictive Biomarker of Neoadjuvant Chemotherapy for Triple Negative Breast Cancer. Cancers, 2020, 12, 2758. | 1.7 | 33 |
| 34 | The E2F Pathway Score as a Predictive Biomarker of Response to Neoadjuvant Therapy in ER+/HER2 ⁺ Breast Cancer. Cells, 2020, 9, 1643. | 1.8 | 76 |
| 35 | Plasmacytoid Dendritic Cell (pDC) Infiltration Correlate with Tumor Infiltrating Lymphocytes, Cancer Immunity, and Better Survival in Triple Negative Breast Cancer (TNBC) More Strongly than Conventional Dendritic Cell (cDC). Cancers, 2020, 12, 3342. | 1.7 | 62 |
| 36 | Degree of Early Estrogen Response Predict Survival after Endocrine Therapy in Primary and Metastatic ER-Positive Breast Cancer. Cancers, 2020, 12, 3557. | 1.7 | 41 |

| # | ARTICLE | IF | CITATIONS |
|----|--|-----|-----------|
| 37 | Transcriptomic Profile of Lymphovascular Invasion, a Known Risk Factor of Pancreatic Ductal Adenocarcinoma Metastasis. <i>Cancers</i> , 2020, 12, 2033. | 1.7 | 24 |
| 38 | Intra-Tumoral Angiogenesis Is Associated with Inflammation, Immune Reaction and Metastatic Recurrence in Breast Cancer. <i>International Journal of Molecular Sciences</i> , 2020, 21, 6708. | 1.8 | 56 |
| 39 | Intratumoral Adipocyte-High Breast Cancer Enrich for Metastatic and Inflammation-Related Pathways but Associated with Less Cancer Cell Proliferation. <i>International Journal of Molecular Sciences</i> , 2020, 21, 5744. | 1.8 | 39 |
| 40 | High Expression of NRF2 Is Associated with Increased Tumor-Infiltrating Lymphocytes and Cancer Immunity in ER-Positive/HER2-Negative Breast Cancer. <i>Cancers</i> , 2020, 12, 3856. | 1.7 | 32 |
| 41 | Inflammation Is Associated with Worse Outcome in the Whole Cohort but with Better Outcome in Triple-Negative Subtype of Breast Cancer Patients. <i>Journal of Immunology Research</i> , 2020, 2020, 1-17. | 0.9 | 34 |
| 42 | A Novel Four-Gene Score to Predict Pathologically Complete (R0) Resection and Survival in Pancreatic Cancer. <i>Cancers</i> , 2020, 12, 3635. | 1.7 | 20 |
| 43 | High Expression of miR-34a Associated with Less Aggressive Cancer Biology but Not with Survival in Breast Cancer. <i>International Journal of Molecular Sciences</i> , 2020, 21, 3045. | 1.8 | 35 |
| 44 | Immune Cytolytic Activity for Comprehensive Understanding of Immune Landscape in Hepatocellular Carcinoma. <i>Cancers</i> , 2020, 12, 1221. | 1.7 | 46 |
| 45 | A Novel 4-gene Score to Predict Survival, Distant Metastasis and Response to Neoadjuvant Therapy in Breast Cancer. <i>Cancers</i> , 2020, 12, 1148. | 1.7 | 46 |
| 46 | High Expression of microRNA-143 is Associated with Favorable Tumor Immune Microenvironment and Better Survival in Estrogen Receptor Positive Breast Cancer. <i>International Journal of Molecular Sciences</i> , 2020, 21, 3213. | 1.8 | 38 |
| 47 | Biologically Aggressive Phenotype and Anti-cancer Immunity Counterbalance in Breast Cancer with High Mutation Rate. <i>Scientific Reports</i> , 2020, 10, 1852. | 1.6 | 65 |
| 48 | G2M Cell Cycle Pathway Score as a Prognostic Biomarker of Metastasis in Estrogen Receptor (ER)-Positive Breast Cancer. <i>International Journal of Molecular Sciences</i> , 2020, 21, 2921. | 1.8 | 100 |
| 49 | Molecular Biological Features of Nottingham Histological Grade 3 Breast Cancers. <i>Annals of Surgical Oncology</i> , 2020, 27, 4475-4485. | 0.7 | 34 |
| 50 | KRAS signaling enriched triple negative breast cancer is associated with favorable tumor immune microenvironment and better survival. <i>American Journal of Cancer Research</i> , 2020, 10, 897-907. | 1.4 | 54 |
| 51 | Triple-Negative Breast Cancer with High Levels of Annexin A1 Expression Is Associated with Mast Cell Infiltration, Inflammation, and Angiogenesis. <i>International Journal of Molecular Sciences</i> , 2019, 20, 4197. | 1.8 | 81 |
| 52 | Tumor Infiltrating Lymphocytes and Macrophages Improve Survival in Microsatellite Unstable Colorectal Cancer. <i>Scientific Reports</i> , 2019, 9, 13455. | 1.6 | 80 |
| 53 | High expression of bone morphogenetic protein (BMP) 6 and BMP7 are associated with higher immune cell infiltration and better survival in estrogen receptor-positive breast cancer. <i>Oncology Reports</i> , 2019, 42, 1413-1421. | 1.2 | 38 |
| 54 | Pancreatic adenocarcinomas with mature blood vessels have better overall survival. <i>Scientific Reports</i> , 2019, 9, 1310. | 1.6 | 77 |

| # | ARTICLE | IF | CITATIONS |
|----|--|-----|-----------|
| 55 | Estrogen Receptor Positive Breast Cancer with High Expression of Androgen Receptor has Less Cytolytic Activity and Worse Response to Neoadjuvant Chemotherapy but Better Survival. <i>International Journal of Molecular Sciences</i> , 2019, 20, 2655. | 1.8 | 59 |
| 56 | High expression of Annexin A2 is associated with DNA repair, metabolic alteration, and worse survival in pancreatic ductal adenocarcinoma. <i>Surgery</i> , 2019, 166, 150-156. | 1.0 | 29 |
| 57 | Tumor Heterogeneity Correlates with Less Immune Response and Worse Survival in Breast Cancer Patients. <i>Annals of Surgical Oncology</i> , 2019, 26, 2191-2199. | 0.7 | 127 |
| 58 | Late recurrence of breast cancer is associated with pro-cancerous immune microenvironment in the primary tumor. <i>Scientific Reports</i> , 2019, 9, 16942. | 1.6 | 44 |
| 59 | Expression of MicroRNA-9 is Associated With Overall Survival in Breast Cancer Patients. <i>Journal of Surgical Research</i> , 2019, 233, 426-435. | 0.8 | 33 |
| 60 | Low DMT1 Expression Associates With Increased Oxidative Phosphorylation and Early Recurrence in Hepatocellular Carcinoma. <i>Journal of Surgical Research</i> , 2019, 234, 343-352. | 0.8 | 17 |
| 61 | High expression of polo-like kinase 1 is associated with TP53 inactivation, DNA repair deficiency, and worse prognosis in ER positive Her2 negative breast cancer. <i>American Journal of Translational Research (discontinued)</i> , 2019, 11, 6507-6521. | 0.0 | 24 |
| 62 | ABCC1-Exported Sphingosine-1-phosphate, Produced by Sphingosine Kinase 1, Shortens Survival of Mice and Patients with Breast Cancer. <i>Molecular Cancer Research</i> , 2018, 16, 1059-1070. | 1.5 | 58 |
| 63 | Novel MicroRNA-Based Risk Score Identified by Integrated Analyses to Predict Metastasis and Poor Prognosis in Breast Cancer. <i>Annals of Surgical Oncology</i> , 2018, 25, 4037-4046. | 0.7 | 34 |
| 64 | Cytolytic Activity Score to Assess Anticancer Immunity in Colorectal Cancer. <i>Annals of Surgical Oncology</i> , 2018, 25, 2323-2331. | 0.7 | 107 |
| 65 | High expression of SLCO2B1 is associated with prostate cancer recurrence after radical prostatectomy. <i>Oncotarget</i> , 2018, 9, 14207-14218. | 0.8 | 35 |
| 66 | Clinical Relevance of microRNA Expressions in Breast Cancer Validated Using the Cancer Genome Atlas (TCGA). <i>Annals of Surgical Oncology</i> , 2017, 24, 2943-2949. | 0.7 | 51 |
| 67 | Overexpression of suppressive microRNAs, miR-30a and miR-200c are associated with improved survival of breast cancer patients. <i>Scientific Reports</i> , 2017, 7, 15945. | 1.6 | 62 |
| 68 | Doxorubicin effect is enhanced by sphingosine-1-phosphate signaling antagonist in breast cancer. <i>Journal of Surgical Research</i> , 2017, 219, 202-213. | 0.8 | 46 |
| 69 | Validation of olfactory deficit as a biomarker of Alzheimer disease. <i>Neurology: Clinical Practice</i> , 2017, 7, 5-14. | 0.8 | 78 |
| 70 | Tamoxifen sensitivity-related microRNA-342 is a useful biomarker for breast cancer survival. <i>Oncotarget</i> , 2017, 8, 99978-99989. | 0.8 | 38 |