Taiki Yamaji

List of Publications by Year in descending order

Source: https://exaly.com/author-pdf/4693611/publications.pdf

Version: 2024-02-01

		109321	82547
189	7,127	35	72
papers	citations	h-index	g-index
191	191	191	13295
all docs	docs citations	times ranked	citing authors

#	Article	IF	CITATIONS
1	Association analysis identifies 65 new breast cancer risk loci. Nature, 2017, 551, 92-94.	27.8	1,099
2	Metagenomic and metabolomic analyses reveal distinct stage-specific phenotypes of the gut microbiota in colorectal cancer. Nature Medicine, 2019, 25, 968-976.	30.7	748
3	Genome-wide association study identifies 112 new loci for body mass index in the Japanese population. Nature Genetics, 2017, 49, 1458-1467.	21.4	380
4	Large-scale genome-wide association study in a Japanese population identifies novel susceptibility loci across different diseases. Nature Genetics, 2020, 52, 669-679.	21.4	304
5	Population-specific and trans-ancestry genome-wide analyses identify distinct and shared genetic risk loci for coronary artery disease. Nature Genetics, 2020, 52, 1169-1177.	21.4	206
6	Natural History of Pulmonary Subsolid Nodules: A Prospective Multicenter Study. Journal of Thoracic Oncology, 2016, 11, 1012-1028.	1.1	184
7	Identification of 28 new susceptibility loci for type 2 diabetes in the Japanese population. Nature Genetics, 2019, 51, 379-386.	21.4	164
8	Identification of six new genetic loci associated with atrial fibrillation in the Japanese population. Nature Genetics, 2017, 49, 953-958.	21.4	136
9	Association of Pancreatic Fatty Infiltration With Pancreatic Ductal Adenocarcinoma. Clinical and Translational Gastroenterology, 2014, 5, e53.	2.5	126
10	Characterizing rare and low-frequency height-associated variants in the Japanese population. Nature Communications, 2019, 10, 4393.	12.8	123
11	Association of Animal and Plant Protein Intake With All-Cause and Cause-Specific Mortality in a Japanese Cohort. JAMA Internal Medicine, 2019, 179, 1509.	5.1	120
12	Interaction between Adiponectin and Leptin Influences the Risk of Colorectal Adenoma. Cancer Research, 2010, 70, 5430-5437.	0.9	115
13	Genome-wide association study identifies seven novel susceptibility loci for primary open-angle glaucoma. Human Molecular Genetics, 2018, 27, 1486-1496.	2.9	111
14	European polygenic risk score for prediction of breast cancer shows similar performance in Asian women. Nature Communications, 2020, 11, 3833.	12.8	88
15	Fruit and vegetable consumption and squamous cell carcinoma of the esophagus in Japan: The JPHC study. International Journal of Cancer, 2008, 123, 1935-1940.	5.1	83
16	Prediction of the 10â€year probability of gastric cancer occurrence in the <scp>J</scp> apanese population: the <scp>JPHC</scp> study cohort <scp>II</scp> . International Journal of Cancer, 2016, 138, 320-331.	5.1	78
17	Genetic polymorphisms of ADH1B, ADH1C and ALDH2, alcohol consumption, and the risk of gastric cancer: the Japan Public Health Center-based prospective study. Carcinogenesis, 2015, 36, 223-231.	2.8	69
18	Association of green tea consumption with mortality due to all causes and major causes of death in a Japanese population: the Japan Public Health Center-based Prospective Study (JPHC Study). Annals of Epidemiology, 2015, 25, 512-518.e3.	1.9	66

#	Article	IF	CITATIONS
19	Investigations in the possibility of early detection of colorectal cancer by gas chromatography/triple-quadrupole mass spectrometry. Oncotarget, 2017, 8, 17115-17126.	1.8	66
20	Plasma 25-hydroxyvitamin D concentration and subsequent risk of total and site specific cancers in Japanese population: large case-cohort study within Japan Public Health Center-based Prospective Study cohort. BMJ: British Medical Journal, 2018, 360, k671.	2.3	61
21	Increased Levels of Branched-Chain Amino Acid Associated With Increased Risk of Pancreatic Cancer in a Prospective Case–Control Study of a Large Cohort. Gastroenterology, 2018, 155, 1474-1482.e1.	1.3	59
22	Association of coffee intake with total and cause-specific mortality in a Japanese population: the Japan Public Health Center–based Prospective Study. American Journal of Clinical Nutrition, 2015, 101, 1029-1037.	4.7	58
23	Visceral Fat Volume and the Prevalence of Colorectal Adenoma. American Journal of Epidemiology, 2009, 170, 1502-1511.	3.4	54
24	Plasma cytokine levels and the presence of colorectal cancer. PLoS ONE, 2019, 14, e0213602.	2.5	54
25	GWAS identifies two novel colorectal cancer loci at 16q24.1 and 20q13.12. Carcinogenesis, 2018, 39, 652-660.	2.8	52
26	Diabetes and cancer risk: A Mendelian randomization study. International Journal of Cancer, 2020, 146, 712-719.	5.1	52
27	Fermented Soy Product Intake Is Inversely Associated with the Development of High Blood Pressure: The Japan Public Health Center-Based Prospective Study. Journal of Nutrition, 2017, 147, 1749-1756.	2.9	51
28	Association between GWAS-identified lung adenocarcinoma susceptibility loci andEGFRmutations in never-smoking Asian women, and comparison with findings from Western populations. Human Molecular Genetics, 2016, 26, ddw414.	2.9	50
29	12 new susceptibility loci for prostate cancer identified by genome-wide association study in Japanese population. Nature Communications, 2019, 10, 4422.	12.8	49
30	Fish, <i>n</i> â^ 3 polyunsaturated fatty acids and <i>n</i> â^ 6 polyunsaturated fatty acids in breast cancer risk: The <scp>J</scp> apan <scp>P</scp> ublic <scp>H</scp> ealth <scp>C</scp> enterâ€based prospective study. International Journal of Cancer, 2015, 137, 2915-2926.	take and 5.1	48
31	Identification of novel breast cancer susceptibility loci in meta-analyses conducted among Asian and European descendants. Nature Communications, 2020, 11, 1217.	12.8	46
32	Transethnic Meta-Analysis of Genome-Wide Association Studies Identifies Three New Loci and Characterizes Population-Specific Differences for Coronary Artery Disease. Circulation Genomic and Precision Medicine, 2020, 13, e002670.	3.6	44
33	Physical inactivity, prolonged sedentary behaviors, and use of visual display terminals as potential risk factors for dry eye disease: JPHC-NEXT study. Ocular Surface, 2020, 18, 56-63.	4.4	42
34	Development of a prediction model for 10-year risk of hepatocellular carcinoma in middle-aged Japanese: The Japan Public Health Center-based Prospective Study Cohort II. Preventive Medicine, 2012, 55, 137-143.	3.4	41
35	Fish, n–3 PUFA consumption, and pancreatic cancer risk in Japanese: a large, population-based, prospective cohort study. American Journal of Clinical Nutrition, 2015, 102, 1490-1497.	4.7	39
36	High hemoglobin A1c levels within the nonâ€diabetic range are associated with the risk of all cancers. International Journal of Cancer, 2016, 138, 1741-1753.	5.1	39

#	Article	IF	Citations
37	Impact of Alcohol Intake and Drinking Patterns on Mortality From All Causes and Major Causes of Death in a Japanese Population. Journal of Epidemiology, 2018, 28, 140-148.	2.4	39
38	Genomeâ€wide association study identifies gastric cancer susceptibility loci at 12q24.11â€12 and 20q11.21. Cancer Science, 2018, 109, 4015-4024.	3.9	39
39	Dietary fiber intake and total and cause-specific mortality: the Japan Public Health Center-based prospective study. American Journal of Clinical Nutrition, 2020, 111, 1027-1035.	4.7	38
40	Association between adherence to the Japanese diet and all-cause and cause-specific mortality: the Japan Public Health Center-based Prospective Study. European Journal of Nutrition, 2021, 60, 1327-1336.	3.9	37
41	Association Between Plasma 25-Hydroxyvitamin D and Colorectal Adenoma According to Dietary Calcium Intake and Vitamin D Receptor Polymorphism. American Journal of Epidemiology, 2012, 175, 236-244.	3.4	35
42	Validating the dietary inflammatory index using inflammatory biomarkers in a Japanese population: A cross-sectional study of the JPHC-FFQ validation study. Nutrition, 2020, 69, 110569.	2.4	35
43	Cruciferous Vegetable Intake Is Inversely Associated with Lung Cancer Risk among Current Nonsmoking Men in the Japan Public Health Center (JPHC) Study. Journal of Nutrition, 2017, 147, 841-849.	2.9	34
44	Perceived stress level and risk of cancer incidence in a Japanese population: the Japan Public Health Center (JPHC)-based Prospective Study. Scientific Reports, 2017, 7, 12964.	3.3	34
45	Genome-wide association meta-analysis identifies GP2 gene risk variants for pancreatic cancer. Nature Communications, 2020, 11, 3175.	12.8	34
46	Hepatitis B and C virus infection and risk of lymphoid malignancies: A population-based cohort study (JPHC Study). Cancer Epidemiology, 2015, 39, 562-566.	1.9	33
47	Hepatitis B and C Virus Infection and Risk of Pancreatic Cancer: A Population-Based Cohort Study (JPHC) Tj ETQq	1 <u>1 0</u> .784	314 rgBT /0\
48	Association between green tea/coffee consumption and biliary tract cancer: A populationâ€based cohort study in Japan. Cancer Science, 2016, 107, 76-83.	3.9	31
49	The Japan Public Health Center-based Prospective Study for the Next Generation (JPHC-NEXT): Study Design and Participants. Journal of Epidemiology, 2020, 30, 46-54.	2.4	30
50	Low-dose CT lung cancer screening in never-smokers and smokers: results of an eight-year observational study. Translational Lung Cancer Research, 2020, 9, 10-22.	2.8	30
51	Association of dietary diversity with total mortality and major causes of mortality in the Japanese population: JPHC study. European Journal of Clinical Nutrition, 2020, 74, 54-66.	2.9	29
52	Dietary consumption of antioxidant vitamins and subsequent lung cancer risk: The <scp>J</scp> apan <scp>P</scp> ublic <scp>H</scp> ealth <scp>C</scp> enterâ€based prospective study. International Journal of Cancer, 2018, 142, 2441-2460.	5.1	28
53	Low carbohydrate diet and all cause and cause-specific mortality. Clinical Nutrition, 2021, 40, 2016-2024.	5.0	28
54	Methionine Synthase A2756G Polymorphism Interacts with Alcohol and Folate Intake to Influence the Risk of Colorectal Adenoma. Cancer Epidemiology Biomarkers and Prevention, 2009, 18, 267-274.	2.5	27

#	Article	IF	CITATIONS
55	Polygenic risk scores for prediction of breast cancer risk in Asian populations. Genetics in Medicine, 2022, 24, 586-600.	2.4	27
56	Circulating inflammatory markers and colorectal cancer risk: A prospective caseâ€cohort study in Japan. International Journal of Cancer, 2018, 143, 2767-2776.	5.1	26
57	Dietary Heterocyclic Amine Intake, <i>NAT2</i> Genetic Polymorphism, and Colorectal Adenoma Risk: The Colorectal Adenoma Study in Tokyo. Cancer Epidemiology Biomarkers and Prevention, 2015, 24, 613-620.	2.5	25
58	Body mass index and colorectal cancer risk: A Mendelian randomization study. Cancer Science, 2021, 112, 1579-1588.	3.9	25
59	Fiber intake and risk of subsequent prostate cancer in Japanese men. American Journal of Clinical Nutrition, 2015, 101, 118-125.	4.7	24
60	Circulating sex hormone levels and colorectal cancer risk in Japanese postmenopausal women: The JPHC nested caseâ€"control study. International Journal of Cancer, 2019, 145, 1238-1244.	5.1	24
61	Dietary patterns and prostate cancer risk in Japanese: the Japan Public Health Center-based Prospective Study (JPHC Study). Cancer Causes and Control, 2018, 29, 589-600.	1.8	23
62	Dietary Inflammatory Index Is Associated With Inflammation in Japanese Men. Frontiers in Nutrition, 2021, 8, 604296.	3.7	23
63	<scp>C</scp> offee intake and the risk of colorectal adenoma: The colorectal adenoma study in Tokyo. International Journal of Cancer, 2015, 137, 463-470.	5.1	22
64	Coffee and green tea consumption in relation to brain tumor risk in a Japanese population. International Journal of Cancer, 2016, 139, 2714-2721.	5.1	22
65	Plasma tea catechins and risk of cardiovascular disease in middle-aged Japanese subjects: The JPHC study. Atherosclerosis, 2018, 277, 90-97.	0.8	22
66	Revisit of an unanswered question by pooled analysis of eight cohort studies in Japan: Does cigarette smoking and alcohol drinking have interaction for the risk of esophageal cancer?. Cancer Medicine, 2019, 8, 6414-6425.	2.8	22
67	Adjustment of Cell-Type Composition Minimizes Systematic Bias in Blood DNA Methylation Profiles Derived by DNA Collection Protocols. PLoS ONE, 2016, 11, e0147519.	2.5	21
68	Vitamin D Receptor Gene Polymorphism and the Risk of Colorectal Cancer: A Nested Case-Control Study. PLoS ONE, 2016, 11, e0164648.	2.5	21
69	Inclusion of a Genetic Risk Score into a Validated Risk Prediction Model for Colorectal Cancer in Japanese Men Improves Performance. Cancer Prevention Research, 2017, 10, 535-541.	1.5	21
70	The relationship between vegetable/fruit consumption and gallbladder/bile duct cancer: A populationâ€based cohort study in <scp>J</scp> apan. International Journal of Cancer, 2017, 140, 1009-1019.	5.1	21
71	Helicobacter pylori infection, atrophic gastritis, and risk of pancreatic cancer: A population-based cohort study in a large Japanese population: the JPHC Study. Scientific Reports, 2019, 9, 6099.	3.3	21
72	Coping behaviors and suicide in the middle-aged and older Japanese general population: the Japan Public Health Center-based Prospective Study. Annals of Epidemiology, 2014, 24, 199-205.	1.9	20

#	Article	IF	CITATIONS
73	<i>CYP1A1</i> , <i>GSTM1</i> and <i>GSTT1</i> genetic polymorphisms and gastric cancer risk among Japanese: A nested caseâ€"control study within a largeâ€scale populationâ€based prospective study. International Journal of Cancer, 2016, 139, 759-768.	5.1	20
74	Dietary patterns and colorectal cancer risk in middle-aged adults: AÂlarge population-based prospective cohort study. Clinical Nutrition, 2018, 37, 1019-1026.	5.0	20
75	Prediagnostic circulating inflammation biomarkers and esophageal squamous cell carcinoma: A case–cohort study in Japan. International Journal of Cancer, 2020, 147, 686-691.	5.1	19
76	High-Negative Anti– <i>Helicobacter pylori</i> IgG Antibody Titers and Long-Term Risk of Gastric Cancer: Results from a Large-Scale Population-Based Cohort Study in Japan. Cancer Epidemiology Biomarkers and Prevention, 2020, 29, 420-426.	2.5	19
77	Fermented soy products intake and risk of cardiovascular disease and total cancer incidence: The Japan Public Health Center-based Prospective study. European Journal of Clinical Nutrition, 2021, 75, 954-968.	2.9	19
78	Cigarette smoking, alcohol drinking, and oral cavity and pharyngeal cancer in the Japanese: a population-based cohort study in Japan. European Journal of Cancer Prevention, 2018, 27, 171-179.	1.3	19
79	Coping strategies and cancer incidence and mortality: The Japan Public Health Center-based prospective study. Cancer Epidemiology, 2016, 40, 126-133.	1.9	18
80	Dietary fiber intake and risk of breast cancer defined by estrogen and progesterone receptor status: the Japan Public Health Center-based Prospective Study. Cancer Causes and Control, 2017, 28, 569-578.	1.8	18
81	Smoking, Alcohol Consumption, and Risks for Biliary Tract Cancer and Intrahepatic Bile Duct Cancer. Journal of Epidemiology, 2019, 29, 180-186.	2.4	18
82	Cruciferous vegetable intake and mortality in middle-aged adults: A prospective cohort study. Clinical Nutrition, 2019, 38, 631-643.	5.0	18
83	Plasma Isoflavones and Risk of Primary Liver Cancer in Japanese Women and Men with Hepatitis Virus Infection: A Nested Case–Control Study. Cancer Epidemiology Biomarkers and Prevention, 2015, 24, 532-537.	2.5	17
84	Evaluation of the degree of pancreatic fatty infiltration by area-based assessment of CT images: comparison with histopathology-based and CT attenuation index-based assessments. Japanese Journal of Radiology, 2016, 34, 667-676.	2.4	17
85	Effect of body-mass index on the risk of gastric cancer: A population-based cohort study in A Japanese population. Cancer Epidemiology, 2019, 63, 101622.	1.9	17
86	Lowâ€carbohydrate diet and risk of cancer incidence: The Japan Public Health Centerâ€based prospective study. Cancer Science, 2022, 113, 744-755.	3.9	17
87	Plasma adiponectin levels, ADIPOQ variants, and incidence of type 2 diabetes: A nested case-control study. Diabetes Research and Clinical Practice, 2017, 127, 254-264.	2.8	16
88	Female reproductive factors and risk of all-cause and cause-specific mortality among women: The Japan Public Health Center–based Prospective Study (JPHC study). Annals of Epidemiology, 2018, 28, 597-604.e6.	1.9	16
89	Genome-wide association meta-analysis and Mendelian randomization analysis confirm the influence of ALDH2 on sleep durationin the Japanese population. Sleep, 2019, 42, .	1.1	16
90	Association of BMI and height with the risk of endometrial cancer, overall and by histological subtype: a population-based prospective cohort study in Japan. European Journal of Cancer Prevention, 2019, 28, 196-202.	1.3	16

#	Article	IF	Citations
91	Development of a risk prediction model for lung cancer: The Japan Public Health Centerâ€based Prospective Study. Cancer Science, 2018, 109, 854-862.	3.9	15
92	Adult height and all-cause and cause-specific mortality in the Japan Public Health Center-based Prospective Study (JPHC). PLoS ONE, 2018, 13, e0197164.	2.5	15
93	Tuberculosis infection and lung adenocarcinoma: Mendelian randomization and pathway analysis of genome-wide association study data from never-smoking Asian women. Genomics, 2020, 112, 1223-1232.	2.9	15
94	The association between complete and partial non-response to psychosocial questions and suicide: the JPHC Study. European Journal of Public Health, 2015, 25, 424-430.	0.3	14
95	Alcohol consumption, genetic variants in the alcohol- and folate metabolic pathways and colorectal cancer risk: the JPHC Study. Scientific Reports, 2016, 6, 36607.	3.3	14
96	Body mass index change during adulthood and risk of oesophageal squamous-cell carcinoma in a Japanese population: the Japan Public Health (JPHC)-based prospective study. British Journal of Cancer, 2017, 117, 1715-1722.	6.4	14
97	Alcohol consumption and bladder cancer risk with or without the flushing response: The Japan Public Health Centerâ€based Prospective Study. International Journal of Cancer, 2017, 141, 2480-2488.	5.1	14
98	Coffee and green tea consumption and subsequent risk of acute myeloid leukemia and myelodysplastic syndromes in Japan. International Journal of Cancer, 2018, 142, 1130-1138.	5.1	14
99	Association between serum liver enzymes and allâ€cause mortality: The Japan Public Health Centerâ€based Prospective Study. Liver International, 2019, 39, 1566-1576.	3.9	14
100	Variations in the estimated intake of acrylamide from food in the Japanese population. Nutrition Journal, 2020, 19, 17.	3.4	14
101	ldentification of a novel uterine leiomyoma GWAS locus in a Japanese population. Scientific Reports, 2020, 10, 1197.	3.3	14
102	Relationship between unhealthy sleep status and dry eye symptoms in a Japanese population: The JPHC-NEXT study. Ocular Surface, 2021, 21, 306-312.	4.4	14
103	Fermented and nonfermented soy foods and the risk of breast cancer in a Japanese populationâ€based cohort study. Cancer Medicine, 2021, 10, 757-771.	2.8	14
104	Risk of thyroid cancer in relation to height, weight, and body mass index in Japanese individuals: a population-based cohort study. Cancer Medicine, 2018, 7, 2200-2210.	2.8	13
105	Plasma and tumoral glypicanâ€3 levels are correlated in patients with hepatitis C virusâ€related hepatocellular carcinoma. Cancer Science, 2020, 111, 334-342.	3.9	13
106	Association of <i>Escherichia coli</i> containing polyketide synthase in the gut microbiota with colorectal neoplasia in Japan. Cancer Science, 2022, 113, 277-286.	3.9	13
107	Glycemic index and glycemic load and risk of colorectal cancer: a population-based cohort study (JPHC Study). Cancer Causes and Control, 2016, 27, 583-593.	1.8	12
108	Smoking and subsequent risk of leukemia in Japan: The Japan Public Health Center-based Prospective Study. Journal of Epidemiology, 2017, 27, 305-310.	2.4	12

#	Article	IF	CITATIONS
109	Menstrual and reproductive factors in the risk of thyroid cancer in Japanese women: the Japan Public Health Center-Based Prospective Study. European Journal of Cancer Prevention, 2018, 27, 361-369.	1.3	11
110	Changes in Smoking Status and Mortality From All Causes and Lung Cancer: A Longitudinal Analysis of a Population-based Study in Japan. Journal of Epidemiology, 2019, 29, 11-17.	2.4	11
111	Fruit and vegetable intake and pancreatic cancer risk in a populationâ€based cohort study in Japan. International Journal of Cancer, 2019, 144, 1858-1866.	5.1	11
112	Cross-Sectional Association Between Employment Status and Self-Rated Health Among Middle-Aged Japanese Women: The Influence of Socioeconomic Conditions and Work-Life Conflict. Journal of Epidemiology, 2020, 30, 396-403.	2.4	11
113	Occupational sitting time and subsequent risk of cancer: The Japan Public Health Centerâ€based Prospective Study. Cancer Science, 2020, 111, 974-984.	3.9	11
114	Fat mass and obesity-associated gene polymorphisms, pre-diagnostic plasma adipokine levels and the risk of colorectal cancer: The Japan Public Health Center-based Prospective Study. PLoS ONE, 2020, 15, e0229005.	2.5	11
115	Associations between changes in fruit and vegetable consumption and weight change in Japanese adults. European Journal of Nutrition, 2021, 60, 217-227.	3.9	11
116	Coffee Consumption and Lung Cancer Risk: The Japan Public Health Center-Based Prospective Study. Journal of Epidemiology, 2018, 28, 207-213.	2.4	10
117	Comparison between the impact of fermented and unfermented soy intake on the risk of liver cancer: the JPHC Study. European Journal of Nutrition, 2021, 60, 1389-1401.	3.9	10
118	Reproductive Factors and Lung Cancer Risk among Never-Smoking Japanese Women with 21 Years of Follow-Up: A Cohort Study. Cancer Epidemiology Biomarkers and Prevention, 2021, 30, 1185-1192.	2.5	10
119	Association between meat intake and mortality due to all-cause and major causes of death in a Japanese population. PLoS ONE, 2020, 15, e0244007.	2.5	10
120	Long-term exposure to fine particle matter and all-cause mortality and cause-specific mortality in Japan: the JPHC Study. BMC Public Health, 2022, 22, 466.	2.9	10
121	Metabolome analysis for pancreatic cancer risk in nested caseâ€control study: Japan Public Health Centerâ€based prospective Study. Cancer Science, 2018, 109, 1672-1681.	3.9	9
122	Identification of two novel breast cancer loci through large-scale genome-wide association study in the Japanese population. Scientific Reports, 2019, 9, 17332.	3.3	9
123	The Association Between Habitual Sleep Duration and Mortality According to Sex and Age: The Japan Public Health Center-based Prospective Study. Journal of Epidemiology, 2021, 31, 109-118.	2.4	9
124	Working cancer survivors' physical and mental characteristics compared to cancer-free workers in Japan: a nationwide general population-based study. Journal of Cancer Survivorship, 2021, 15, 912-921.	2.9	9
125	Association between C-reactive protein and risk of overall and 18 site-specific cancers in a Japanese case-cohort. British Journal of Cancer, 2022, 126, 1481-1489.	6.4	9
126	Trends in cancer prognosis in a population-based cohort survey: Can recent advances in cancer therapy affect the prognosis?. Cancer Epidemiology, 2015, 39, 97-103.	1.9	8

#	Article	IF	CITATIONS
127	Association of plasma C-reactive protein level with the prevalence of colorectal adenoma: the Colorectal Adenoma Study in Tokyo. Scientific Reports, 2017, 7, 4456.	3.3	8
128	Genome-wide association study (GWAS) of ovarian cancer in Japanese predicted regulatory variants in 22q13.1. PLoS ONE, 2018, 13, e0209096.	2.5	8
129	Higher Dietary Non-enzymatic Antioxidant Capacity Is Associated with Decreased Risk of All-Cause and Cardiovascular Disease Mortality in Japanese Adults. Journal of Nutrition, 2019, 149, 1967-1976.	2.9	8
130	Circulating Inflammation Markers and Risk of Gastric and Esophageal Cancers: A Case–Cohort Study Within the Japan Public Health Center–Based Prospective Study. Cancer Epidemiology Biomarkers and Prevention, 2019, 28, 829-832.	2.5	8
131	Soy food and isoflavones are not associated with changes in serum lipids and glycohemoglobin concentrations among Japanese adults: a cohort study. European Journal of Nutrition, 2020, 59, 2075-2087.	3.9	8
132	Doneness preferences, meat and meat-derived heterocyclic amines intake, and N-acetyltransferase 2 polymorphisms: association with colorectal adenoma in Japanese Brazilians. European Journal of Cancer Prevention, 2020, 29, 7-14.	1.3	8
133	Inclusion of a geneâ€environment interaction between alcohol consumption and the aldehyde dehydrogenase 2 genotype in a risk prediction model for upper aerodigestive tract cancer in Japanese men. Cancer Science, 2020, 111, 3835-3844.	3.9	8
134	Association of dietary intakes of vitamin B12, vitamin B6, folate, and methionine with the risk of esophageal cancer: the Japan Public Health Center-based (JPHC) prospective study. BMC Cancer, 2021, 21, 982.	2.6	8
135	Sugary drink consumption and risk of kidney and bladder cancer in Japanese adults. Scientific Reports, 2021, 11, 21701.	3.3	8
136	Comparison of weighed food record procedures for the reference methods in two validation studies of food frequency questionnaires. Journal of Epidemiology, 2017, 27, 331-337.	2.4	7
137	Association between dietary sugar intake and colorectal adenoma among cancer screening examinees in Japan. Cancer Science, 2020, 111, 3862-3872.	3.9	7
138	Sugary Drink Consumption and Subsequent Colorectal Cancer Risk: The Japan Public Health Center–Based Prospective Cohort Study. Cancer Epidemiology Biomarkers and Prevention, 2021, 30, 782-788.	2.5	7
139	Alcohol consumption, tobacco smoking, and subsequent risk of renal cell carcinoma: The JPHC study. Cancer Science, 2021, 112, 5068-5077.	3.9	7
140	Fruit and vegetable consumption and risk of esophageal cancer in the Asian region: a systematic review and meta-analysis. Esophagus, 2022, 19, 27-38.	1.9	7
141	Cruciferous vegetable intake and colorectal cancer risk: Japan public health center-based prospective study. European Journal of Cancer Prevention, 2019, 28, 420-427.	1.3	6
142	Family history of cancer and subsequent risk of cancer: A largeâ€scale populationâ€based prospective study in Japan. International Journal of Cancer, 2020, 147, 331-337.	5.1	6
143	Soy and isoflavone consumption and subsequent risk of prostate cancer mortality: the Japan Public Health Center-based Prospective Study. International Journal of Epidemiology, 2020, 49, 1553-1561.	1.9	6
144	Metabolic Syndrome, Physical Activity, and Inflammation: A Cross-Sectional Analysis of 110 Circulating Biomarkers in Japanese Adults. Cancer Epidemiology Biomarkers and Prevention, 2020, 29, 1639-1646.	2.5	6

#	Article	IF	Citations
145	Association Between Birth Weight and Risk of Pregnancy-Induced Hypertension and Gestational Diabetes in Japanese Women: JPHC-NEXT Study. Journal of Epidemiology, 2022, 32, 168-173.	2.4	6
146	Functional annotation of the 2q35 breast cancer risk locus implicates a structural variant in influencing activity of a long-range enhancer element. American Journal of Human Genetics, 2021, 108, 1190-1203.	6.2	6
147	Prediagnostic circulating inflammation-related biomarkers and gastric cancer: A case-cohort study in Japan. Cytokine, 2021, 144, 155558.	3.2	6
148	Food frequency questionnaire reproducibility for middle-aged and elderly Japanese. Asia Pacific Journal of Clinical Nutrition, 2019, 28, 362-370.	0.4	6
149	Meat consumption and gastric cancer risk: The Japan Public Health Center-based Prospective Study. American Journal of Clinical Nutrition, 2021, , .	4.7	6
150	Coffee and Green Tea Consumption and Subsequent Risk of Malignant Lymphoma and Multiple Myeloma in Japan: The Japan Public Health Center-based Prospective Study. Cancer Epidemiology Biomarkers and Prevention, 2017, 26, 1352-1356.	2.5	5
151	Female reproductive factors and risk of lymphoid neoplasm: The Japan Public Health Centerâ€based Prospective Study. Cancer Science, 2019, 110, 1442-1452.	3.9	5
152	Plasma Câ€peptide and glycated albumin and subsequent risk of cancer: From a large prospective caseâ€cohort study in Japan. International Journal of Cancer, 2019, 144, 718-729.	5.1	5
153	Soy Intake and Colorectal Cancer Risk: Results from a Pooled Analysis of Prospective Cohort Studies Conducted in China and Japan. Journal of Nutrition, 2020, 150, 2442-2450.	2.9	5
154	Estimation of the performance of a risk prediction model for gastric cancer occurrence in Japan: Evidence from a small external population. Cancer Epidemiology, 2020, 67, 101766.	1.9	5
155	Dietary glycemic index, glycemic load, and endometrial cancer risk: The Japan Public Health Centerâ€based Prospective Study. Cancer Science, 2021, 112, 3682-3690.	3.9	5
156	Association of sugary drink consumption with all-cause and cause-specific mortality: the Japan Public Health Center-based Prospective Study. Preventive Medicine, 2021, 148, 106561.	3.4	5
157	Exploring predictive biomarkers from clinical genome-wide association studies via multidimensional hierarchical mixture models. European Journal of Human Genetics, 2019, 27, 140-149.	2.8	4
158	Relationship between Meat/Fish Consumption and Biliary Tract Cancer: The Japan Public Health Center–Based Prospective Study. Cancer Epidemiology Biomarkers and Prevention, 2020, 29, 95-102.	2.5	4
159	Soy Food Intake and Pancreatic Cancer Risk: The Japan Public Health Center–based Prospective Study. Cancer Epidemiology Biomarkers and Prevention, 2020, 29, 1214-1221.	2.5	4
160	Body Mass Index, Height, Weight Change, and Subsequent Lung Cancer Risk: The Japan Public Health Center–Based Prospective Study. Cancer Epidemiology Biomarkers and Prevention, 2021, 30, 1708-1716.	2.5	4
161	A Personal Breast Cancer Risk Stratification Model Using Common Variants and Environmental Risk Factors in Japanese Females. Cancers, 2021, 13, 3796.	3.7	4
162	Vegetable and fruit intake and the risk of bladder cancer: Japan Public Health Center-based prospective study. British Journal of Cancer, 2022, 126, 1647-1658.	6.4	4

#	Article	IF	CITATIONS
163	The association between plasma C-peptide concentration and the risk of prostate cancer: a nested case–control study within a Japanese population-based prospective study. European Journal of Cancer Prevention, 2018, 27, 461-467.	1.3	3
164	Female reproductive factors and risk of external causes of death among women: The Japan Public Health Center-based Prospective Study (JPHC Study). Scientific Reports, 2019, 9, 14329.	3.3	3
165	Lack of social support and social trust as potential risk factors for dry eye disease: JPHC-NEXT study. Ocular Surface, 2019, 17, 278-284.	4.4	3
166	Impact of alcohol drinking on cancer risk with consideration of flushing response: The Japan Public Health Center-based Prospective Study Cohort (JPHC study). Preventive Medicine, 2020, 133, 106026.	3.4	3
167	Effectiveness of Screening Using Fecal Occult Blood Testing and Colonoscopy on the Risk of Colorectal Cancer: The Japan Public Health Center-based Prospective Study. Journal of Epidemiology, 2023, 33, 91-100.	2.4	3
168	Apolipoprotein A2 Isoforms in Relation to the Risk of Myocardial Infarction: A Nested Case-Control Analysis in the JPHC Study. Journal of Atherosclerosis and Thrombosis, 2021, 28, 483-490.	2.0	3
169	Long-term Response of <i>Helicobacter pylori</i> Antibody Titer After Eradication Treatment in Middle-aged Japanese: JPHC-NEXT Study. Journal of Epidemiology, 2023, 33, 1-7.	2.4	3
170	The potential for reducing alcohol consumption to prevent esophageal cancer morbidity in Asian heavy drinkers: a systematic review and meta-analysis. Esophagus, 2021, 19, 39.	1.9	3
171	Association between coffee consumption and risk of prostate cancer in Japanese men: a population-based cohort study in Japan. Cancer Epidemiology Biomarkers and Prevention, 2021, , cebp.0484.2021.	2.5	3
172	Association of B Vitamins and Methionine Intake with the Risk of Gastric Cancer: The Japan Public Health Center–based Prospective Study. Cancer Prevention Research, 2022, 15, 101-110.	1.5	3
173	Association between Meat, Fish, and Fatty Acid Intake and Non-Hodgkin Lymphoma Incidence: The Japan Public Health Center–Based Prospective Study. Journal of Nutrition, 2022, 152, 1895-1906.	2.9	3
174	Germline HOXB13 mutations p.G84E and p.R217C do not confer an increased breast cancer risk. Scientific Reports, 2020, 10, 9688.	3.3	2
175	Risk of stroke in cancer survivors using a propensity score-matched cohort analysis. Scientific Reports, 2021, 11, 5599.	3.3	2
176	Dietary glycemic index, glycemic load and mortality: Japan Public Health Center-based prospective study. European Journal of Nutrition, 2021, 60, 4607-4620.	3.9	2
177	Circulating Inflammation Markers and Pancreatic Cancer Risk: A Prospective Case-Cohort Study in Japan. Cancer Epidemiology Biomarkers and Prevention, 2022, 31, 236-241.	2.5	2
178	Dietary fibre intake is associated with reduced risk of lung cancer: a Japan public health centre-based prospective study (JPHC). International Journal of Epidemiology, 2022, 51, 1142-1152.	1.9	2
179	Body mass index and height in relation to brain tumor risk in a Japanese population. Annals of Epidemiology, 2020, 51, 1-6.	1.9	1
180	Validity of dietary isothiocyanate intake estimates from a food frequency questionnaire using 24 h urinary isothiocyanate excretion as an objective biomarker: the JPHC-NEXT protocol area. European Journal of Clinical Nutrition, 2021, , .	2.9	1

#	Article	IF	Citations
181	Total, animal, and plant protein intake and pneumonia mortality in the Japan Public Health Center–based Prospective Study. American Journal of Clinical Nutrition, 2022, 115, 781-789.	4.7	1
182	Relevance of the MHC region for breast cancer susceptibility in Asians. Breast Cancer, 2022, 29, 869-879.	2.9	1
183	Association of Plasma Iron Status with Subsequent Risk of Total and Site-Specific Cancer: A Large Case–Cohort Study within JPHC Study. Cancer Prevention Research, 2022, 15, 669-678.	1.5	1
184	Low <i>MICA </i> gene expression confers an increased risk of Graves' disease: a Mendelian randomization study. Thyroid, 2021, , .	4.5	0
185	COT-6 Body mass index and height in relation to brain tumor risk in a Japanese population. Neuro-Oncology Advances, 2021, 3, vi29-vi29.	0.7	0
186	Title is missing!. , 2020, 15, e0244007.		0
187	Title is missing!. , 2020, 15, e0244007.		0
188	Title is missing!. , 2020, 15, e0244007.		0
189	Title is missing!. , 2020, 15, e0244007.		0