

# Resnati Massimo

## List of Publications by Year in descending order

Source: <https://exaly.com/author-pdf/4695792/publications.pdf>

Version: 2024-02-01

29  
papers

3,033  
citations

394286

19  
h-index

477173

29  
g-index

29  
all docs

29  
docs citations

29  
times ranked

3459  
citing authors

#	ARTICLE	IF	CITATIONS
1	A novel endothelial-specific membrane protein is a marker of cell-cell contacts.. Journal of Cell Biology, 1992, 118, 1511-1522.	2.3	602
2	The High Mobility Group (Hmg) Boxes of the Nuclear Protein Hmg1 Induce Chemotaxis and Cytoskeleton Reorganization in Rat Smooth Muscle Cells. Journal of Cell Biology, 2001, 152, 1197-1206.	2.3	435
3	The fibrinolytic receptor for urokinase activates the G protein-coupled chemotactic receptor FPRL1/LXA4R. Proceedings of the National Academy of Sciences of the United States of America, 2002, 99, 1359-1364.	3.3	345
4	The role of integrins in the maintenance of endothelial monolayer integrity.. Journal of Cell Biology, 1991, 112, 479-490.	2.3	278
5	A urokinase-sensitive region of the human urokinase receptor is responsible for its chemotactic activity. EMBO Journal, 1997, 16, 7279-7286.	3.5	210
6	Cytotoxicity of some catalysts commonly used in the synthesis of copolymers for biomedical use. Journal of Materials Science: Materials in Medicine, 1994, 5, 393-396.	1.7	189
7	Src-Dependence and Pertussis-Toxin Sensitivity of Urokinase Receptor-Dependent Chemotaxis and Cytoskeleton Reorganization in Rat Smooth Muscle Cells. Blood, 1999, 94, 649-662.	0.6	111
8	Domain 2 of the Urokinase Receptor Contains an Integrin-interacting Epitope with Intrinsic Signaling Activity. Journal of Biological Chemistry, 2005, 280, 24792-24803.	1.6	103
9	Urokinase/urokinase receptor and vitronectin/ $\alpha$ v $\beta$ 3 integrin induce chemotaxis and cytoskeleton reorganization through different signaling pathways. Oncogene, 2001, 20, 2032-2043.	2.6	100
10	A plastic SQSTM1/p62-dependent autophagic reserve maintains proteostasis and determines proteasome inhibitor susceptibility in multiple myeloma cells. Autophagy, 2015, 11, 1161-1178.	4.3	82
11	Metalloproteases Cleave the Urokinase-Type Plasminogen Activator Receptor in the D1-D2 Linker Region and Expose Epitopes not Present in the intact Soluble Receptor. Thrombosis and Haemostasis, 2002, 88, 298-306.	1.8	77
12	The amyloidogenic light chain is a stressor that sensitizes plasma cells to proteasome inhibitor toxicity. Blood, 2017, 129, 2132-2142.	0.6	70
13	The soluble D2D388-274 fragment of the urokinase receptor inhibits monocyte chemotaxis and integrin-dependent cell adhesion. Journal of Cell Science, 2004, 117, 2909-2916.	1.2	69
14	PAI-1 inhibits urokinase-induced chemotaxis by internalizing the urokinase receptor. FEBS Letters, 2001, 505, 249-254.	1.3	63
15	The urokinase receptor: Structure, regulation and inhibitor-mediated internalization. Fibrinolysis, 1994, 8, 182-188.	0.5	56
16	Myb-Binding Protein 1A (MYBBP1A) Is Essential for Early Embryonic Development, Controls Cell Cycle and Mitosis, and Acts as a Tumor Suppressor. PLoS ONE, 2012, 7, e39723.	1.1	43
17	uPA/uPAR System Is Active in Immature Dendritic Cells Derived from CD14+CD34+ Precursors and Is Down-Regulated upon Maturation. Journal of Immunology, 2000, 164, 712-718.	0.4	31
18	Endothelial integrins and their role in maintaining the integrity of the vessel wall. Kidney International, 1993, 43, 61-65.	2.6	28

#	ARTICLE	IF	CITATIONS
19	The Interaction of the Tumor Suppressor FAM46C with p62 and FNDC3 Proteins Integrates Protein and Secretory Homeostasis. <i>Cell Reports</i> , 2020, 32, 108162.	2.9	24
20	Oncogenic HoxB7 requires TALE cofactors and is inactivated by a dominant-negative Pbx1 mutant in a cell-specific manner. <i>Cancer Letters</i> , 2008, 266, 144-155.	3.2	23
21	HIV-1 Infected Lymphoid Organs Upregulate Expression and Release of the Cleaved Form of uPAR That Modulates Chemotaxis and Virus Expression. <i>PLoS ONE</i> , 2013, 8, e70606.	1.1	18
22	Autophagy mediates epithelial cancer chemoresistance by reducing p62/SQSTM1 accumulation. <i>PLoS ONE</i> , 2018, 13, e0201621.	1.1	15
23	Biosynthesis and apical localization of the urokinase receptor in polarized MDCK epithelial cells. <i>FEBS Letters</i> , 1995, 369, 207-211.	1.3	13
24	Specific immunofluorimetric assay detecting the chemotactic epitope of the urokinase receptor (uPAR). <i>Journal of Immunological Methods</i> , 2006, 308, 192-202.	0.6	13
25	The Autophagic Process Occurs in Human Bone Metastasis and Implicates Molecular Mechanisms Differently Affected by Rab5a in the Early and Late Stages. <i>International Journal of Molecular Sciences</i> , 2016, 17, 443.	1.8	12
26	A multimodal molecular imaging approach targeting urokinase plasminogen activator receptor for the diagnosis, resection and surveillance of urothelial cell carcinoma. <i>European Journal of Cancer</i> , 2021, 146, 11-20.	1.3	8
27	Side-by-Side Comparison of uPAR-Targeting Optical Imaging Antibodies and Antibody Fragments for Fluorescence-Guided Surgery of Solid Tumors. <i>Molecular Imaging and Biology</i> , 2021, , 1.	1.3	6
28	Requirement of the enzymatic and signaling activities of plasmin for phorbol-ester-induced scattering of colon cancer cells. <i>Experimental Cell Research</i> , 2006, 312, 2203-2213.	1.2	5
29	Src-Dependence and Pertussis-Toxin Sensitivity of Urokinase Receptor-Dependent Chemotaxis and Cytoskeleton Reorganization in Rat Smooth Muscle Cells. <i>Blood</i> , 1999, 94, 649-662.	0.6	4