

Guntram Suske

List of Publications by Year in Descending Order

Source: <https://exaly.com/author-pdf/4697027/guntram-suske-publications-by-year.pdf>

Version: 2024-04-27

This document has been generated based on the publications and citations recorded by exaly.com. For the latest version of this publication list, visit the link given above.

The third column is the impact factor (IF) of the journal, and the fourth column is the number of citations of the article.

35
papers

3,335
citations

25
h-index

36
g-index

36
ext. papers

3,517
ext. citations

7.1
avg, IF

4.86
L-index

#	Paper	IF	Citations
35	MGA, L3MBTL2 and E2F6 determine genomic binding of the non-canonical Polycomb repressive complex PRC1.6. <i>PLoS Genetics</i> , 2018 , 14, e1007193	6	54
34	Transcription factor Sp2 potentiates binding of the TALE homeoproteins Pbx1:Prep1 and the histone-fold domain protein Nf-y to composite genomic sites. <i>Journal of Biological Chemistry</i> , 2018 , 293, 19250-19262	5.4	4
33	NF-Y and SP transcription factors - New insights in a long-standing liaison. <i>Biochimica Et Biophysica Acta - Gene Regulatory Mechanisms</i> , 2017 , 1860, 590-597	6	20
32	Sp2 is the only glutamine-rich specificity protein with minor impact on development and differentiation in myelinating glia. <i>Journal of Neurochemistry</i> , 2017 , 140, 245-256	6	2
31	Zinc finger independent genome-wide binding of Sp2 potentiates recruitment of histone-fold protein Nf-y distinguishing it from Sp1 and Sp3. <i>PLoS Genetics</i> , 2015 , 11, e1005102	6	29
30	Sp1/Sp3 transcription factors regulate hallmarks of megakaryocyte maturation and platelet formation and function. <i>Blood</i> , 2015 , 125, 1957-67	2.2	45
29	SUMOylation of the polycomb group protein L3MBTL2 facilitates repression of its target genes. <i>Nucleic Acids Research</i> , 2014 , 42, 3044-58	20.1	19
28	Ligand binding reduces SUMOylation of the peroxisome proliferator-activated receptor [[PPAR]] activation function 1 (AF1) domain. <i>PLoS ONE</i> , 2013 , 8, e66947	3.7	23
27	Genome-wide localization and expression profiling establish Sp2 as a sequence-specific transcription factor regulating vitally important genes. <i>Nucleic Acids Research</i> , 2012 , 40, 7844-57	20.1	27
26	Differential regulation of sense and antisense promoter activity at the Csf1R locus in B cells by the transcription factor PAX5. <i>Experimental Hematology</i> , 2011 , 39, 730-40.e1-2	3.1	4
25	Specificity protein 2 (Sp2) is essential for mouse development and autonomous proliferation of mouse embryonic fibroblasts. <i>PLoS ONE</i> , 2010 , 5, e9587	3.7	18
24	Epigenetic silencing of spermatocyte-specific and neuronal genes by SUMO modification of the transcription factor Sp3. <i>PLoS Genetics</i> , 2010 , 6, e1001203	6	30
23	SUMO-modified Sp3 represses transcription by provoking local heterochromatic gene silencing. <i>EMBO Reports</i> , 2008 , 9, 899-906	6.5	68
22	Identification of SUMO-dependent chromatin-associated transcriptional repression components by a genome-wide RNAi screen. <i>Molecular Cell</i> , 2008 , 29, 742-54	17.6	93
21	Sp1/Sp3 compound heterozygous mice are not viable: impaired erythropoiesis and severe placental defects. <i>Developmental Dynamics</i> , 2007 , 236, 2235-44	2.9	49
20	Transcription factor Sp3 knockout mice display serious cardiac malformations. <i>Molecular and Cellular Biology</i> , 2007 , 27, 8571-82	4.8	46
19	Mammalian SP/KLF transcription factors: bring in the family. <i>Genomics</i> , 2005 , 85, 551-6	4.3	287

18	Complexity of translationally controlled transcription factor Sp3 isoform expression. <i>Journal of Biological Chemistry</i> , 2004 , 279, 42095-105	5.4	89
17	E2F and Sp1/Sp3 Synergize but are not sufficient to activate the MYCN gene in neuroblastomas. <i>Journal of Biological Chemistry</i> , 2004 , 279, 5110-7	5.4	39
16	Impaired hematopoiesis in mice lacking the transcription factor Sp3. <i>Blood</i> , 2003 , 102, 858-66	2.2	38
15	Transcription factor Sp3 is silenced through SUMO modification by PIAS1. <i>EMBO Journal</i> , 2002 , 21, 5206-15	15	222
14	Complex phenotype of mice homozygous for a null mutation in the Sp4 transcription factor gene. <i>Genes To Cells</i> , 2001 , 6, 689-97	2.3	49
13	Transcription factor Sp3 is regulated by acetylation. <i>Nucleic Acids Research</i> , 2001 , 29, 4994-5000	20.1	119
12	Impaired ossification in mice lacking the transcription factor Sp3. <i>Mechanisms of Development</i> , 2001 , 106, 77-83	1.7	95
11	Characterization and promoter analysis of the mouse gene for transcription factor Sp4. <i>Gene</i> , 2001 , 264, 19-27	3.8	21
10	Transcription factor Sp3 is essential for post-natal survival and late tooth development. <i>EMBO Journal</i> , 2000 , 19, 655-61	13	163
9	Transient transfection of Schneider cells in the study of transcription factors. <i>Methods in Molecular Biology</i> , 2000 , 130, 175-87	1.4	28
8	A tale of three fingers: the family of mammalian Sp/XKLF transcription factors. <i>Nucleic Acids Research</i> , 1999 , 27, 2991-3000	20.1	535
7	Combinatorial action of HNF3 and Sp family transcription factors in the activation of the rabbit uteroglobin/CC10 promoter. <i>Journal of Biological Chemistry</i> , 1998 , 273, 9821-8	5.4	61
6	Human Sp3 transcriptional regulator gene (SP3) maps to chromosome 2q31. <i>Genomics</i> , 1996 , 37, 410-2	4.3	15
5	High Sp1/Sp3 ratios in epithelial cells during epithelial differentiation and cellular transformation correlate with the activation of the HPV-16 promoter. <i>Virology</i> , 1996 , 224, 281-91	3.6	133
4	Functional analyses of the transcription factor Sp4 reveal properties distinct from Sp1 and Sp3. <i>Journal of Biological Chemistry</i> , 1995 , 270, 24989-94	5.4	179
3	Sp1 trans-activation of cell cycle regulated promoters is selectively repressed by Sp3. <i>Biochemistry</i> , 1995 , 34, 16503-8	3.2	179
2	Human Sp4 transcription factor gene (SP4) maps to chromosome 7p15. <i>Genomics</i> , 1995 , 26, 631-3	4.3	7
1	Cloning by recognition site screening of two novel GT box binding proteins: a family of Sp1 related genes. <i>Nucleic Acids Research</i> , 1992 , 20, 5519-25	20.1	545

