

Zhang Zhang

List of Publications by Year in descending order

Source: <https://exaly.com/author-pdf/4704393/publications.pdf>

Version: 2024-02-01

34
papers

633
citations

623734

14
h-index

610901

24
g-index

36
all docs

36
docs citations

36
times ranked

1094
citing authors

#	ARTICLE	IF	CITATIONS
1	Folate Receptor β Associated With Triple-Negative Breast Cancer and Poor Prognosis. Archives of Pathology and Laboratory Medicine, 2014, 138, 890-895.	2.5	110
2	miR-483-5p determines mitochondrial fission and cisplatin sensitivity in tongue squamous cell carcinoma by targeting FIS1. Cancer Letters, 2015, 362, 183-191.	7.2	67
3	Mitochondrial fission determines cisplatin sensitivity in tongue squamous cell carcinoma through the BRCA1-miR-593-5p-MFF axis. Oncotarget, 2015, 6, 14885-14904.	1.8	50
4	Chemotherapy-Induced Long Non-coding RNA 1 Promotes Metastasis and Chemo-Resistance of TSCC via the Wnt/ β -Catenin Signaling Pathway. Molecular Therapy, 2018, 26, 1494-1508.	8.2	47
5	Consistent Amplification of FRS2 and MDM2 in Low-grade Osteosarcoma. American Journal of Surgical Pathology, 2018, 42, 1143-1155.	3.7	34
6	Automated quantitative analysis of Ki-67 staining and HE images recognition and registration based on whole tissue sections in breast carcinoma. Diagnostic Pathology, 2020, 15, 65.	2.0	26
7	CD49f Can Act as a Biomarker for Local or Distant Recurrence in Breast Cancer. Journal of Breast Cancer, 2017, 20, 142.	1.9	24
8	Infantile hepatic hemangioendothelioma: A clinicopathologic study in a Chinese population. World Journal of Gastroenterology, 2010, 16, 4549.	3.3	22
9	ALDH+/CD44+ cells in breast cancer are associated with worse prognosis and poor clinical outcome. Experimental and Molecular Pathology, 2016, 100, 145-150.	2.1	22
10	The Presence of EpCAM-/CD49f+Cells in Breast Cancer Is Associated with a Poor Clinical Outcome. Journal of Breast Cancer, 2015, 18, 242.	1.9	21
11	Amplification of <i>FRS2</i> in atypical lipomatous tumour/well-differentiated liposarcoma and de-differentiated liposarcoma: a clinicopathological and genetic study of 146 cases. Histopathology, 2018, 72, 1145-1155.	2.9	20
12	Pathological features and clinical outcomes of breast cancer according to levels of oestrogen receptor expression. Histopathology, 2014, 65, 508-516.	2.9	18
13	MiR-4653-3p and its target gene FRS2 are prognostic biomarkers for hormone receptor positive breast cancer patients receiving tamoxifen as adjuvant endocrine therapy. Oncotarget, 2016, 7, 61166-61182.	1.8	18
14	Hormone Receptor and Human Epidermal Growth Factor Receptor 2 Detection in Invasive Breast Carcinoma: A Retrospective Study of 12,467 Patients From 19 Chinese Representative Clinical Centers. Clinical Breast Cancer, 2020, 20, e65-e74.	2.4	18
15	Application of <i>COL1A1</i> PDGFB fusion gene detection by fluorescence <i>in situ</i> hybridization in biopsy tissue of dermatofibrosarcoma protuberans. Journal of Dermatology, 2017, 44, 798-802.	1.2	15
16	Genomic Pathology and Biomarkers in Breast Cancer. Critical Reviews in Oncogenesis, 2017, 22, 411-426.	0.4	12
17	Primary pure carcinoid tumors of the testis: Clinicopathological and immunophenotypical characteristics of 11 cases. Oncology Letters, 2015, 9, 2017-2022.	1.8	11
18	A novel sclerosing atypical lipomatous tumor/well-differentiated liposarcoma in a 7-year-old girl: report of a case with molecular confirmation. Human Pathology, 2018, 71, 41-46.	2.0	11

#	ARTICLE	IF	CITATIONS
19	Liposarcoma in children and young adults: a clinicopathologic and molecular study of 23 cases in one of the largest institutions of China. <i>Virchows Archiv Fur Pathologische Anatomie Und Physiologie Und Fur Klinische Medizin</i> , 2021, 479, 537-549.	2.8	10
20	External quality assessment (EQA) program for the immunohistochemical detection of ER, PR and Ki-67 in breast cancer: results of an interlaboratory reproducibility ring study in China. <i>BMC Cancer</i> , 2019, 19, 978.	2.6	9
21	Prevalence of protein C receptor (PROCR) is associated with inferior clinical outcome in Breast invasive ductal carcinoma. <i>Pathology Research and Practice</i> , 2017, 213, 1173-1179.	2.3	8
22	Synovial sarcoma showing loss of a green signal in SS18 fluorescence in situ hybridization: a clinicopathological and molecular study of 12 cases. <i>Virchows Archiv Fur Pathologische Anatomie Und Physiologie Und Fur Klinische Medizin</i> , 2017, 471, 799-807.	2.8	8
23	The differences of clinicopathologic characteristics among subgroups of reclassified HER2 fluorescence in situ hybridization (FISH) according to the ASCO/CAP 2018 breast cancer HER2 testing guidelines. <i>Journal of Clinical Pathology</i> , 2020, 73, 283-290.	2.0	8
24	Clinicopathologic characteristics of HER2-positive metaplastic squamous cell carcinoma of the breast. <i>Journal of Clinical Pathology</i> , 2022, 75, 18-23.	2.0	7
25	Neoadjuvant pyrotinib plus trastuzumab and nab-paclitaxel for HER2-positive early or locally advanced breast cancer: an exploratory phase II trial. <i>Gland Surgery</i> , 2022, 11, 216-225.	1.1	7
26	Traumatic neuroma in mastectomy scar. <i>Medicine (United States)</i> , 2019, 98, e15142.	1.0	6
27	Primary cardiac dedifferentiated liposarcoma with homologous and heterologous differentiation: a case report. <i>International Journal of Clinical and Experimental Pathology</i> , 2015, 8, 9662-6.	0.5	6
28	An Advanced Automated Image Analysis Model for Scoring of ER, PR, HER-2 and Ki-67 in Breast Carcinoma. <i>IEEE Access</i> , 2021, 9, 108441-108451.	4.2	5
29	A decision tree-based prediction model for fluorescence <i>in situ</i> hybridization <i>HER2</i> gene status in HER2 immunohistochemistry-2+ breast cancers: a 2538-case multicenter study on consecutive surgical specimens. <i>Journal of Cancer</i> , 2018, 9, 2327-2333.	2.5	4
30	Genomic landscape of secretory carcinoma of the breast with axillary lymph node metastasis. <i>Pathology Research and Practice</i> , 2022, 231, 153790.	2.3	4
31	High expression of microRNA20b is associated with malignant clinicopathological features and poor prognosis in breast phyllodes tumor. <i>International Journal of Clinical Oncology</i> , 2020, 25, 2025-2034.	2.2	2
32	Atypical spindle cell lipomatous tumor/atypical pleomorphic lipomatous tumor and atypical mammary-type myofibroblastoma. <i>Chinese Medical Journal</i> , 2021, Publish Ahead of Print, 2370-2372.	2.3	1
33	Low-grade fibromyxoid sarcoma: a clinicopathologic and molecular study of 10 genetically confirmed cases. <i>International Journal of Clinical and Experimental Pathology</i> , 2018, 11, 5860-5868.	0.5	1
34	Biomarker profile of invasive lobular carcinoma: pleomorphic versus classic subtypes, clinicopathological characteristics and prognosis analyses. <i>Breast Cancer Research and Treatment</i> , 0, , ,	2.5	1