
Martin Wermke

List of Publications by Year in descending order

Source: https://exaly.com/author-pdf/4714926/publications.pdf Version: 2024-02-01



#	Article	IF	CITATIONS
1	Comparison of FACS and PCR for Detection of BCMA-CAR-T Cells. International Journal of Molecular Sciences, 2022, 23, 903.	1.8	7
2	Using stroma-anchoring cytokines to augment ADCC: a phase 1 trial of F16IL2 and BI 836858 for posttransplant AML relapse. Blood Advances, 2022, 6, 3684-3696.	2.5	5
3	The impact of TP53 co-mutations and immunologic microenvironment on outcome of lung cancer with EGFR exon 20 insertions. European Journal of Cancer, 2022, 170, 106-118.	1.3	15
4	Phase I trial of the DLL3/CD3 bispecific T-cell engager BI 764532 in DLL3-positive small-cell lung cancer and neuroendocrine carcinomas. Future Oncology, 2022, 18, 2639-2649.	1.1	14
5	Phase 1 Expansion Cohort of Ramucirumab Plus Pembrolizumab in Advanced Treatment-Naive NSCLC. Journal of Thoracic Oncology, 2021, 16, 289-298.	0.5	35
6	Phase I First-in-Human Dose Escalation Study of the oral SF3B1 modulator H3B-8800 in myeloid neoplasms. Leukemia, 2021, 35, 3542-3550.	3.3	97
7	Proof of concept for a rapidly switchable universal CAR-T platform with UniCAR-T-CD123 in relapsed/refractory AML. Blood, 2021, 137, 3145-3148.	0.6	70
8	Efficacy of Immune Checkpoint Inhibitors Alone or in Combination With Chemotherapy in NSCLC Harboring ERBB2 Mutations. Journal of Thoracic Oncology, 2021, 16, 1952-1958.	0.5	32
9	Selpercatinib in RET fusion-positive non-small-cell lung cancer (SIREN): a retrospective analysis of patients treated through an access program. Therapeutic Advances in Medical Oncology, 2021, 13, 175883592110196.	1.4	27
10	503â€Clinical activity of ICT01, an anti-BTN3A-targeted, γ9Î′2-activating mAb, alone and in combination with pembrolizumab in patients with advanced/refractory solid tumors: EVICTION trial. , 2021, 9, A535-A535.		0
11	Gemtuzumab Ozogamicin Plus Midostaurin in Combination with Standard Intensive Induction Therapy in Newly Diagnosed AML: Results from a Phase-I Study. Blood, 2021, 138, 2324-2324.	0.6	4
12	A Phase 1b Study of Blinatumomab Including Subcutaneous Administration in Relapsed / Refractory (R/R) Indolent Non Hodgkin's Lymphoma (NHL). Blood, 2021, 138, 2436-2436.	0.6	5
13	Allogeneic Hematopoietic Cell Transplantation Outcomes of Patients with R/R AML or Higher-Risk MDS Treated with the TIM-3 Inhibitor MBG453 (Sabatolimab) and Hypomethylating Agents. Blood, 2021, 138, 3677-3677.	0.6	5
14	Targeted Therapy in BRAF p.K601E–Driven NSCLC: Case Report and Literature Review. JCO Precision Oncology, 2020, 4, 1163-1166.	1.5	8
15	Long-term in vivo imaging reveals tumor-specific dissemination and captures host tumor interaction in zebrafish xenografts. Scientific Reports, 2020, 10, 13254.	1.6	20
16	Phase I experience with rogaratinib in patients (pts) with urothelial carcinoma (UC) selected based on <i>FGFR</i> mRNA overexpression Journal of Clinical Oncology, 2020, 38, 527-527.	0.8	4
17	Comparative RNAi Screens in Isogenic Human Stem Cells Reveal SMARCA4 as a Differential Regulator. Stem Cell Reports, 2019, 12, 1084-1098.	2.3	10
18	Enhanced labile plasma iron in hematopoietic stem cell transplanted patients promotes Aspergillus outgrowth. Blood Advances, 2019, 3, 1695-1700.	2.5	19

MARTIN WERMKE

#	Article	IF	CITATIONS
19	Enhanced labile plasma iron and outcome in acute myeloid leukaemia and myelodysplastic syndrome after allogeneic haemopoietic cell transplantation (ALLIVE): a prospective, multicentre, observational trial. Lancet Haematology,the, 2018, 5, e201-e210.	2.2	44
20	STK3 is a therapeutic target for a subset of acute myeloid leukemias. Oncotarget, 2018, 9, 25458-25473.	0.8	10
21	Characterization of a switchable chimeric antigen receptor platform in a pre-clinical solid tumor model. Oncolmmunology, 2017, 6, e1342909.	2.1	22
22	Mammalianâ€ŧarget of rapamycin inhibition with temsirolimus in myelodysplastic syndromes (<scp>MDS</scp>) patients is associated with considerable toxicity: results of the temsirolimus pilot trial by the German <scp>MDS</scp> Study Group (Dâ€MDS). British Journal of Haematology, 2016, 175, 917-924.	1.2	8
23	Selective expansion of regulatory T cells during lenalidomide treatment of myelodysplastic syndrome with isolated deletion 5q. Annals of Hematology, 2016, 95, 1805-1810.	0.8	11
24	RNAi profiling of primary human AML cells identifies ROCK1 as a therapeutic target and nominates fasudil as an antileukemic drug. Blood, 2015, 125, 3760-3768.	0.6	53
25	Iron Overload in Allogeneic Hematopoietic Cell Transplantation Outcome: A Meta-Analysis. Biology of Blood and Marrow Transplantation, 2014, 20, 1248-1251.	2.0	64
26	Reconstitution of Interleukin-17–Producing T Helper Cells after Allogeneic Hematopoietic Cell Transplantation. Biology of Blood and Marrow Transplantation, 2013, 19, 357-365.	2.0	11
27	In Vivo Expansion of Co-Transplanted T Cells Impacts on Tumor Re-Initiating Activity of Human Acute Myeloid Leukemia in NSG Mice. PLoS ONE, 2013, 8, e60680.	1.1	22
28	Response and Long-Term Outcome After Treatment With Third-Party Mesenchymal Stromal Cells - Updated Results In 58 Patients With Steroid-Refractory Acute Graft-Versus Host Disease Blood, 2013, 122, 4612-4612.	0.6	1
29	Reconstitution of 6-Sulfo LacNAc Dendritic Cells After Allogeneic Stem-Cell Transplantation. Transplantation, 2012, 93, 1270-1275.	0.5	5
30	MRI-Based Liver Iron Content Predicts for Nonrelapse Mortality in MDS and AML Patients Undergoing Allogeneic Stem Cell Transplantation. Clinical Cancer Research, 2012, 18, 6460-6468.	3.2	66
31	Reconstitution of IL-17-Producing T Helper Cells After Allogeneic Hematopoietic Cell Transplantation. Blood, 2012, 120, 4167-4167.	0.6	Ο
32	Azacytidine Compromises NK-Cell Activity in AML and MDS Patients Undergoing MRD-Based Pre-Emptive Treatment After Allogeneic Stem Cell Transplantation. Blood, 2012, 120, 4122-4122.	0.6	0
33	Systemic Iron Overload in Patients Undergoing Allogeneic Stem Cell Transplantation – a Magnetic Resonance Imaging Based Study in 81 AML and MDS Patients. Blood, 2011, 118, 489-489.	0.6	1
34	¹⁸⁸ Re antiâ€CD66 radioimmunotherapy combined with reducedâ€intensity conditioning and <i>inâ€vivo</i> T cell depletion in elderly patients undergoing allogeneic haematopoietic cell transplantation. British Journal of Haematology, 2010, 148, 910-917.	1.2	21
35	Genetic Variations of Interleukin-23R (1143A>G) and BPI (A645G), but Not of NOD2, Are Associated with Acute Graft-versus-Host Disease after Allogeneic Transplantation. Biology of Blood and Marrow Transplantation, 2010, 16, 1718-1727.	2.0	22
36	Comparing the Value of Serum Ferritin, Transfusion History and Magnetic Resonance Imaging for the Prediction of Iron Overload In MDS and AML Patients Undergoing Allogeneic Stem Cell Transplantation Blood, 2010, 116, 3493-3493.	0.6	0

MARTIN WERMKE

#	Article	IF	CITATIONS
37	The HCT-Specific Comorbidity Index Fails to Predict Survival After Allogeneic HCT In High Risk AML Patients - A Single Center Experience. Blood, 2010, 116, 1326-1326.	0.6	0
38	BPI A645G SNP but Not NOD2 Genotype Predicts for Acute Lung Injury After Allogeneic Stem Cell Transplantation. Blood, 2010, 116, 2324-2324.	0.6	11
39	Appearance of Mature 6-Sulfo LacNAc+ Dendritic Cells In Early and Late Engraftment After Allogeneic Stem Cell Transplantation Blood, 2010, 116, 3720-3720.	0.6	Ο
40	Comparative analysis of MLL partial tandem duplication and FLT3 internal tandem duplication mutations in 956 adult patients with acute myeloid leukemia. Genes Chromosomes and Cancer, 2003, 37, 237-251.	1.5	133
41	Analysis of FLT3-activating mutations in 979 patients with acute myelogenous leukemia: association with FAB subtypes and identification of subgroups with poor prognosis. Blood, 2002, 99, 4326-4335.	0.6	1,550