

Frank Petrak

List of Publications by Year in descending order

Source: <https://exaly.com/author-pdf/471638/publications.pdf>

Version: 2024-02-01

43
papers

1,640
citations

304743

22
h-index

302126

39
g-index

70
all docs

70
docs citations

70
times ranked

1995
citing authors

#	ARTICLE	IF	CITATIONS
1	Depression and diabetes: treatment and health-care delivery. <i>Lancet Diabetes and Endocrinology</i> , 2015, 3, 472-485.	11.4	156
2	Psychosocial morbidity of Graves' orbitopathy. <i>Clinical Endocrinology</i> , 2005, 63, 395-402.	2.4	152
3	The Association Between Depressive Symptoms and Systemic Inflammation in People With Type 2 Diabetes: Findings From the South London Diabetes Study. <i>Diabetes Care</i> , 2014, 37, 2186-2192.	8.6	95
4	Depression and Anxiety: Their Predictive Function for Weight Loss in Obese Individuals. <i>Obesity Facts</i> , 2009, 2, 227-234.	3.4	86
5	Treatment of depression in diabetes: an update. <i>Current Opinion in Psychiatry</i> , 2009, 22, 211-217.	6.3	79
6	Development and Validation of a New Measure to Evaluate Psychological Resistance to Insulin Treatment. <i>Diabetes Care</i> , 2007, 30, 2199-2204.	8.6	74
7	Influence of depressive and eating disorders on short- and long-term course of weight after surgical and nonsurgical weight loss treatment. <i>Comprehensive Psychiatry</i> , 2011, 52, 301-311.	3.1	70
8	The symptom-specific efficacy of antidepressant medication vs. cognitive behavioral therapy in the treatment of depression: results from an individual patient data meta-analysis. <i>World Psychiatry</i> , 2019, 18, 183-191.	10.4	68
9	Cognitive Behavioral Therapy Versus Sertraline in Patients With Depression and Poorly Controlled Diabetes: The Diabetes and Depression (DAD) Study. <i>Diabetes Care</i> , 2015, 38, 767-775.	8.6	66
10	Psychosocial Factors in Subjects with Thyroid-Associated Ophthalmopathy. <i>Thyroid</i> , 2002, 12, 237-239.	4.5	61
11	Prevalence of psychiatric disorders in an onset cohort of adults with type 1 diabetes. <i>Diabetes/Metabolism Research and Reviews</i> , 2003, 19, 216-222.	4.0	49
12	Psychological insulin resistance in geriatric patients with diabetes mellitus. <i>Patient Education and Counseling</i> , 2014, 94, 417-422.	2.2	45
13	Psychosocial Factors and Diabetes Mellitus: Evidence-Based Treatment Guidelines. <i>Current Diabetes Reviews</i> , 2005, 1, 255-270.	1.3	44
14	Suicidal Ideation in Adult Patients with Atopic Dermatitis: A German Cross-sectional Study. <i>Acta Dermato-Venereologica</i> , 2017, 97, 1189-1195.	1.3	40
15	Determinants of health-related quality of life in patients with persistent somatoform pain disorder. <i>European Journal of Pain</i> , 2003, 7, 463-471.	2.8	37
16	Cortisol, Platelet Serotonin Content, and Platelet Activity in Patients With Major Depression and Type 2 Diabetes. <i>Psychosomatic Medicine</i> , 2015, 77, 145-155.	2.0	31
17	Trajectories of depression in adults with newly diagnosed type 1 diabetes: results from the German Multicenter Diabetes Cohort Study. <i>Diabetologia</i> , 2017, 60, 60-68.	6.3	29
18	Do mental disorders and eating patterns affect long-term weight loss maintenance?. <i>General Hospital Psychiatry</i> , 2010, 32, 132-140.	2.4	26

#	ARTICLE	IF	CITATIONS
19	Psychological Insulin Resistance in Type 2 Diabetes Patients Regarding Oral Antidiabetes Treatment, Subcutaneous Insulin Injections, or Inhaled Insulin. <i>Diabetes Technology and Therapeutics</i> , 2013, 15, 702-710.	4.4	26
20	Behavioral medicine interventions in diabetes. <i>Current Opinion in Psychiatry</i> , 2010, 23, 131-138.	6.3	25
21	Study protocol of the Diabetes and Depression Study (DAD): a multi-center randomized controlled trial to compare the efficacy of a diabetes-specific cognitive behavioral group therapy versus sertraline in patients with major depression and poorly controlled diabetes mellitus. <i>BMC Psychiatry</i> , 2013, 13, 206.	2.6	18
22	Newly diagnosed depression is associated with increased beta-thromboglobulin levels and increased expression of platelet activation markers and platelet derived CD40-CD40L. <i>Journal of Psychiatric Research</i> , 2013, 47, 865-871.	3.1	18
23	Health information, behavior change, and decision support for patients with type 2 diabetes: development of a tailored, preference-sensitive health communication application. <i>Patient Preference and Adherence</i> , 2013, 7, 1091.	1.8	18
24	Can trajectories of glycemic control be predicted by depression, anxiety, or diabetes-related distress in a prospective cohort of adults with newly diagnosed type 1 diabetes? Results of a five-year follow-up from the German multicenter diabetes cohort study (GMDC-Study). <i>Diabetes Research and Clinical Practice</i> , 2018, 141, 106-117.	2.8	17
25	Cognitive behavioural therapy in elderly type 2 diabetes patients with minor depression or mild major depression: study protocol of a randomized controlled trial (MIND-DIA). <i>BMC Geriatrics</i> , 2010, 10, 21.	2.7	15
26	Adaptation to life after surgical removal of the bladder – an application of graphical Markov models for analysing longitudinal data. <i>Statistics in Medicine</i> , 2004, 23, 649-666.	1.6	13
27	hs-CRP Predicts Improvement in Depression in Patients With Type 1 Diabetes and Major Depression Undergoing Depression Treatment: Results From the Diabetes and Depression (DAD) Study. <i>Diabetes Care</i> , 2016, 39, e171-e173.	8.6	13
28	New pathways of increased cardiovascular risk in depression: A pilot study on the association of high-sensitivity C-reactive protein with pro-atherosclerotic markers in patients with depression. <i>Journal of Affective Disorders</i> , 2013, 146, 420-425.	4.1	11
29	Cost-effectiveness analysis of cognitive behaviour therapy for treatment of minor or mild-major depression in elderly patients with type 2 diabetes: study protocol for the economic evaluation alongside the MIND-DIA randomized controlled trial (MIND-DIA CEA). <i>BMC Geriatrics</i> , 2009, 9, 25.	2.7	9
30	Psychotherapeutische Kombinationsbehandlung. <i>Psychotherapeut</i> , 1998, 43, 282-287.	0.1	8
31	Associations between type of treatment and illness-specific locus of control in type 1 diabetes patients. <i>Psychology and Health</i> , 1993, 8, 383-391.	2.2	6
32	Stationäre verhaltenstherapeutisch-psychodynamische Kombinationsbehandlung. <i>Psychotherapeut</i> , 1999, 44, 241-247.	0.1	5
33	Checkliste zur Bewertung der wissenschaftlichen Qualität kontrollierter psychotherapeutischer Interventionsstudien (CPI). <i>Psychotherapeut</i> , 1999, 44, 390-393.	0.1	4
34	Data on diabetes-specific distress are needed to improve the quality of diabetes care. <i>Lancet, The</i> , 2021, 397, 2149.	13.7	4
35	Stationäre verhaltenstherapeutisch-psychodynamische Kombinationsbehandlung. <i>Psychotherapeut</i> , 1999, 44, 108-115.	0.1	2
36	Central serotonergic activity correlates with salivary cortisol after waking in depressed patients. <i>Psychopharmacology</i> , 2011, 217, 605-607.	3.1	2

#	ARTICLE	IF	CITATIONS
37	Reply to "Comments on Petrak et al.'s "Psychological Insulin Resistance in Type 2 Diabetes Patients Regarding Oral Antidiabetes Treatment, Subcutaneous Insulin Injections, or Inhaled Insulin": Considerations for Language and Communication". Diabetes Technology and Therapeutics, 2013, 15, 1055-1056.	4.4	2
38	Ängste und Angststörungen bei Diabetes mellitus. , 2013, , 145-157.		1
39	Psychodiabetologie in der Rehabilitation. , 2020, , 327-342.		1
40	Diabetes mellitus und Depression. , 2017, , 1-6.		0
41	Anpassung des Therapiemanuals an verschiedene klinische Settings. , 2017, , 45-47.		0
42	Diabetes mellitus und Altersdepression. , 2017, , 209-218.		0
43	Typ-2-Diabetes mellitus und Depression bei Älteren Menschen. , 2017, , 3-8.		0