## Yuejin Wang

## List of Publications by Year in descending order

Source: https://exaly.com/author-pdf/4719475/publications.pdf

Version: 2024-02-01

79	2,834	29 h-index	48
papers	citations		g-index
80	80	80	2313 citing authors
all docs	docs citations	times ranked	

#	Article	IF	CITATIONS
1	Genome-Wide Identification and Expression Analysis of Thioredoxin (Trx) Genes in Seed Development of Vitis vinifera. Journal of Plant Growth Regulation, 2022, 41, 3030-3045.	2.8	5
2	Genome Sequence Resource for <i>Colletotrichum viniferum</i> , the Cause of Grapevine Ripe Rot in China. Molecular Plant-Microbe Interactions, 2022, 35, 90-93.	1.4	4
3	Cloning and activity analysis of the highly expressed gene VviABCG20 promoter in seed and its activity is negatively regulated by the transcription factor VviDof14. Plant Science, 2022, 315, 111152.	1.7	2
4	Grafting with rootstocks promotes phenolic compound accumulation in grape berry skin during development based on integrative multi-omics analysis. Horticulture Research, 2022, 9, .	2.9	17
5	Genome-wide identification and characterisation of phenylalanine ammonia-lyase gene family in grapevine. Journal of Horticultural Science and Biotechnology, 2021, 96, 456-468.	0.9	18
6	Overexpression of two <i>CDPKs</i> from wild Chinese grapevine enhances powdery mildew resistance in <i>Vitis vinifera</i> and Arabidopsis. New Phytologist, 2021, 230, 2029-2046.	3.5	23
7	Importin-αs are required for the nuclear localization and function of the Plasmopara viticola effector PvAVH53. Horticulture Research, 2021, 8, 46.	2.9	15
8	<i>VvHDZ28</i> positively regulate salicylic acid biosynthesis during seed abortion in Thompson Seedless. Plant Biotechnology Journal, 2021, 19, 1824-1838.	4.1	24
9	VqMYB154 promotes polygene expression and enhances resistance to pathogens in Chinese wild grapevine. Horticulture Research, 2021, 8, 151.	2.9	30
10	Glyoxalase lâ€4 functions downstream of NAC72 to modulate downy mildew resistance in grapevine. Plant Journal, 2021, 108, 394-410.	2.8	18
11	VqBGH40a isolated from Chinese wild Vitis quinquangularis degrades trans-piceid and enhances trans-resveratrol. Plant Science, 2021, 310, 110989.	1.7	4
12	The co-expression of genes involved in seed coat and endosperm development promotes seed abortion in grapevine. Planta, 2021, 254, 87.	1.6	6
13	Genome-wide identification, phylogenetic analysis, and expression profiling of glycine-rich RNA-binding protein (GRPs) genes in seeded and seedless grapes (Vitis vinifera). Physiology and Molecular Biology of Plants, 2021, 27, 2231-2243.	1.4	6
14	The process of embryo abortion of stenospermocarpic grape and it develops into plantlet in vitro using embryo rescue. Plant Cell, Tissue and Organ Culture, 2020, 143, 389-409.	1.2	18
15	CRISPR/Cas9-mediated mutagenesis of VvMLO3 results in enhanced resistance to powdery mildew in grapevine (Vitis vinifera). Horticulture Research, 2020, 7, 116.	2.9	113
16	Expression and functional analysis of VviABCG14 from Vitis vinifera suggest the role in cytokinin transport and the interaction with VviABCG7. Plant Physiology and Biochemistry, 2020, 153, 1-10.	2.8	9
17	The WRKY53 transcription factor enhances stilbene synthesis and disease resistance by interacting with MYB14 and MYB15 in Chinese wild grape. Journal of Experimental Botany, 2020, 71, 3211-3226.	2.4	42
18	Heterologous expression of Chinese wild grapevine VqERFs in Arabidopsis thaliana enhance resistance to Pseudomonas syringae pv. tomato DC3000 and to Botrytis cinerea. Plant Science, 2020, 293, 110421.	1.7	25

#	Article	IF	CITATIONS
19	The grape ubiquitin ligase VpRH2 is a negative regulator in response to ABA treatment. Planta, 2020, 251, 88.	1.6	3
20	Insight Into Function and Subcellular Localization of Plasmopara viticola Putative RxLR Effectors. Frontiers in Microbiology, 2020, 11, 692.	1.5	16
21	Transcription factor VqERF114 regulates stilbene synthesis in Chinese wild Vitis quinquangularis by interacting with VqMYB35. Plant Cell Reports, 2019, 38, 1347-1360.	2.8	24
22	<i>VpSTS29/STS2</i> enhances fungal tolerance in grapevine through a positive feedback loop. Plant, Cell and Environment, 2019, 42, 2979-2998.	2.8	25
23	VqbZIP1 isolated from Chinese wild Vitis quinquangularis is involved in the ABA signaling pathway and regulates stilbene synthesis. Plant Science, 2019, 287, 110202.	1.7	16
24	Expression of stilbene synthase VqSTS6 from wild Chinese Vitis quinquangularis in grapevine enhances resveratrol production and powdery mildew resistance. Planta, 2019, 250, 1997-2007.	1.6	14
25	Molecular cloning and functional characterization of a seed-specific $Vv\hat{l}^2VPE$ gene promoter from Vitis vinifera. Planta, 2019, 250, 657-665.	1.6	8
26	Genome-wide analysis of glyoxalase-like gene families in grape (Vitis vinifera L.) and their expression profiling in response to downy mildew infection. BMC Genomics, 2019, 20, 362.	1.2	19
27	Dynamic translocation of stilbene synthase VpSTS29 from a Chinese wild Vitis species upon UV irradiation. Phytochemistry, 2019, 159, 137-147.	1.4	3
28	Genome-wide Identification, Phylogenetic Analysis, and Expression Profiling of CONSTANS-like (COL) Genes in Vitis vinifera. Journal of Plant Growth Regulation, 2019, 38, 631-643.	2.8	12
29	Improved in vitro Vitis vinifera L. embryo development of F1 progeny of †Delight†× †Ruby seedless†putrescine and marker-assisted selection. In Vitro Cellular and Developmental Biology - Plant, 2018, 54, 291-301.	™ using 0.9	15
30	Grapevine Vp <scp>PR</scp> 10.1 functions in resistance to <i>Plasmopara viticola</i> through triggering a cell deathâ€ike defence response by interacting with Vp <scp>VDAC</scp> 3. Plant Biotechnology Journal, 2018, 16, 1488-1501.	4.1	43
31	Ectopic expression of VpSTS29, a stilbene synthase gene from Vitis pseudoreticulata, indicates STS presence in cytosolic oil bodies. Planta, 2018, 248, 89-103.	1.6	16
32	CRISPR/Cas9â€mediated efficient targeted mutagenesis in grape in the first generation. Plant Biotechnology Journal, 2018, 16, 844-855.	4.1	270
33	Ectopic expression of Arabidopsis broad-spectrum resistance gene RPW8.2 improves the resistance to powdery mildew in grapevine (Vitis vinifera). Plant Science, 2018, 267, 20-31.	1.7	28
34	Current Progress and Future Prospects for the Clustered Regularly Interspaced Short Palindromic Repeats (CRISPR) Genome Editing Technology in Fruit Tree Breeding. Critical Reviews in Plant Sciences, 2018, 37, 233-258.	2.7	10
35	VlbZIP30 of grapevine functions in dehydration tolerance via the abscisic acid core signaling pathway. Horticulture Research, 2018, 5, 49.	2.9	20
36	Vacuolar processing enzyme (VvβVPE) from Vitis vinifera, processes seed proteins during ovule development, and accelerates seed germination in VvβVPE heterologously over-expressed Arabidopsis. Plant Science, 2018, 274, 420-431.	1.7	18

#	Article	IF	CITATIONS
37	Abnormal Somatic Embryo Reduction and Recycling in Grapevine Regeneration. Journal of Plant Growth Regulation, 2017, 36, 912-918.	2.8	9
38	VpPUB24, a novel gene from Chinese grapevine, Vitis pseudoreticulata, targets VpICE1 to enhance cold tolerance. Journal of Experimental Botany, 2017, 68, 2933-2949.	2.4	30
39	RING-H2-type E3 gene VpRH2 from Vitis pseudoreticulata improves resistance to powdery mildew by interacting with VpGRP2A. Journal of Experimental Botany, 2017, 68, 1669-1687.	2.4	32
40	VpuR9, a novel RING-type ubiquitin ligase gene from Vitis pseudoreticulata, is involved in powdery mildew response in transgenic V. vinifera plants. Plant Cell, Tissue and Organ Culture, 2017, 131, 41-49.	1.2	6
41	Overexpression of VpEIFP1, a novel F-box/Kelch-repeat protein from wild Chinese Vitis pseudoreticulata, confers higher tolerance to powdery mildew by inducing thioredoxin z proteolysis. Plant Science, 2017, 263, 142-155.	1.7	49
42	Analysis of the grape (Vitis vinifera L.) thaumatin-like protein (TLP) gene family and demonstration that TLP29 contributes to disease resistance. Scientific Reports, 2017, 7, 4269.	1.6	75
43	Evolution and expression analysis reveal the potential role of the HD-Zip gene family in regulation of embryo abortion in grapes (Vitis vinifera L.). BMC Genomics, 2017, 18, 744.	1.2	51
44	Insights into the Mechanisms Underlying Ultraviolet-C Induced Resveratrol Metabolism in Grapevine (V. amurensis Rupr.) cv. "Tonghua-3â€, Frontiers in Plant Science, 2016, 7, 503.	1.7	38
45	Identification and Characterization of Erysiphe necator-Responsive MicroRNAs in Chinese Wild Vitis pseudoreticulata by High-Throughput Sequencing. Frontiers in Plant Science, 2016, 7, 621.	1.7	50
46	The Novel Gene VpPR4-1 from Vitis pseudoreticulata Increases Powdery Mildew Resistance in Transgenic Vitis vinifera L Frontiers in Plant Science, 2016, 7, 695.	1.7	52
47	Expression of the Grape VqSTS21 Gene in Arabidopsis Confers Resistance to Osmotic Stress and Biotrophic Pathogens but Not Botrytis cinerea. Frontiers in Plant Science, 2016, 7, 1379.	1.7	23
48	Gene Cloning, Expression and Enzyme Activity of Vitis vinifera Vacuolar Processing Enzymes (VvVPEs). PLoS ONE, 2016, 11, e0160945.	1.1	13
49	Transcriptome analyses of seed development in grape hybrids reveals a possible mechanism influencing seed size. BMC Genomics, 2016, 17, 898.	1.2	39
50	Ectopic expression of a grape aspartic protease gene, AP13, in Arabidopsis thaliana improves resistance to powdery mildew but increases susceptibility to Botrytis cinerea. Plant Science, 2016, 248, 17-27.	1.7	47
51	Functional analysis of the fruit-specific promoter of VqSTS6 from the Chinese wild grape, Vitis quinquangularis. Agri Gene, 2016, 1, 38-45.	1.9	6
52	A stilbene synthase allele from a Chinese wild grapevine confers resistance to powdery mildew by recruiting salicylic acid signalling for efficient defence. Journal of Experimental Botany, 2016, 67, 5841-5856.	2.4	45
53	VqDUF642, a gene isolated from the Chinese grape Vitis quinquangularis, is involved in berry development and pathogen resistance. Planta, 2016, 244, 1075-1094.	1.6	22
54	Expression of a grape (Vitis vinifera) bZIP transcription factor, VlbZIP36, in Arabidopsis thaliana confers tolerance of drought stress during seed germination and seedling establishment. Plant Science, 2016, 252, 311-323.	1.7	31

#	Article	IF	CITATIONS
55	Genetic transformation of grape varieties and rootstocks via organogenesis. Plant Cell, Tissue and Organ Culture, 2016, 126, 541-552.	1.2	26
56	Genetic transformation of a fruit-specific, highly expressed stilbene synthase gene from Chinese wild Vitis quinquangularis. Planta, 2016, 243, 1041-1053.	1.6	45
57	Resistance evaluation of Chinese wild Vitis genotypes against Botrytis cinerea and different responses of resistant and susceptible hosts to the infection. Frontiers in Plant Science, 2015, 6, 854.	1.7	58
58	Histological responses to downy mildew in resistant and susceptible grapevines. Protoplasma, 2015, 252, 259-270.	1.0	44
59	Establishment of a picloram-induced somatic embryogenesis system in Vitis vinifera cv. chardonnay and genetic transformation of a stilbene synthase gene from wild-growing Vitis species. Plant Cell, Tissue and Organ Culture, 2015, 121, 397-412.	1.2	43
60	Constitutive expression of a grape aspartic protease gene in transgenic Arabidopsis confers osmotic stress tolerance. Plant Cell, Tissue and Organ Culture, 2015, 121, 275-287.	1.2	43
61	Gibberellin-induced changes in the transcriptome of grapevine (Vitis labrusca $\tilde{A}-V$ . vinifera) cv. Kyoho flowers. BMC Genomics, 2015, 16, 128.	1.2	77
62	Breeding new seedless grapes using in ovulo embryo rescue and marker-assisted selection. In Vitro Cellular and Developmental Biology - Plant, 2015, 51, 241-248.	0.9	22
63	In vitro induction of tetraploids in VitisÂ×ÂMuscadinia hybrids. Plant Cell, Tissue and Organ Culture, 2015, 122, 675-683.	1.2	28
64	Embryo rescue technique and its applications for seedless breeding in grape. Plant Cell, Tissue and Organ Culture, 2015, 120, 861-880.	1.2	29
65	Molecular Characteristics and Biochemical Functions of VpPR10s from Vitis pseudoreticulata Associated with Biotic and Abiotic Stresses. International Journal of Molecular Sciences, 2014, 15, 19162-19182.	1.8	24
66	Transcriptome sequencing and metabolite analysis reveals the role of delphinidin metabolism in flower colour in grape hyacinth. Journal of Experimental Botany, 2014, 65, 3157-3164.	2.4	185
67	Construction of Flower-specific Chimeric Promoters and Analysis of Their Activities in Transgenic Torenia. Plant Molecular Biology Reporter, 2014, 32, 234-245.	1.0	21
68	The comparative analysis of the potential relationship between resveratrol and stilbene synthase gene family in the development stages of grapes (Vitis quinquangularis and Vitis vinifera). Plant Physiology and Biochemistry, 2014, 74, 24-32.	2.8	70
69	A circulatory system useful both for long-term somatic embryogenesis and genetic transformation in Vitis vinifera L. cv. Thompson Seedless. Plant Cell, Tissue and Organ Culture, 2014, 118, 157-168.	1.2	45
70	Agrobacterium-mediated transformation of embryogenic cell suspension cultures and plant regeneration in Lilium tenuifolium orientalÂ×Âtrumpet â€~Robina'. Acta Physiologiae Plantarum, 2014, 36, 2047-2057.	1.0	18
71	The metacaspase gene family of Vitis vinifera L.: Characterization and differential expression during ovule abortion in stenospermocarpic seedless grapes. Gene, 2013, 528, 267-276.	1.0	46
72	Flower-specific expression of the Phalaenopsis flavonoid 3′, 5′-hydoxylase modifies flower color pigmentation in Petunia and Lilium. Plant Cell, Tissue and Organ Culture, 2013, 115, 263-273.	1.2	40

## YUEJIN WANG

#	Article	IF	CITATIONS
73	The Chinese wild grapevine ( <i>Vitis pseudoreticulata</i> ) E3 ubiquitin ligase <i>Erysiphe necator</i> â€induced RING finger protein 1 (EIRP1) activates plant defense responses by inducing proteolysis of the VpWRKY11 transcription factor. New Phytologist, 2013, 200, 834-846.	3.5	117
74	A Novel Heat Shock Transcription Factor, VpHsf1, from Chinese Wild Vitis pseudoreticulata is Involved in Biotic and Abiotic Stresses. Plant Molecular Biology Reporter, 2013, 31, 240-247.	1.0	40
75	Continuous biosynthesis of abscisic acid (ABA) may be required for maintaining dormancy of isolated embryos and intact seeds of Euonymus alatus. Plant Cell, Tissue and Organ Culture, 2012, 108, 493-500.	1.2	3
76	Expression pattern, genomic structure, and promoter analysis of the gene encoding stilbene synthase from Chinese wild Vitis pseudoreticulata. Journal of Experimental Botany, 2011, 62, 2745-2761.	2.4	66
77	VpRFP1, a novel C4C4-type RING finger protein gene from Chinese wild Vitis pseudoreticulata, functions as a transcriptional activator in defence response of grapevine. Journal of Experimental Botany, 2011, 62, 5671-5682.	2.4	43
78	Isolation and expression analysis of a novel pathogenesis-related protein 10 gene from Chinese wild Vitis pseudoreticulata induced by Uncinula necator. Biologia (Poland), 2010, 65, 653-659.	0.8	18
79	Characterization of a novel stilbene synthase promoter involved in pathogen- and stress-inducible expression from Chinese wild Vitis pseudoreticulata. Planta, 2010, 231, 475-487.	1.6	125