Sadao Suzuki

List of Publications by Year in descending order

Source: https://exaly.com/author-pdf/473040/publications.pdf Version: 2024-02-01



#	Article	IF	CITATIONS
1	No association between HPV vaccine and reported post-vaccination symptoms in Japanese young women: Results of the Nagoya study. Papillomavirus Research (Amsterdam, Netherlands), 2018, 5, 96-103.	4.5	128
2	Profile of Participants and Genotype Distributions of 108 Polymorphisms in a Cross-Sectional Study of Associations of Genotypes With Lifestyle and Clinical Factors: A Project in the Japan Multi-Institutional Collaborative Cohort (J-MICC) Study. Journal of Epidemiology, 2011, 21, 223-235.	1.1	92
3	Effects of smoking and drinking habits on the incidence of periodontal disease and tooth loss among Japanese males: a 4-yr longitudinal study. Journal of Periodontal Research, 2006, 41, 560-566.	1.4	81
4	Genome-wide meta-analysis identifies multiple novel loci associated with serum uric acid levels in Japanese individuals. Communications Biology, 2019, 2, 115.	2.0	66
5	The conditional relative odds ratio provided less biased results for comparing diagnostic test accuracy in meta-analyses. Journal of Clinical Epidemiology, 2004, 57, 461-469.	2.4	51
6	Dietary Habits and Risk of Ovarian Cancer Death in a Large-Scale Cohort Study (JACC Study) in Japan. Nutrition and Cancer, 2007, 57, 138-145.	0.9	46
7	Obesity/Weight Gain and Breast Cancer Risk: Findings From the Japan Collaborative Cohort Study for the Evaluation of Cancer Risk. Journal of Epidemiology, 2013, 23, 139-145.	1.1	45
8	Study Profile of the Japan Multi-institutional Collaborative Cohort (J-MICC) Study. Journal of Epidemiology, 2021, 31, 660-668.	1.1	41
9	Effect of Physical Activity on Breast Cancer Risk: Findings of the Japan Collaborative Cohort Study. Cancer Epidemiology Biomarkers and Prevention, 2008, 17, 3396-3401.	1.1	38
10	Genomewide Association Study of Leisure-Time Exercise Behavior in Japanese Adults. Medicine and Science in Sports and Exercise, 2018, 50, 2433-2441.	0.2	36
11	A genome-wide association study in the Japanese population identifies the 12q24 locus for habitual coffee consumption: The J-MICC Study. Scientific Reports, 2018, 8, 1493.	1.6	32
12	PURE RED CELL APLASIA INDUCED BY FK506. Transplantation, 1996, 61, 831,832.	0.5	27
13	Effect of the PPARG2 Pro12Ala Polymorphism and Clinical Risk Factors for Diabetes Mellitus on HbA1c in the Japanese General Population. Journal of Epidemiology, 2012, 22, 523-531.	1.1	24
14	Associations of Nutrient Patterns with the Prevalence of Metabolic Syndrome: Results from the Baseline Data of the Japan Multi-Institutional Collaborative Cohort Study. Nutrients, 2019, 11, 990.	1.7	24
15	Association of Dietary Acid Load with the Prevalence of Metabolic Syndrome among Participants in Baseline Survey of the Japan Multi-Institutional Collaborative Cohort Study. Nutrients, 2020, 12, 1605.	1.7	23
16	Accuracy of selfâ€reported height, weight and waist circumference in a Japanese sample. Obesity Science and Practice, 2017, 3, 417-424.	1.0	20
17	Low Intake of Vegetables and Fruits and Risk of Colorectal Cancer: The Japan Collaborative Cohort Study. Journal of Epidemiology, 2014, 24, 353-360.	1.1	19
18	Genome-Wide Association Study of Renal Function Traits: Results from the Japan Multi-Institutional Collaborative Cohort Study. American Journal of Nephrology, 2018, 47, 304-316.	1.4	18

SADAO SUZUKI

#	Article	IF	CITATIONS
19	eNOS genotype modifies the effect of leisure-time physical activity on serum triglyceride levels in a Japanese population. Lipids in Health and Disease, 2012, 11, 150.	1.2	16
20	Genome-wide association meta-analysis and Mendelian randomization analysis confirm the influence of ALDH2 on sleep durationin the Japanese population. Sleep, 2019, 42, .	0.6	16
21	Diagnostic accuracy of glycohemoglobin A1c (HbA1c) for postprandial hyperglycemia was equivalent to that of fasting blood glucose. Journal of Clinical Epidemiology, 2005, 58, 1052-1057.	2.4	15
22	Gene-Gene Combination Effect and Interactions among ABCA1, APOA1, SR-B1, and CETP Polymorphisms for Serum High-Density Lipoprotein-Cholesterol in the Japanese Population. PLoS ONE, 2013, 8, e82046.	1.1	15
23	Conditional Relative Odds Ratio and Comparison of Accuracy of Diagnostic Tests Based on 2×2 Tables. Journal of Epidemiology, 2006, 16, 145-153.	1.1	14
24	Community social capital and the onset of functional disability among older adults in Japan: a multilevel longitudinal study using Japan Gerontological Evaluation Study (JAGES) data. BMJ Open, 2019, 9, e029279.	0.8	14
25	GWAS analysis reveals a significant contribution of PSCA to the risk of Heliobacter pylori-induced gastric atrophy. Carcinogenesis, 2019, 40, 661-668.	1.3	13
26	A genomeâ€wide association study of coping behaviors suggests <i>FBXO45</i> is associated with emotional expression. Genes, Brain and Behavior, 2019, 18, e12481.	1.1	13
27	Re. Epidemiology, 2016, 27, e19.	1.2	12
28	Association of genetic risk score and chronic kidney disease in a Japanese population. Nephrology, 2019, 24, 670-673.	0.7	12
29	Genetic Variants of <i>RAMP2</i> and <i>CLR</i> are Associated with Stroke. Journal of Atherosclerosis and Thrombosis, 2017, 24, 1267-1281.	0.9	11
30	The interaction between ABCA1 polymorphism and physical activity on the HDL-cholesterol levels in a Japanese population. Journal of Lipid Research, 2020, 61, 86-94.	2.0	11
31	Impact of <i>PSCA</i> Polymorphisms on the Risk of Duodenal Ulcer. Journal of Epidemiology, 2021, 31, 12-20.	1.1	9
32	Diabetes Mellitus and Risk of Colorectal Cancer Mortality in Japan: the Japan Collaborative Cohort Study. Asian Pacific Journal of Cancer Prevention, 2016, 17, 4681-4688.	0.5	9
33	Insulin-like growth factor (IGF)-I, IGF-II, IGF binding protein-3, and risk of colorectal cancer: a nested case-control study in the Japan Collaborative Cohort study. Asian Pacific Journal of Cancer Prevention, 2009, 10 Suppl, 45-9.	0.5	9
34	Independent relationships of daily life activity and leisure-time exercise with metabolic syndrome and its traits in the general Japanese population. Endocrine, 2019, 64, 552-563.	1.1	8
35	Polymorphisms of genes involved in lipid metabolism and risk of chronic kidney disease in Japanese - cross-sectional data from the J-MICC study. Lipids in Health and Disease, 2014, 13, 162.	1.2	7
36	Association Between Second-hand Smoke Exposure and Depressive Symptoms Among Japanese Adults: A Cross-sectional Study. Journal of Epidemiology, 2020, 30, 566-573.	1.1	7

SADAO SUZUKI

#	Article	IF	CITATIONS
37	Moderate-to-vigorous Physical Activity and Sedentary Behavior Are Independently Associated With Renal Function: A Cross-sectional Study. Journal of Epidemiology, 2023, 33, 285-293.	1.1	7
38	Change in Food Intake Frequency at Five Years after Baseline in the JACC Study. Journal of Epidemiology, 2005, 15, S48-S55.	1.1	6
39	Cohort profile: Aichi regional sub-cohort of the Japan Environment and Children's Study (JECS-A). BMJ Open, 2019, 9, e028105.	0.8	6
40	Association of skipping breakfast and short sleep duration with the prevalence of metabolic syndrome in the general Japanese population: Baseline data from the Japan Multi-Institutional Collaborative cohort study. Preventive Medicine Reports, 2021, 24, 101613.	0.8	6
41	Associations between Dietary Patterns, ADRβ2 Gln27Glu and ADRβ3 Trp64Arg with Regard to Serum Triglyceride Levels: J-MICC Study. Nutrients, 2016, 8, 545.	1.7	5
42	Differential Effect of Polymorphisms on Body Mass Index Across the Life Course of Japanese: The Japan Multi-Institutional Collaborative Cohort Study. Journal of Epidemiology, 2021, 31, 172-179.	1.1	5
43	Development of de novo malignancies following renal transplantation: a single-center study. Transplantation Proceedings, 1994, 26, 938-40.	0.3	5
44	Associations Between hOGG1 Ser326Cys Polymorphism and Increased Body Mass Index and Fasting Glucose Level in the Japanese General Population. Journal of Epidemiology, 2014, 24, 379-384.	1.1	4
45	Letter to the editor: Safety concerns with human papilloma virus immunization in Japan: Analysis and evaluation of Nagoya City's surveillance data for adverse events. Japan Journal of Nursing Science, 2019, 16, 500-502.	0.5	4
46	Inhibitory effects of cyclosporin on exocrine and endocrine functions of the rat pancreas. Transplantation Proceedings, 1989, 21, 913-4.	0.3	4
47	Neighbourhood relationships moderate the positive association between family caregiver burden and psychological distress in Japanese adults: a cross-sectional study. Public Health, 2020, 185, 80-86.	1.4	3
48	Genetic polymorphisms in <i>CYP19A1</i> and <i>ESR1</i> are associated with serum CK activity after prolonged running in men. Journal of Applied Physiology, 2022, 132, 966-973.	1.2	3
49	Sex-specific Relationship Between Stress Coping Strategies and All-cause Mortality: Japan Multi-Institutional Collaborative Cohort Study. Journal of Epidemiology, 2023, 33, 236-245.	1.1	2
50	The association of reproductive history with hypertension and obesity according to menopausal status: the J-MICC Study. Hypertension Research, 2022, 45, 708-714.	1.5	2
51	Association of perceived stress and coping strategies with the renal function in middle-aged and older Japanese men and women. Scientific Reports, 2022, 12, 291.	1.6	1
52	Letter to the Editor: Authors' response to letter to the Editor: Safety concerns with human papilloma virus immunization in Japan: Analysis and evaluation of Nagoya City's surveillance data for adverse events. Japan Journal of Nursing Science, 2020, 17, e12309.	0.5	0
53	Population-Based Impact of Smoking, Drinking, and Genetic Factors on HDL-Cholesterol Levels in J-MICC Study Participants. Journal of Epidemiology, 2021, , .	1.1	0
54	Assessing the Relationship Between High-sensitivity C-reactive Protein and Kidney Function Employing Mendelian Randomization in the Japanese Community-based J-MICC Study. Journal of Epidemiology, 2021, , .	1.1	0

#	Article	IF	CITATIONS
55	Does Direct-to-Consumer Personal Genetic Testing Improve Gynecological Cancer Screening Uptake among Never-Screened Attendees? A Randomized Controlled Study. International Journal of Environmental Research and Public Health, 2021, 18, 12333.	1.2	0
56	Efficacy of Low-Dose Aspirin in Colorectal Cancer Risk Prevention is Dependent on <i>ADH1B</i> and <i>ALDH2</i> Genotype in Japanese Familial Adenomatous Polyposis Patients. Cancer Research Communications, 2022, 2, 483-488.	0.7	0