

Richard A James

List of Publications by Year in descending order

Source: <https://exaly.com/author-pdf/4732457/publications.pdf>

Version: 2024-02-01

44
papers

8,350
citations

109137

35
h-index

264894

42
g-index

44
all docs

44
docs citations

44
times ranked

6730
citing authors

#	ARTICLE	IF	CITATIONS
1	Impact of Varying Light and Dew on Ground Cover Estimates from Active NDVI, RGB, and LiDAR. <i>Plant Phenomics</i> , 2021, 2021, 9842178.	2.5	3
2	Transcriptional variation is associated with differences in shoot sodium accumulation in distinct barley varieties. <i>Environmental and Experimental Botany</i> , 2019, 166, 103812.	2.0	5
3	Evaluation of the Phenotypic Repeatability of Canopy Temperature in Wheat Using Continuous-Terrestrial and Airborne Measurements. <i>Frontiers in Plant Science</i> , 2019, 10, 875.	1.7	36
4	Deeper roots associated with cooler canopies, higher normalized difference vegetation index, and greater yield in three wheat populations grown on stored soil water. <i>Journal of Experimental Botany</i> , 2019, 70, 4963-4974.	2.4	43
5	High Throughput Determination of Plant Height, Ground Cover, and Above-Ground Biomass in Wheat with LiDAR. <i>Frontiers in Plant Science</i> , 2018, 9, 237.	1.7	206
6	Methodology for High-Throughput Field Phenotyping of Canopy Temperature Using Airborne Thermography. <i>Frontiers in Plant Science</i> , 2016, 7, 1808.	1.7	118
7	Tissue tolerance: an essential but elusive trait for salt-tolerant crops. <i>Functional Plant Biology</i> , 2016, 43, 1103.	1.1	162
8	High-throughput phenotyping technologies allow accurate selection of stay-green. <i>Journal of Experimental Botany</i> , 2016, 67, 4919-4924.	2.4	75
9	Rhizosheaths on wheat grown in acid soils: phosphorus acquisition efficiency and genetic control. <i>Journal of Experimental Botany</i> , 2016, 67, 3709-3718.	2.4	42
10	The barley anion channel, <i>HvALMT1</i> , has multiple roles in guard cell physiology and grain metabolism. <i>Physiologia Plantarum</i> , 2015, 153, 183-193.	2.6	40
11	Early vigour improves phosphate uptake in wheat. <i>Journal of Experimental Botany</i> , 2015, 66, 7089-7100.	2.4	46
12	Can citrate efflux from roots improve phosphorus uptake by plants? Testing the hypothesis with near-isogenic lines of wheat. <i>Physiologia Plantarum</i> , 2014, 151, 230-242.	2.6	71
13	Impact of ancestral wheat sodium exclusion genes <i>Nax1</i> and <i>Nax2</i> on grain yield of durum wheat on saline soils. <i>Functional Plant Biology</i> , 2012, 39, 609.	1.1	86
14	Infrared Thermography in Plant Phenotyping for Salinity Tolerance. , 2012, 913, 173-189.		23
15	Aluminium tolerance of root hairs underlies genotypic differences in rhizosheath size of wheat (<i>Triticum aestivum</i>) grown on acid soil. <i>New Phytologist</i> , 2012, 195, 609-619.	3.5	95
16	Wheat grain yield on saline soils is improved by an ancestral Na ⁺ transporter gene. <i>Nature Biotechnology</i> , 2012, 30, 360-364.	9.4	690
17	<i>Hordeum marinum</i> -wheat amphiploids maintain higher leaf K ⁺ :Na ⁺ and suffer less leaf injury than wheat parents in saline conditions. <i>Plant and Soil</i> , 2011, 348, 365-377.	1.8	28
18	Major genes for Na ⁺ exclusion, <i>Nax1</i> and <i>Nax2</i> (wheat <i>HKT1;4</i> and <i>HKT1;5</i>), decrease Na ⁺ accumulation in bread wheat leaves under saline and waterlogged conditions. <i>Journal of Experimental Botany</i> , 2011, 62, 2939-2947.	2.4	394

#	ARTICLE	IF	CITATIONS
19	Characterisation of HvALMT1 function in transgenic barley plants. <i>Functional Plant Biology</i> , 2011, 38, 163.	1.1	35
20	Stomatal conductance as a screen for osmotic stress tolerance in durum wheat growing in saline soil. <i>Functional Plant Biology</i> , 2010, 37, 255.	1.1	288
21	New phenotyping methods for screening wheat and barley for beneficial responses to water deficit. <i>Journal of Experimental Botany</i> , 2010, 61, 3499-3507.	2.4	359
22	A new screening method for osmotic component of salinity tolerance in cereals using infrared thermography. <i>Functional Plant Biology</i> , 2009, 36, 970.	1.1	173
23	Genetic variation in tolerance to the osmotic stress component of salinity stress in durum wheat. <i>Functional Plant Biology</i> , 2008, 35, 111.	1.1	126
24	Cell-specific localization of Na ⁺ in roots of durum wheat and possible control points for salt exclusion. <i>Plant, Cell and Environment</i> , 2008, 31, 1565-1574.	2.8	90
25	Osmotic adjustment leads to anomalously low estimates of relative water content in wheat and barley. <i>Functional Plant Biology</i> , 2008, 35, 1172.	1.1	100
26	HKT1;5-Like Cation Transporters Linked to Na ⁺ Exclusion Loci in Wheat, Nax2 and Kna1. <i>Plant Physiology</i> , 2007, 143, 1918-1928.	2.3	378
27	Approaches to increasing the salt tolerance of wheat and other cereals. <i>Journal of Experimental Botany</i> , 2006, 57, 1025-1043.	2.4	1,484
28	Physiological Characterization of Two Genes for Na ⁺ Exclusion in Durum Wheat, Nax1 and Nax2. <i>Plant Physiology</i> , 2006, 142, 1537-1547.	2.3	350
29	Photosynthetic capacity is related to the cellular and subcellular partitioning of Na ⁺ , K ⁺ and Cl ⁻ in salt-affected barley and durum wheat. <i>Plant, Cell and Environment</i> , 2006, 29, 2185-2197.	2.8	180
30	A Sodium Transporter (HKT7) Is a Candidate for Nax1, a Gene for Salt Tolerance in Durum Wheat. <i>Plant Physiology</i> , 2006, 142, 1718-1727.	2.3	266
31	Control of Sodium Transport in Durum Wheat. <i>Plant Physiology</i> , 2005, 137, 807-818.	2.3	264
32	Screening methods for salinity tolerance: a case study with tetraploid wheat. <i>Plant and Soil</i> , 2003, 253, 201-218.	1.8	609
33	Genetic control of sodium exclusion in durum wheat. <i>Australian Journal of Agricultural Research</i> , 2003, 54, 627.	1.5	115
34	Avenues for increasing salt tolerance of crops, and the role of physiologically based selection traits. <i>Plant and Soil</i> , 2002, 243, 93-105.		61
35	Stomatal control in tomato with ABA-deficient roots: response of grafted plants to soil drying. <i>Journal of Experimental Botany</i> , 2002, 53, 1503-1514.	2.4	191
36	Effect of salinity on water relations and growth of wheat genotypes with contrasting sodium uptake. <i>Functional Plant Biology</i> , 2002, 29, 1065.	1.1	93

#	ARTICLE	IF	CITATIONS
37	Factors affecting CO ₂ assimilation, leaf injury and growth in salt-stressed durum wheat. <i>Functional Plant Biology</i> , 2002, 29, 1393.	1.1	259
38	Title is missing!. <i>Plant and Soil</i> , 2002, 247, 93-105.	1.8	252
39	Stomatal control in tomato with ABA-deficient roots: response of grafted plants to soil drying. <i>Journal of Experimental Botany</i> , 2002, 53, 1503-14.	2.4	165
40	Genetic variation for improving the salt tolerance of durum wheat. <i>Australian Journal of Agricultural Research</i> , 2000, 51, 69.	1.5	218
41	Does water and phosphorus uptake limit leaf growth of <i>Rhizoctonia</i> -infected wheat seedlings?. <i>Plant and Soil</i> , 1999, 209, 157-166.	1.8	20
42	Contribution of <i>Rhizoctonia</i> to reduced seedling growth of direct-drilled wheat: studies with intact cores. <i>Australian Journal of Agricultural Research</i> , 1997, 48, 1231.	1.5	8
43	Reduced growth and yield of wheat with conservation cropping. II. Soil biological factors limit growth under direct drilling. <i>Australian Journal of Agricultural Research</i> , 1995, 46, 75.	1.5	56
44	Stored xylem sap from wheat and barley in drying soil contains a transpiration inhibitor with a large molecular size. <i>Plant, Cell and Environment</i> , 1993, 16, 867-872.	2.8	47