

Andrea Icks

List of Publications by Year in descending order

Source: <https://exaly.com/author-pdf/4733302/publications.pdf>

Version: 2024-02-01

77
papers

1,849
citations

279487

23
h-index

288905

40
g-index

87
all docs

87
docs citations

87
times ranked

2254
citing authors

#	ARTICLE	IF	CITATIONS
1	Socio-economic inequalities in glycaemic control in recently diagnosed adults with type 1 and type 2 diabetes. <i>Diabetic Medicine</i> , 2022, 39, e14833.	1.2	3
2	Study protocol for a randomised controlled trial of a web-based behavioural lifestyle programme for emPOWERment in early Multiple Sclerosis (POWER@MS1). <i>BMJ Open</i> , 2021, 11, e041720.	0.8	5
3	Mental Health Specialist Video Consultations Versus Treatment-as-Usual for Patients With Depression or Anxiety Disorders in Primary Care: Randomized Controlled Feasibility Trial. <i>JMIR Mental Health</i> , 2021, 8, e22569.	1.7	16
4	Health care utilization and excess costs after pelvic fractures among older people in Germany. <i>Osteoporosis International</i> , 2021, 32, 2061-2072.	1.3	5
5	Mental health specialist video consultations versus treatment as usual in patients with depression or anxiety disorders in primary care: study protocol for an individually randomised superiority trial (the) Tj ETQq1 1 0.784314 rgBT /Over	1.3	5
6	Effects of a Brief Web-Based "Social Norms" Intervention on Alcohol, Tobacco and Cannabis Use Among German University Students: Results of a Cluster-Controlled Trial Conducted at Eight Universities. <i>Frontiers in Public Health</i> , 2021, 9, 659875.	1.3	5
7	High Depressive Symptoms in Previously Undetected Diabetes " 10-Year Follow-Up Results of the Heinz Nixdorf Recall Study. <i>Clinical Epidemiology</i> , 2021, Volume 13, 429-438.	1.5	4
8	BÄrgerwissenschaftliche Forschungsansätze in Medizin und Gesundheitsforschung. <i>TATuP - Zeitschrift für Technikfolgenabschätzung in Theorie Und Praxis</i> , 2021, 30, 63-69.	0.2	2
9	Patient time costs due to self-management in diabetes may be as high as direct medical costs: results from the population-based KORA survey FF4 in Germany. <i>Diabetic Medicine</i> , 2020, 37, 895-897.	1.2	6
10	Different information needs in subgroups of people with diabetes mellitus: a latent class analysis. <i>BMC Public Health</i> , 2020, 20, 1901.	1.2	4
11	Standardized risk management in catheterization procedures for non-ST-segment elevation myocardial infarction: associations with in-hospital clinical outcomes. <i>European Heart Journal</i> , 2020, 41, .	1.0	0
12	Which patients with type 2 diabetes mellitus are perceived as "difficult" by general practitioners? <i>Primary Care Diabetes</i> , 2019, 13, 353-359.	0.9	7
13	Effectiveness of a complex intervention to improve participation and activities in nursing home residents with joint contractures (JointConEval): study protocol of a multicentre cluster-randomised controlled trial [DRKS-ID:DRKS00015185]. <i>Trials</i> , 2019, 20, 305.	0.7	2
14	Effect of person-centred care on antipsychotic drug use in nursing homes (EPCentCare): a cluster-randomised controlled trial. <i>Age and Ageing</i> , 2019, 48, 419-425.	0.7	26
15	Implementation of a multicomponent intervention to prevent physical restraints in nursing homes (IMPRINT): A pragmatic cluster randomized controlled trial. <i>International Journal of Nursing Studies</i> , 2019, 96, 27-34.	2.5	52
16	P41...Effectiveness of ACP to improve patient-centred care: study protocol of a cluster-randomised intervention trial focussing on nursing home residents (BEVOR-Study)., 2019, , .		1
17	Healthcare provision, functional ability and quality of life after proximal femoral fracture - "ProFem": Study protocol of a population-based, prospective study based on individually linked survey and statutory health insurance data. <i>BMJ Open</i> , 2019, 9, e028144.	0.8	1
18	Mental health specialist video consultations for patients with depression or anxiety disorders in primary care: protocol for a randomised controlled feasibility trial. <i>BMJ Open</i> , 2019, 9, e030003.	0.8	24

#	ARTICLE	IF	CITATIONS
19	Time spent on self-management by people with diabetes: results from the population-based KORA survey in Germany. <i>Diabetic Medicine</i> , 2019, 36, 970-981.	1.2	8
20	Incidence and relative risk of renal replacement therapy (RRT) in people with and without diabetes between 2002 and 2016 in a German region. <i>Diabetologie Und Stoffwechsel</i> , 2019, 14, .	0.0	0
21	Secondary data in diabetes surveillance - co-operation projects and definition of references on the documented prevalence of diabetes.. , 2019, 4, 50-63.		1
22	Information needs in people with diabetes mellitus: a systematic review. <i>Systematic Reviews</i> , 2018, 7, 27.	2.5	30
23	What information needs do people with recently diagnosed diabetes mellitus have and what are the associated factors? A cross-sectional study in Germany. <i>BMJ Open</i> , 2018, 8, e017895.	0.8	12
24	How do patients with diabetes report their comorbidity? Comparison with administrative data. <i>Clinical Epidemiology</i> , 2018, Volume 10, 499-509.	1.5	10
25	General practitioners'™ attitudes towards patients with poorly controlled type 2 diabetes: a qualitative study. <i>BMC Family Practice</i> , 2018, 19, 49.	2.9	12
26	Healthcare costs of Type 2 diabetes in Germany. <i>Diabetic Medicine</i> , 2017, 34, 855-861.	1.2	61
27	Time cost of diabetes: Development of a questionnaire to assess time spent on diabetes self-care. <i>Journal of Diabetes and Its Complications</i> , 2017, 31, 260-266.	1.2	8
28	Cost burden of type 2 diabetes in Germany: results from the population-based KORA studies. <i>BMJ Open</i> , 2016, 6, e012527.	0.8	50
29	Determinants of patients'™ needs in asthma treatment: a cross-sectional study. <i>Npj Primary Care Respiratory Medicine</i> , 2016, 26, 16044.	1.1	7
30	Regional Disparities in Diabetes Care for Pediatric Patients with Type 1 Diabetes. A Cross-sectional DPV Multicenter Analysis of 24%928 German Children and Adolescents. <i>Experimental and Clinical Endocrinology and Diabetes</i> , 2016, 124, 111-119.	0.6	14
31	Drug-based pain management in people with dementia after hip or pelvic fractures: a systematic review protocol. <i>Systematic Reviews</i> , 2016, 5, 113.	2.5	2
32	Direct costs in impaired glucose regulation: results from the population-based Heinz Nixdorf Recall study. <i>BMJ Open Diabetes Research and Care</i> , 2016, 4, e000172.	1.2	6
33	Cohort profile: the German Diabetes Study (GDS). <i>Cardiovascular Diabetology</i> , 2016, 15, 59.	2.7	97
34	Assessment of information needs in diabetes: Development and evaluation of a questionnaire. <i>Primary Care Diabetes</i> , 2016, 10, 287-292.	0.9	13
35	Studies analysing the need for health-related information in Germany - a systematic review. <i>BMC Health Services Research</i> , 2015, 15, 407.	0.9	27
36	Effect of person-centred care on antipsychotic drug use in nursing homes (EPCentCare): study protocol for a cluster-randomised controlled trial. <i>Implementation Science</i> , 2015, 10, 82.	2.5	23

#	ARTICLE	IF	CITATIONS
37	Epidemiology of Pelvic Fractures in Germany: Considerably High Incidence Rates among Older People. PLoS ONE, 2015, 10, e0139078.	1.1	96
38	Implementation of a multicomponent intervention to prevent physical restraints in nursing home residents (IMPRINT): study protocol for a cluster-randomised controlled trial. BMC Geriatrics, 2015, 15, 86.	1.1	18
39	Incidence of hip fractures in Germany, 1995–2010. Archives of Osteoporosis, 2013, 8, 140.	1.0	56
40	Diabetes incidence does not differ between subjects with and without high depressive symptoms – 5-year follow-up results of the Heinz Nixdorf Recall Study. Diabetic Medicine, 2013, 30, 65-69.	1.2	8
41	Medication Costs by Glucose Tolerance Stage in Younger and Older Women and Men: Results from the Population-based KORA Survey in Germany. Experimental and Clinical Endocrinology and Diabetes, 2013, 121, 614-623.	0.6	3
42	Identification of a Multistate Continuous-Time Nonhomogeneous Markov Chain Model for Patients with Decreased Renal Function. Medical Decision Making, 2013, 33, 298-306.	1.2	16
43	Drug Costs in Prediabetes and Undetected Diabetes Compared With Diagnosed Diabetes and Normal Glucose Tolerance: Results From the Population-Based KORA Survey in Germany. Diabetes Care, 2013, 36, e53-e54.	4.3	13
44	Patient time costs attributable to healthcare use in diabetes: results from the population-based KORA survey in Germany. Diabetic Medicine, 2013, 30, 1245-1249.	1.2	5
45	Frequency and Cost of Diabetic Ketoacidosis in Germany – Study in 12,001 Paediatric Patients. Experimental and Clinical Endocrinology and Diabetes, 2013, 121, 58-59.	0.6	24
46	Risk for High Depressive Symptoms in Diagnosed and Previously Undetected Diabetes: 5-Year Follow-Up Results of the Heinz Nixdorf Recall Study. PLoS ONE, 2013, 8, e56300.	1.1	10
47	Time-Dependent Impact of Diabetes on Mortality in Patients With Stroke. Diabetes Care, 2012, 35, 1868-1875.	4.3	37
48	Pre-test of questions on health-related resource use and expenditure, using behaviour coding and cognitive interviewing techniques. BMC Health Services Research, 2012, 12, 303.	0.9	23
49	Correction factor for the analysis of the hip fracture incidence – differences between age, sex, region, and calendar year. Wiener Klinische Wochenschrift, 2012, 124, 391-394.	1.0	10
50	Blood glucose testing and primary prevention of Type 2 diabetes – evaluation of the effect of evidence-based patient information: a randomized controlled trial. Diabetic Medicine, 2012, 29, 1011-1020.	1.2	6
51	Time-dependent impact of diabetes on the mortality of patients on renal replacement therapy: A population-based study in Germany (2002–2009). Diabetes Research and Clinical Practice, 2011, 92, 380-385.	1.1	6
52	P2-269 Regional differences in the prevalence of type 2 diabetes mellitus: results from five population-based cohort studies in Germany (DIAB-CORE consortium). Journal of Epidemiology and Community Health, 2011, 65, A296-A296.	2.0	0
53	Time-Dependent Impact of Diabetes on Mortality in Patients After Major Lower Extremity Amputation. Diabetes Care, 2011, 34, 1350-1354.	4.3	88
54	Is it really a true reversal?. Wiener Klinische Wochenschrift, 2010, 122, 545-545.	1.0	1

#	ARTICLE	IF	CITATIONS
55	Within-trial economic evaluation of diabetes-specific cognitive behaviour therapy in patients with type 2 diabetes and subthreshold depression. <i>BMC Public Health</i> , 2010, 10, 625.	1.2	6
56	Reduced incidence of blindness in relation to diabetes mellitus in southern Germany?. <i>Diabetic Medicine</i> , 2010, 27, 1138-1143.	1.2	25
57	Costs of dialysis—a regional population-based analysis. <i>Nephrology Dialysis Transplantation</i> , 2010, 25, 1647-1652.	0.4	66
58	Incidence of Lower-limb Amputations in the Diabetic Compared to the Non-diabetic Population. Findings from Nationwide Insurance Data, Germany, 2005–2007. <i>Experimental and Clinical Endocrinology and Diabetes</i> , 2009, 117, 500-504.	0.6	99
59	Hip fractures and area level socioeconomic conditions: a population-based study. <i>BMC Public Health</i> , 2009, 9, 114.	1.2	31
60	Differences in trends in estimated incidence of myocardial infarction in non-diabetic and diabetic people: Monitoring Trends and Determinants on Cardiovascular Diseases (MONICA)/Cooperative Health Research in the Region of Augsburg (KORA) registry. <i>Diabetologia</i> , 2009, 52, 1836-1841.	2.9	29
61	Cost-effectiveness analysis of cognitive behaviour therapy for treatment of minor or mild-major depression in elderly patients with type 2 diabetes: study protocol for the economic evaluation alongside the MIND-DIA randomized controlled trial (MIND-DIA CEA). <i>BMC Geriatrics</i> , 2009, 9, 25.	1.1	9
62	Trend of hip fracture incidence in Germany 1995–2004: a population-based study. <i>Osteoporosis International</i> , 2008, 19, 1139-1145.	1.3	182
63	Hip fracture incidence in the elderly in Austria: An epidemiological study covering the years 1994 to 2006. <i>BMC Geriatrics</i> , 2008, 8, 35.	1.1	54
64	Are symptoms of depression more common in diabetes? Results from the Heinz Nixdorf Recall study. <i>Diabetic Medicine</i> , 2008, 25, 1330-1336.	1.2	46
65	Economics in Pediatric Type 1 Diabetes - Results from Recently Published Studies. <i>Experimental and Clinical Endocrinology and Diabetes</i> , 2007, 115, 448-454.	0.6	20
66	REMARKABLE DECLINE IN HIP FRACTURE INCIDENCE IN YOUNGER AGE GROUPS IN GERMANY. <i>American Journal of Public Health</i> , 2007, 97, 1733-1734.	1.5	8
67	Persistent social disparities in the risk of hospital admission of paediatric diabetic patients in Germany-prospective data from 1277 diabetic children and adolescents. <i>Diabetic Medicine</i> , 2007, 24, 440-442.	1.2	28
68	Cost comparison analysis: pentaerythrithyl tetranitrate (PETN) and isosorbide dinitrate (ISDN) prescribed to diabetic patients in primary care practices in Germany. <i>International Journal of Clinical Pharmacology and Therapeutics</i> , 2007, 45, 516-523.	0.3	6
69	Incremental Prescription and Drug Costs during the Years Preceding Diabetes Diagnosis in Primary Care Practices in Germany. <i>Experimental and Clinical Endocrinology and Diabetes</i> , 2006, 114, 348-355.	0.6	13
70	Increased Use of Hip Protectors in Nursing Homes: Economic Analysis of a Cluster Randomized, Controlled Trial. <i>Journal of the American Geriatrics Society</i> , 2005, 53, 2153-2158.	1.3	24
71	Cost-Effectiveness of Type 2 Diabetes Screening: Results from Recently Published Studies. <i>Gesundheitswesen</i> , 2005, 67, 167-171.	0.8	13
72	Direct Costs of Pediatric Diabetes Care in Germany and Their Predictors. <i>Experimental and Clinical Endocrinology and Diabetes</i> , 2004, 112, 302-309.	0.6	33

#	ARTICLE	IF	CITATIONS
73	Hospitalization among diabetic children and adolescents and non-diabetic control subjects: a prospective population-based study. <i>Diabetologia</i> , 2001, 44, B87-B92.	2.9	17
74	Hospitalization Among Diabetic Children and Adolescents and the General Population in Germany. <i>Diabetes Care</i> , 2001, 24, 435-440.	4.3	29
75	Incidence of Blindness in Relation to Diabetes: A population-based study. <i>Diabetes Care</i> , 1997, 20, 1147-1153.	4.3	79
76	Blindness Due to Diabetes: Population-based Age- and Sex-specific Incidence Rates. , 1997, 14, 571-575.		46
77	Evaluation of the effects of the population-based intervention "10.000 Steps Duesseldorf" promoting physical activity in community-dwelling adults residing in two cities in North Rhine-Westphalia: Study protocol (Preprint). <i>JMIR Research Protocols</i> , 0, , .	0.5	0