
David P Sonne

List of Publications by Year in descending order

Source: https://exaly.com/author-pdf/4740441/publications.pdf Version: 2024-02-01



#	Article	IF	CITATIONS
1	Effect of single doses of citalopram and reboxetine on urethral pressure: A randomized, doubleâ€blind, placebo―and activeâ€controlled threeâ€period crossover study in healthy women. Neurourology and Urodynamics, 2022, 41, 1482-1488.	0.8	2
2	Protocol for a randomised, double-blinded, placebo-controlled, double-dummy 6-week clinical trial comparing the treatment effects of the glucagon-like peptide 1 receptor agonist liraglutide versus the bile acid sequestrant colesevelam on bile acid malabsorption. BMJ Open, 2021, 11, e044711.	0.8	3
3	MECHANISMS IN ENDOCRINOLOGY: FXR signalling: a novel target in metabolic diseases. European Journal of Endocrinology, 2021, 184, R193-R205.	1.9	14
4	What is on the horizon for type 2 diabetes pharmacotherapy? – An overview of the antidiabetic drug development pipeline. Expert Opinion on Drug Discovery, 2020, 15, 1253-1265.	2.5	6
5	Glucagon-Like Peptide 2 Inhibits Postprandial Gallbladder Emptying in Man: A Randomized, Double-Blinded, Crossover Study. Clinical and Translational Gastroenterology, 2020, 11, e00257.	1.3	8
6	Remission of Bile Acid Malabsorption Symptoms Following Treatment With the Glucagon-Like Peptide 1 Receptor Agonist Liraglutide. Gastroenterology, 2019, 157, 569-571.	0.6	16
7	Glucoseâ€lowering effects and mechanisms of the bile acidâ€sequestering resin sevelamer. Diabetes, Obesity and Metabolism, 2018, 20, 1623-1631.	2.2	21
8	Determinants of Fasting Hyperglucagonemia in Patients with Type 2 Diabetes and Nondiabetic Control Subjects. Metabolic Syndrome and Related Disorders, 2018, 16, 530-536.	0.5	22
9	Restoration of enteroendocrine and pancreatic function after internal hernia and short bowel syndrome in a young woman with gastric bypass - a 2-year follow-up. Physiological Reports, 2018, 6, e13686.	0.7	1
10	Model-Based Prediction of Plasma Concentration and Enterohepatic Circulation of Total Bile Acids in Humans. CPT: Pharmacometrics and Systems Pharmacology, 2018, 7, 603-612.	1.3	12
11	Effects of liraglutide on gallbladder emptying: A randomized, placeboâ€controlled trial in adults with overweight or obesity. Diabetes, Obesity and Metabolism, 2018, 20, 2557-2564.	2.2	28
12	Metformin-induced glucagon-like peptide-1 secretion contributes to the actions of metformin in type 2 diabetes. JCI Insight, 2018, 3, .	2.3	86
13	Bile acid sequestrants for glycemic control in patients with type 2 diabetes: A systematic review with meta-analysis of randomized controlled trials. Journal of Diabetes and Its Complications, 2017, 31, 918-927.	1.2	72
14	Dipeptidyl-peptidase (DPP)-4 inhibitors and glucagon-like peptide (GLP)-1 analogues for prevention or delay of type 2 diabetes mellitus and its associated complications in people at increased risk for the development of type 2 diabetes mellitus. The Cochrane Library, 2017, 5, CD012204.	1.5	31
15	Postprandial Plasma Concentrations of ProANP in Patients with Type 2 Diabetes and Healthy Controls. Clinical Chemistry, 2017, 63, 1040-1041.	1.5	5
16	Evidence connecting old, new and neglected glucoseâ€lowering drugs to bile acidâ€induced <scp>GLP</scp> â€l secretion: <scp>A</scp> review. Diabetes, Obesity and Metabolism, 2017, 19, 1214-1222.	2.2	14
17	Clinical relevance of the bile acid receptor TGR5 in metabolism. Lancet Diabetes and Endocrinology,the, 2017, 5, 224-233.	5.5	105
18	Involvement of glucagonâ€like peptideâ€1 in the glucoseâ€lowering effect of metformin. Diabetes, Obesity and Metabolism, 2016, 18, 955-961.	2.2	50

DAVID P SONNE

#	Article	IF	CITATIONS
19	Cholecystokinin-Induced Gallbladder Emptying and Metformin Elicit Additive Glucagon-Like Peptide-1 Responses. Journal of Clinical Endocrinology and Metabolism, 2016, 101, 2076-2083.	1.8	24
20	Postprandial Plasma Concentrations of Individual Bile Acids and FGF-19 in Patients With Type 2 Diabetes. Journal of Clinical Endocrinology and Metabolism, 2016, 101, 3002-3009.	1.8	100
21	Comment on Xu et al. Effects of Metformin on Metabolite Profiles and LDL Cholesterol in Patients With Type 2 Diabetes. Diabetes Care 2015;38:1858–1867. Diabetes Care, 2015, 38, e215-e215.	4.3	5
22	Pancreatic Amylase and Lipase Plasma Concentrations Are Unaffected by Increments in Endogenous GLP-1 Levels Following Liquid Meal Tests. Diabetes Care, 2015, 38, e71-e72.	4.3	11
23	MECHANISMS IN ENDOCRINOLOGY: Bile acid sequestrants in type 2 diabetes: potential effects on GLP1 secretion. European Journal of Endocrinology, 2014, 171, R47-R65.	1.9	62
24	On the role of gallbladder emptying and incretin hormones for nutrient-mediated TSH suppression in patients with type 2 diabetes. Endocrine Connections, 2014, 3, 193-199.	0.8	2
25	Sulfonylurea versus metformin monotherapy in patients with type 2 diabetes: a Cochrane systematic review and meta-analysis of randomized clinical trials and trial sequential analysis. CMAJ Open, 2014, 2, E162-E175.	1.1	73
26	Postprandial gallbladder emptying in patients with type 2 diabetes: potential implications for bile-induced secretion of glucagon-like peptide 1. European Journal of Endocrinology, 2014, 171, 407-419.	1.9	56
27	Postprandial gut hormone responses and glucose metabolism in cholecystectomized patients. American Journal of Physiology - Renal Physiology, 2013, 304, G413-G419.	1.6	43
28	Cardiovascular effects of alpha-linolenic acid – a possible role of glucagon-like peptide-1. Experimental Biology and Medicine, 2013, 238, 1116-1117.	1.1	0