Brendan Carvalho

List of Publications by Year in Descending Order

Source: https://exaly.com/author-pdf/4750700/brendan-carvalho-publications-by-year.pdf

Version: 2024-04-27

This document has been generated based on the publications and citations recorded by exaly.com. For the latest version of this publication list, visit the link given above.

The third column is the impact factor (IF) of the journal, and the fourth column is the number of citations of the article.

212 5,135 39 61 g-index

232 6,446 3.6 ext. papers ext. citations avg, IF L-index

| # | Paper | IF | Citations |
|-----|---|------|-----------|
| 212 | Limitations of Gravimetric Quantitative Blood Loss during Cesarean Delivery <i>AJP Reports</i> , 2022 , 12, e36-e40 | 1.2 | O |
| 211 | Patient Preferences for Outcomes Associated With Labor Epidural Analgesia <i>Cureus</i> , 2022 , 14, e22599 | 1.2 | O |
| 210 | Accuracy of visual estimation of blood loss in obstetrics using clinical reconstructions: an observational simulation cohort study <i>International Journal of Obstetric Anesthesia</i> , 2022 , 50, 103539 | 2.1 | |
| 209 | Development and validation of a Portuguese version of Obstetric Quality of Recovery-10 (ObsQoR-10-Portuguese) <i>Anaesthesia, Critical Care & Description Medicine</i> , 2022 , 101085 | 3 | O |
| 208 | Regional Anesthesia for Labor, Operative Vaginal Delivery and Cesarean Delivery: A Narrative Review. <i>Obstetric Anesthesia Digest</i> , 2021 , 41, 164-165 | 0.2 | |
| 207 | The Generalizability of Patients' Preferences and Concerns regarding Anesthesia Care for Cesarean Delivery: A Prospective Survey <i>Anesthesiology Research and Practice</i> , 2021 , 2021, 9002061 | 1.1 | |
| 206 | Superficial Cervical Plexus Block for Awake Large-Bore Central Line Placement in Parturients: A Case Series. <i>A&A Practice</i> , 2021 , 15, e01429 | 0.8 | O |
| 205 | Society for Obstetric Anesthesia and Perinatology: Consensus Statement and Recommendations for Enhanced Recovery After Cesarean. <i>Anesthesia and Analgesia</i> , 2021 , 132, 1362-1377 | 3.9 | 43 |
| 204 | Use of Patient-Reported Outcome Measures to Assess Outpatient Postpartum Recovery: A Systematic Review. <i>JAMA Network Open</i> , 2021 , 4, e2111600 | 10.4 | 9 |
| 203 | Integrated trajectories of the maternal metabolome, proteome, and immunome predict labor onset. <i>Science Translational Medicine</i> , 2021 , 13, | 17.5 | 17 |
| 202 | Quadratus lumborum block vs. transversus abdominis plane block for caesarean delivery: a systematic review and network meta-analysis. <i>Anaesthesia</i> , 2021 , 76, 393-403 | 6.6 | 20 |
| 201 | Comment on the editorial relating to: Transversus abdominis plane block compared with wound infiltration for postoperative analgesia following Cesarean delivery: a systematic review and network meta-analysis. <i>Canadian Journal of Anaesthesia</i> , 2021 , 68, 581-582 | 3 | 1 |
| 200 | Acute Pain Burden and Opioid Dose Requirements after Cesarean Delivery in Parturients with Preexisting Chronic Back Pain and Migraine. <i>Anesthesiology Research and Practice</i> , 2021 , 2021, 3305579 | 1.1 | |
| 199 | A systematic review of patient-reported outcome measures to assess postpartum pain using Consensus Based Standards for the Selection of Health Measurement Instruments (COSMIN) guidelines. <i>British Journal of Anaesthesia</i> , 2021 , 127, 264-274 | 5.4 | 7 |
| 198 | An observational pilot study of a novel loss of resistance syringe for locating the epidural space. <i>International Journal of Obstetric Anesthesia</i> , 2021 , 47, 102984 | 2.1 | 2 |
| 197 | Estimating Obstetric Anaesthesia Workload: Number of Deliveries Compared to Time-Based Workload <i>Turkish Journal of Anaesthesiology and Reanimation</i> , 2021 , 49, 292-297 | 0.7 | 1 |
| 196 | A randomized trial to investigate needle redirections/re-insertions using a handheld ultrasound device versus traditional palpation for spinal anesthesia in obese women undergoing cesarean delivery. <i>International Journal of Obstetric Anesthesia</i> , 2021 , 103229 | 2.1 | |

| 195 | A Peripheral Immune Signature of Labor Induction. Frontiers in Immunology, 2021, 12, 725989 | 8.4 | 4 |
|-----|--|------------------|----|
| 194 | Intravenous oxytocin dosing regimens for postpartum hemorrhage prevention following cesarean delivery: a systematic review and meta-analysis. <i>American Journal of Obstetrics and Gynecology</i> , 2021 , 225, 250.e1-250.e38 | 6.4 | 3 |
| 193 | Proposed domains for assessing postpartum recovery: a concept elicitation study. <i>BJOG: an International Journal of Obstetrics and Gynaecology</i> , 2021 , | 3.7 | 1 |
| 192 | Impact of enhanced recovery after cesarean delivery on maternal outcomes: A systematic review and meta-analysis. <i>Anaesthesia, Critical Care & Design Medicine</i> , 2021 , 40, 100935 | 3 | 4 |
| 191 | Evidence-based guidance for use of intrathecal morphine as an alternative to diamorphine for Caesarean delivery analgesia. <i>British Journal of Anaesthesia</i> , 2021 , 127, 501-505 | 5.4 | 2 |
| 190 | Regional anaesthesia for labour, operative vaginal delivery and caesarean delivery: a narrative review. <i>Anaesthesia</i> , 2021 , 76 Suppl 1, 136-147 | 6.6 | 10 |
| 189 | Evaluation of Domains of Patient-Reported Outcome Measures for Recovery After Childbirth: A Scoping and Systematic Review. <i>JAMA Network Open</i> , 2020 , 3, e205540 | 10.4 | 17 |
| 188 | Pain management during labor and vaginal birth. <i>Best Practice and Research in Clinical Obstetrics and Gynaecology</i> , 2020 , 67, 100-112 | 4.6 | 14 |
| 187 | Enhanced recovery after surgery for cesarean delivery. <i>Current Opinion in Obstetrics and Gynecology</i> , 2020 , 32, 113-120 | 2.4 | 5 |
| 186 | Enhanced recovery after caesarean delivery versus standard care studies: a systematic review of interventions and outcomes. <i>International Journal of Obstetric Anesthesia</i> , 2020 , 43, 72-86 | 2.1 | 18 |
| 185 | Comparison of spontaneous versus operative vaginal delivery using Obstetric Quality of Recovery-10 (ObsQoR-10): An observational cohort study. <i>Journal of Clinical Anesthesia</i> , 2020 , 63, 1097 | 8 ^{1.9} | 16 |
| 184 | Administration of Neuraxial Anesthesia in Adults with Pre-Existing Bleeding Disorders and Tendencies: Methodology for Delphi Consensus Recommendations. <i>Blood</i> , 2020 , 136, 29-29 | 2.2 | |
| 183 | Confounders Not General Anesthesia Likely Explain Greater Postpartum Depression. <i>Anesthesia and Analgesia</i> , 2020 , 131, e248-e250 | 3.9 | 2 |
| 182 | Antibiotic Prophylaxis for Cesarean Delivery: A Survey of Anesthesiologists. <i>Anesthesiology Research and Practice</i> , 2020 , 2020, 3741608 | 1.1 | O |
| 181 | Electrical stimulation test for epidural catheter placement after receiving intrathecal local anesthetic during a combined spinal-epidural technique for Cesarean delivery: confirmation of the spinal nerve root as epidural site of action. <i>Canadian Journal of Anaesthesia</i> , 2020 , 67, 498-499 | 3 | |
| 180 | Prevention of hypotension after spinal anaesthesia for caesarean section: a systematic review and network meta-analysis of randomised controlled trials. <i>Anaesthesia</i> , 2020 , 75, 109-121 | 6.6 | 29 |
| 179 | Virtual reality successfully provides anxiolysis to laboring women undergoing epidural placement. Journal of Clinical Anesthesia, 2020 , 61, 109635 | 1.9 | 2 |
| 178 | Prospective observational investigation of body habitus measurements and relationship to epidural depth in term pregnant women. <i>Acta Anaesthesiologica Scandinavica</i> , 2020 , 64, 677-684 | 1.9 | 1 |

| 177 | Conversion of labour epidural analgesia to surgical anaesthesia for emergency intrapartum Caesarean section. <i>BJA Education</i> , 2020 , 20, 26-31 | 1.2 | 3 |
|-----|--|------------------|----|
| 176 | Transversus abdominis plane block compared with wound infiltration for postoperative analgesia following Cesarean delivery: a systematic review and network meta-analysis. <i>Canadian Journal of Anaesthesia</i> , 2020 , 67, 1710-1727 | 3 | 17 |
| 175 | Re: Association between opioid use after intrapartum cesarean delivery and repeat cesarean delivery: a retrospective cohort study. <i>International Journal of Obstetric Anesthesia</i> , 2020 , 43, 30 | 2.1 | |
| 174 | Reply to: Re: enhanced recovery for cesarean section: beyond pain control. <i>International Journal of Obstetric Anesthesia</i> , 2020 , 44, 52 | 2.1 | |
| 173 | Transversus Abdominis Plane Block With Liposomal Bupivacaine for Pain After Cesarean Delivery in a Multicenter, Randomized, Double-Blind, Controlled Trial. <i>Anesthesia and Analgesia</i> , 2020 , 131, 1830-1 | 8 3 9 | 21 |
| 172 | Evaluation of inpatient postpartum recovery using the Obstetric Quality of Recovery-10 patient-reported outcome measure: a single-center observational study. <i>American Journal of Obstetrics & Samp; Gynecology MFM</i> , 2020 , 2, 100202 | 7.4 | 12 |
| 171 | Impact of intra-operative dexamethasone after scheduled cesarean delivery: a retrospective study. <i>International Journal of Obstetric Anesthesia</i> , 2020 , 41, 39-46 | 2.1 | 3 |
| 170 | Evaluation of the impact of enhanced recovery after surgery protocol implementation on maternal outcomes following elective cesarean delivery. <i>International Journal of Obstetric Anesthesia</i> , 2020 , 43, 39-46 | 2.1 | 24 |
| 169 | Hemodynamic profiles with and without left uterine displacement: A randomized study in term pregnancies receiving subarachnoid blockade for cesarean delivery. <i>Journal of Clinical Anesthesia</i> , 2020 , 64, 109796 | 1.9 | 2 |
| 168 | The comparative accuracy of a handheld and console ultrasound device for neuraxial depth and landmark assessment. <i>International Journal of Obstetric Anesthesia</i> , 2019 , 39, 68-73 | 2.1 | 8 |
| 167 | Differential Dynamics of the Maternal Immune System in Healthy Pregnancy and Preeclampsia. <i>Frontiers in Immunology</i> , 2019 , 10, 1305 | 8.4 | 25 |
| 166 | Alternative Magnesium Sulfate Dosing Regimens for Women With Preeclampsia: A Population Pharmacokinetic Exposure-Response Modeling and Simulation Study. <i>Journal of Clinical Pharmacology</i> , 2019 , 59, 1519-1526 | 2.9 | 4 |
| 165 | Personalized analgesic management for cesarean delivery. <i>International Journal of Obstetric Anesthesia</i> , 2019 , 40, 91-100 | 2.1 | 7 |
| 164 | Impact of patient choice for different postcesarean delivery analgesic protocols on opioid consumption: a randomized prospective clinical trial. <i>Regional Anesthesia and Pain Medicine</i> , 2019 , 44, 578-585 | 3.4 | 9 |
| 163 | Prospective Observational Investigation of Capnography and Pulse Oximetry Monitoring After Cesarean Delivery With Intrathecal Morphine. <i>Anesthesia and Analgesia</i> , 2019 , 128, 513-522 | 3.9 | 5 |
| 162 | Evaluation of the Obstetric Quality-of-Recovery score (ObsQoR-11) following non-elective caesarean delivery. <i>International Journal of Obstetric Anesthesia</i> , 2019 , 39, 51-59 | 2.1 | 19 |
| 161 | Assessing the Association Between Blood Loss and Postoperative Hemoglobin After Cesarean Delivery: A Prospective Study of 4 Blood Loss Measurement Modalities. <i>Anesthesia and Analgesia</i> , 2019 , 128, 926-932 | 3.9 | 13 |
| 160 | Development and evaluation of an obstetric quality-of-recovery score (ObsQoR-11) after elective Caesarean delivery. <i>British Journal of Anaesthesia</i> , 2019 , 122, 69-78 | 5.4 | 44 |

(2018-2019)

| 159 | Evaluation of patient-reported outcome measures of functional recovery following caesarean section: a systematic review using the consensus-based standards for the selection of health measurement instruments (COSMIN) checklist. <i>Anaesthesia</i> , 2019 , 74, 1439-1455 | 6.6 | 19 |
|-----|--|-----|----|
| 158 | Labor Epidural Analgesia to Cesarean Section Anesthetic Conversion Failure: A National Survey. <i>Anesthesiology Research and Practice</i> , 2019 , 2019, 6381792 | 1.1 | 7 |
| 157 | International consensus statement on the use of uterotonic agents during caesarean section. <i>Anaesthesia</i> , 2019 , 74, 1305-1319 | 6.6 | 45 |
| 156 | Pain relief during labour. <i>Lancet, The</i> , 2019 , 394, e12 | 40 | |
| 155 | Determinants of women's dissatisfaction with anaesthesia care in labour and delivery. <i>Anaesthesia</i> , 2019 , 74, 1112-1120 | 6.6 | 13 |
| 154 | Supraglottic Airway Rescue After Failed Fiberoptic Intubation in a Patient With Osteogenesis Imperfecta: A Case Report. <i>A&A Practice</i> , 2019 , 13, 7-9 | 0.8 | 3 |
| 153 | Transnasal Humidified Rapid-Insufflation Ventilatory Exchange for Elective Laryngeal Surgery During Pregnancy: A Case Report. <i>A&A Practice</i> , 2019 , 13, 426-429 | 0.8 | 1 |
| 152 | Intra-aortic Balloon Pump for Cesarean Hysterectomy and Massive Hemorrhage in a Parturient with Placenta Accreta and Pulmonary Embolus. <i>Journal of Medical Ultrasound</i> , 2019 , 27, 104-106 | 0.8 | 1 |
| 151 | The Crystalloid Co-Load: Clinically as Effective as Colloid Preload for Preventing Hypotension from Spinal Anaesthesia for Caesarean Delivery. <i>Turkish Journal of Anaesthesiology and Reanimation</i> , 2019 , 47, 35-40 | 0.7 | 2 |
| 150 | Preoperative Fasting Times for Patients Undergoing Caesarean Delivery: Before and After a Patient Educational Initiative. <i>Turkish Journal of Anaesthesiology and Reanimation</i> , 2019 , 47, 282-286 | 0.7 | 3 |
| 149 | In vitro intravenous fluid co-load rates with and without an intravenous fluid warming device. <i>International Journal of Obstetric Anesthesia</i> , 2019 , 38, 149-150 | 2.1 | |
| 148 | Evaluation of Opioid Use With Split Doses of Oral Opioids in a Postcesarean Delivery Analgesia Order Set. <i>Obstetrics and Gynecology</i> , 2019 , 134, 120-127 | 4.9 | 9 |
| 147 | Society for Obstetric Anesthesia and Perinatology Consensus Statement: Monitoring Recommendations for Prevention and Detection of Respiratory Depression Associated With Administration of Neuraxial Morphine for Cesarean Delivery Analgesia. <i>Anesthesia and Analgesia</i> , | 3.9 | 34 |
| 146 | Population Pharmacokinetic Modeling to Evaluate Standard Magnesium Sulfate Treatments and Alternative Dosing Regimens for Women With Preeclampsia. <i>Journal of Clinical Pharmacology</i> , 2019 , 59, 374-385 | 2.9 | 7 |
| 145 | Observational Study Using Ultrasound to Assess Midline Labor Epidural Analgesia Placement and Analgesic Efficacy. <i>Journal of Ultrasound in Medicine</i> , 2018 , 37, 1693-1699 | 2.9 | 1 |
| 144 | Does the addition of active body warming to in-line intravenous fluid warming prevent maternal hypothermia during elective caesarean section? A randomised controlled trial. <i>International Journal of Obstetric Anesthesia</i> , 2018 , 35, 115-116 | 2.1 | 1 |
| 143 | High-dose versus low-dose local anaesthetic for transversus abdominis plane block post-Caesarean delivery analgesia: a meta-analysis. <i>British Journal of Anaesthesia</i> , 2018 , 120, 252-263 | 5.4 | 30 |
| 142 | Left Uterine Tilt for Cesarean Delivery Significantly Improves Maternal Hemodynamics and Should Not Be Considered Outdated Dogma. <i>Anesthesiology</i> , 2018 , 128, 858-859 | 4.3 | 3 |

| 141 | The Effect of Glycopyrrolate on the Incidence of Hypotension and Vasopressor Requirement During Spinal Anesthesia for Cesarean Delivery: A Meta-analysis. <i>Anesthesia and Analgesia</i> , 2018 , 126, 552-558 | 3.9 | 6 |
|-----|--|----------------|-----|
| 140 | International consensus statement on the management of hypotension with vasopressors during caesarean section under spinal anaesthesia. <i>Anaesthesia</i> , 2018 , 73, 71-92 | 6.6 | 194 |
| 139 | The Accuracy of a Handheld Ultrasound Device for Neuraxial Depth and Landmark Assessment: A Prospective Cohort Trial. <i>Anesthesia and Analgesia</i> , 2018 , 126, 1995-1998 | 3.9 | 18 |
| 138 | Preoperative anterior thigh temperature does not correlate with perioperative temporal hypothermia during cesarean delivery with spinal anesthesia: Secondary analysis of a randomized control trial. <i>International Journal of Obstetric Anesthesia</i> , 2018 , 33, 40-45 | 2.1 | 4 |
| 137 | Prediction of outliers in pain, analgesia requirement, and recovery of function after childbirth: a prospective observational cohort study. <i>British Journal of Anaesthesia</i> , 2018 , 121, 417-426 | 5.4 | 16 |
| 136 | Modern techniques to optimize neuraxial labor analgesia. <i>Anesthesia and Pain Medicine</i> , 2018 , 13, 233-2 | . 40 .3 | 2 |
| 135 | The Society for Obstetric Anesthesia and Perinatology Consensus Statement on the Anesthetic Management of Pregnant and Postpartum Women Receiving Thromboprophylaxis or Higher Dose Anticoagulants. <i>Anesthesia and Analgesia</i> , 2018 , 126, 928-944 | 3.9 | 107 |
| 134 | International Consensus Statement on the Management of Hypotension With Vasopressors During Caesarean Section Under Spinal Anaesthesia. <i>Obstetric Anesthesia Digest</i> , 2018 , 38, 171-172 | 0.2 | 7 |
| 133 | A Systematic Review Evaluating Neuraxial Morphine and Diamorphine-Associated Respiratory Depression After Cesarean Delivery. <i>Anesthesia and Analgesia</i> , 2018 , 127, 1385-1395 | 3.9 | 25 |
| 132 | Correlation of changes in hemodynamic response as measured by cerebral optical spectrometry with subjective pain ratings in volunteers and patients: a prospective cohort study. <i>Journal of Pain Research</i> , 2018 , 11, 1991-1998 | 2.9 | O |
| 131 | A Systematic Review of Postoperative Pain Outcome Measurements Utilised in Regional Anesthesia Randomized Controlled Trials. <i>Anesthesiology Research and Practice</i> , 2018 , 2018, 9050239 | 1.1 | 7 |
| 130 | In reply. International Journal of Obstetric Anesthesia, 2017, 30, 76 | 2.1 | |
| 129 | Postcesarean delivery analgesia. <i>Bailliereg Best Practice and Research in Clinical Anaesthesiology</i> , 2017 , 31, 69-79 | 4 | 69 |
| 128 | Effect of a High-Rate Versus a Low-Rate Oxytocin Infusion for Maintaining Uterine Contractility During Elective Cesarean Delivery: A Prospective Randomized Clinical Trial. <i>Anesthesia and Analgesia</i> , 2017 , 124, 857-862 | 3.9 | 19 |
| 127 | Essentials of Our Current Understanding: Abdominal Wall Blocks. <i>Regional Anesthesia and Pain Medicine</i> , 2017 , 42, 133-183 | 3.4 | 208 |
| 126 | Nitrous oxide for labor analgesia: Utilization and predictors of conversion to neuraxial analgesia. Journal of Clinical Anesthesia, 2017 , 40, 40-45 | 1.9 | 15 |
| 125 | Patient choice compared with no choice of intrathecal morphine dose for caesarean analgesia: a randomized clinical trial. <i>British Journal of Anaesthesia</i> , 2017 , 118, 762-771 | 5.4 | 18 |
| 124 | A Perspective on Hypercapnia Events After Cesarean Delivery in Women Receiving Intrathecal Morphine. <i>Anesthesia and Analgesia</i> , 2017 , 125, 355-356 | 3.9 | |

(2017-2017)

| 123 | Neuraxial Anesthesia in Obstetric Patients Receiving Thromboprophylaxis With Unfractionated or Low-Molecular-Weight Heparin: A Systematic Review of Spinal Epidural Hematoma. <i>Anesthesia and Analgesia</i> , 2017 , 125, 223-231 | 3.9 | 32 |
|-----|---|---------------------|----------------|
| 122 | Optimization of Maternal Magnesium Sulfate Administration for Fetal Neuroprotection: Application of a Prospectively Constructed Pharmacokinetic Model to the BEAM Cohort. <i>Journal of Clinical Pharmacology</i> , 2017 , 57, 1419-1424 | 2.9 | 9 |
| 121 | Analysis of Physiological Respiratory Variable Alarm Alerts Among Laboring Women Receiving Remifentanil. <i>Anesthesia and Analgesia</i> , 2017 , 124, 1211-1218 | 3.9 | 21 |
| 120 | What's trending now? An analysis of trends in internet searches for labor epidurals. <i>International Journal of Obstetric Anesthesia</i> , 2017 , 30, 52-57 | 2.1 | 10 |
| 119 | Optimal administration of cefazolin prophylaxis for cesarean delivery. <i>Journal of Perinatology</i> , 2017 , 37, 16-20 | 3.1 | 4 |
| 118 | Optimal Pain Management After Cesarean Delivery. <i>Anesthesiology Clinics</i> , 2017 , 35, 107-124 | 2.3 | 74 |
| 117 | Postpartum hemorrhage following vaginal delivery: risk factors and maternal outcomes. <i>Journal of Perinatology</i> , 2017 , 37, 243-248 | 3.1 | 7 |
| 116 | Recovery after Nulliparous Birth: A Detailed Analysis of Pain Analgesia and Recovery of Function. <i>Anesthesiology</i> , 2017 , 127, 684-694 | 4.3 | 34 |
| 115 | Programmed Intermittent Epidural Boluses (PIEB) for Maintenance of Labor Analgesia: An Incremental Step Before the Next Paradigm Shift?. <i>Turkish Journal of Anaesthesiology and Reanimation</i> , 2017 , 45, 73-75 | 0.7 | 2 |
| 114 | Programmed Intermittent Epidural Boluses (PIEB) for Maintenance of Labor Analgesia: A Superior Technique to Continuous Epidural Infusion?. <i>Turkish Journal of Anaesthesiology and Reanimation</i> , 2017 , 45, 65-66 | 0.7 | 8 |
| 113 | Rectus Muscle Reapproximation at Cesarean Delivery and Postoperative Pain: A Randomized Controlled Trial. <i>The Surgery Journal</i> , 2017 , 3, e128-e133 | 0.9 | 5 |
| 112 | Comparative Effectiveness of Lower Leg Compression Devices Versus Sequential Compression Devices to Prevent Postspinal Hypotension During Cesarean Delivery. <i>Anesthesia and Analgesia</i> , 2017 , 124, 696-697 | 3.9 | 4 |
| 111 | . International Journal of Obstetric Anesthesia, 2017 , 32, 93-94 | 2.1 | |
| 110 | General anaesthesia for caesarean section: is the end in sight for thiopental?. <i>British Journal of Hospital Medicine (London, England: 2005)</i> , 2017 , 78, 358 | 0.8 | 1 |
| 109 | Intrathecal Migration of an Epidural Catheter While Using a Programmed Intermittent Epidural Bolus Technique for Labor Analgesia Maintenance: A Case Report. A & A Case Reports, 2017, 9, 357-359 | | 3 |
| 108 | Optimising triggers for patient-assisted remifentanil analgesia during labour. <i>Anaesthesia</i> , 2017 , 72, 115 | 5 6. d15 | 62 |
| 107 | Comparative Echogenicity of an Epidural Catheter and 2 New Catheters Designed for Ultrasound-Guided Continuous Peripheral Nerve Blocks. <i>Journal of Ultrasound in Medicine</i> , 2017 , 36, 257 | 7 7 -257 | 6 ¹ |
| 106 | Clinical and microbiological features of maternal sepsis: a retrospective study. <i>International Journal of Obstetric Anesthesia</i> , 2017 , 29, 26-33 | 2.1 | 8 |

| 105 | Ambient operating room temperature: mother, baby or surgeon?. <i>British Journal of Anaesthesia</i> , 2017 , 119, 839 | 5.4 | 4 |
|-----|---|-----|----|
| 104 | Postpartum tubal ligation: A retrospective review of anesthetic management at a single institution and a practice survey of academic institutions. <i>Journal of Clinical Anesthesia</i> , 2017 , 43, 39-46 | 1.9 | 3 |
| 103 | Checklists and multidisciplinary team performance during simulated obstetric hemorrhage. <i>International Journal of Obstetric Anesthesia</i> , 2016 , 25, 9-16 | 2.1 | 22 |
| 102 | Survey of external cephalic version for breech presentation and neuraxial blockade use. <i>Journal of Clinical Anesthesia</i> , 2016 , 34, 616-22 | 1.9 | 2 |
| 101 | In reply. International Journal of Obstetric Anesthesia, 2016 , 27, 92-3 | 2.1 | |
| 100 | Mapping the Fetomaternal Peripheral Immune System at Term Pregnancy. <i>Journal of Immunology</i> , 2016 , 197, 4482-4492 | 5.3 | 25 |
| 99 | Amniotic fluid embolism: update and review. Current Opinion in Anaesthesiology, 2016, 29, 288-96 | 2.9 | 33 |
| 98 | Programmed intermittent epidural boluses for maintenance of labor analgesia: an impact study. <i>International Journal of Obstetric Anesthesia</i> , 2016 , 26, 32-8 | 2.1 | 35 |
| 97 | Pharmacokinetics and placental transfer of magnesium sulfate in pregnant women. <i>American Journal of Obstetrics and Gynecology</i> , 2016 , 214, 737.e1-9 | 6.4 | 24 |
| 96 | Remifentanil for labor analgesia: an evidence-based narrative review. <i>International Journal of Obstetric Anesthesia</i> , 2016 , 25, 66-74 | 2.1 | 51 |
| 95 | A Prospective Cohort Study Evaluating the Ability of Anticipated Pain, Perceived Analgesic Needs, and Psychological Traits to Predict Pain and Analgesic Usage following Cesarean Delivery. <i>Anesthesiology Research and Practice</i> , 2016 , 2016, 7948412 | 1.1 | 12 |
| 94 | Pharmacokinetics and Pharmacodynamics of Drugs Commonly Used in Pregnancy and Parturition. <i>Anesthesia and Analgesia</i> , 2016 , 122, 786-804 | 3.9 | 42 |
| 93 | The Effect of Intrathecal Morphine Dose on Outcomes After Elective Cesarean Delivery: A Meta-Analysis. <i>Anesthesia and Analgesia</i> , 2016 , 123, 154-64 | 3.9 | 85 |
| 92 | Active Warming Utilizing Combined IV Fluid and Forced-Air Warming Decreases Hypothermia and Improves Maternal Comfort During Cesarean Delivery: A Randomized Control Trial. <i>Anesthesia and Analgesia</i> , 2016 , 122, 1490-7 | 3.9 | 53 |
| 91 | Implementation of Programmed Intermittent Epidural Bolus for the Maintenance of Labor Analgesia. <i>Anesthesia and Analgesia</i> , 2016 , 123, 965-71 | 3.9 | 49 |
| 90 | In reply. International Journal of Obstetric Anesthesia, 2016 , 27, 90 | 2.1 | O |
| 89 | Challenging the 4- to 5-minute rule: from perimortem cesarean to resuscitative hysterotomy. <i>American Journal of Obstetrics and Gynecology</i> , 2016 , 215, 129-31 | 6.4 | 4 |
| 88 | The impact of postpartum hemoglobin levels on maternal quality of life after delivery: a prospective exploratory study. <i>Annals of Hematology</i> , 2016 , 95, 2049-2055 | 3 | 11 |

| 87 | In reply. International Journal of Obstetric Anesthesia, 2016 , 28, 96 | 2.1 | |
|----|---|-------------------|-----|
| 86 | Methylnaltrexone to prevent intrathecal morphine-induced pruritus after Caesarean delivery: a multicentre, randomized clinical trial. <i>British Journal of Anaesthesia</i> , 2015 , 114, 469-76 | 5.4 | 10 |
| 85 | Scheduled acetaminophen with as-needed opioids compared to as-needed acetaminophen plus opioids for post-cesarean pain management. <i>International Journal of Obstetric Anesthesia</i> , 2015 , 24, 210 | 0-6 ⁻¹ | 64 |
| 84 | Cardiac Arrest in Pregnancy: A Scientific Statement From the American Heart Association. <i>Circulation</i> , 2015 , 132, 1747-73 | 16.7 | 198 |
| 83 | The Effect of patient warming during Caesarean delivery on maternal and neonatal outcomes: a meta-analysis. <i>British Journal of Anaesthesia</i> , 2015 , 115, 500-10 | 5.4 | 58 |
| 82 | Utilization of spinal anesthesia for external cephalic version: a clinical practice cohort analysis. <i>International Journal of Obstetric Anesthesia</i> , 2015 , 24, 389-90 | 2.1 | |
| 81 | Drug labeling in the practice of obstetric anesthesia. <i>American Journal of Obstetrics and Gynecology</i> , 2015 , 212, 24-7 | 6.4 | 6 |
| 80 | Magnesium sulfate exposure and neonatal intensive care unit admission at term. <i>Journal of Perinatology</i> , 2015 , 35, 181-5 | 3.1 | 4 |
| 79 | Survey of Accepted Practice following Failed Intubation for Emergency Caesarean Delivery. <i>Anesthesiology Research and Practice</i> , 2015 , 2015, 192315 | 1.1 | 2 |
| 78 | Second-line uterotonics and the risk of hemorrhage-related morbidity. <i>American Journal of Obstetrics and Gynecology</i> , 2015 , 212, 642.e1-7 | 6.4 | 28 |
| 77 | The impact of breastfeeding on postpartum pain after vaginal and cesarean delivery. <i>Journal of Clinical Anesthesia</i> , 2015 , 27, 33-8 | 1.9 | 11 |
| 76 | Preferred spoken language mediates differences in neuraxial labor analgesia utilization among racial and ethnic groups. <i>International Journal of Obstetric Anesthesia</i> , 2014 , 23, 161-7 | 2.1 | 9 |
| 75 | In reply. Journal of Hand Surgery, 2014 , 39, 1888-9 | 2.6 | |
| 74 | Risk factors for obstetric morbidity in patients with uterine atony undergoing caesarean delivery. <i>British Journal of Anaesthesia</i> , 2014 , 113, 661-8 | 5.4 | 26 |
| 73 | Pharmacokinetics of prophylactic cefazolin in parturients undergoing cesarean delivery. <i>Antimicrobial Agents and Chemotherapy</i> , 2014 , 58, 3504-13 | 5.9 | 30 |
| 72 | The dilemma of vaginal breech delivery worldwide. <i>Lancet, The</i> , 2014 , 384, 1183 | 40 | 1 |
| 71 | Incidence of failure of continuous peripheral nerve catheters for postoperative analgesia in upper extremity surgery. <i>Journal of Hand Surgery</i> , 2014 , 39, 324-9 | 2.6 | 46 |
| 70 | Evaluation of a prospectively administered written questionnaire to reduce the incidence of suspected latex anaphylaxis during elective cesarean delivery. <i>International Journal of Obstetric Apesthesia</i> 2014 23 335-40 | 2.1 | 5 |

| 69 | In response. Anesthesia and Analgesia, 2014 , 118, 884-5 | 3.9 | |
|----|--|-----|-----|
| 68 | The Society for Obstetric Anesthesia and Perinatology consensus statement on the management of cardiac arrest in pregnancy. <i>Anesthesia and Analgesia</i> , 2014 , 118, 1003-16 | 3.9 | 108 |
| 67 | Prospective longitudinal cohort questionnaire assessment of labouring women's preference both pre- and post-delivery for either reduced pain intensity for a longer duration or greater pain intensity for a shorter duration. <i>British Journal of Anaesthesia</i> , 2014 , 113, 468-73 | 5.4 | 13 |
| 66 | Strategies to Optimize Pain Control Following Cesarean Delivery. <i>Refresher Courses in Anesthesiology</i> , 2014 , 42, 23-30 | | |
| 65 | "Pseudo" shearing of a peripheral nerve catheter after interscalene block. <i>Regional Anesthesia and Pain Medicine</i> , 2014 , 39, 556-7 | 3.4 | |
| 64 | A prospective observational study evaluating the ability of prelabor psychological tests to predict labor pain, epidural analgesic consumption, and maternal satisfaction. <i>Anesthesia and Analgesia</i> , 2014 , 119, 632-640 | 3.9 | 25 |
| 63 | Transversus abdominis plane blocks for rescue analgesia following Cesarean delivery: a case series. <i>Canadian Journal of Anaesthesia</i> , 2013 , 60, 299-303 | 3 | 25 |
| 62 | The effect of low concentrations versus high concentrations of local anesthetics for labour analgesia on obstetric and anesthetic outcomes: a meta-analysis. <i>Canadian Journal of Anaesthesia</i> , 2013 , 60, 840-54 | 3 | 68 |
| 61 | Postoperative subcutaneous instillation of low-dose ketorolac but not hydromorphone reduces wound exudate concentrations of interleukin-6 and interleukin-10 and improves analgesia following cesarean delivery. <i>Journal of Pain</i> , 2013 , 14, 48-56 | 5.2 | 22 |
| 60 | Intrathecal morphine 100 and 200 g for post-cesarean delivery analgesia: a trade-off between analgesic efficacy and side effects. <i>International Journal of Obstetric Anesthesia</i> , 2013 , 22, 36-41 | 2.1 | 35 |
| 59 | Peripartum outcomes and anaesthetic management of parturients with moderate to complex congenital heart disease or pulmonary hypertension*. <i>Anaesthesia</i> , 2013 , 68, 52-9 | 6.6 | 19 |
| 58 | Evaluation of experimental pain tests to predict labour pain and epidural analgesic consumption. <i>British Journal of Anaesthesia</i> , 2013 , 110, 600-6 | 5.4 | 14 |
| 57 | Response times for emergency cesarean delivery: use of simulation drills to assess and improve obstetric team performance. <i>Journal of Perinatology</i> , 2013 , 33, 259-63 | 3.1 | 27 |
| 56 | Transport decreases the quality of cardiopulmonary resuscitation during simulated maternal cardiac arrest. <i>Anesthesia and Analgesia</i> , 2013 , 116, 162-7 | 3.9 | 34 |
| 55 | Brief report: a cost analysis of neuraxial anesthesia to facilitate external cephalic version for breech fetal presentation. <i>Anesthesia and Analgesia</i> , 2013 , 117, 155-9 | 3.9 | 15 |
| 54 | The effect of OPRM1 and COMT genotypes on the analgesic response to intravenous fentanyl labor analgesia. <i>Anesthesia and Analgesia</i> , 2013 , 116, 386-91 | 3.9 | 29 |
| 53 | Does intrathecal analgesia reduce the subsequent minimum local analgesic concentration of epidural bupivacaine?. <i>Anaesthesia</i> , 2012 , 67, 1054-5; author reply 1055-6 | 6.6 | |
| 52 | Failed epidural top-up for cesarean delivery for failure to progress in labor: the case against single-shot spinal anesthesia. <i>International Journal of Obstetric Anesthesia</i> , 2012 , 21, 357-9 | 2.1 | 16 |

(2010-2012)

| 51 | Intrathecal fentanyl added to bupivacaine and morphine for cesarean delivery may induce a subtle acute opioid tolerance. <i>International Journal of Obstetric Anesthesia</i> , 2012 , 21, 29-34 | 2.1 | 21 |
|----|--|------|----|
| 50 | Can We Predict Postoperative Pain Prior to Patients Undergoing Surgery?. <i>Journal of Pain & Relief</i> , 2012 , 01, | О | 6 |
| 49 | Non-invasive haemoglobin measurement in patients undergoing elective Caesarean section. <i>British Journal of Anaesthesia</i> , 2012 , 108, 271-7 | 5.4 | 54 |
| 48 | Collecting and measuring wound exudate biochemical mediators in surgical wounds. <i>Journal of Visualized Experiments</i> , 2012 , | 1.6 | 2 |
| 47 | Prior Epidural Lidocaine Alters the Pharmacokinetics and Drug Effects of Extended-Release Epidural Morphine (DepoDur¶After Cesarean Delivery. <i>Survey of Anesthesiology</i> , 2012 , 56, 298-299 | | |
| 46 | Neuraxial morphine and respiratory depression: finding the right balance. <i>Drugs</i> , 2011 , 71, 1807-19 | 12.1 | 83 |
| 45 | Neuraxial blockade for external cephalic version: a systematic review. <i>International Journal of Obstetric Anesthesia</i> , 2011 , 20, 299-306 | 2.1 | 27 |
| 44 | ED(50) and ED(95) of intrathecal bupivacaine in morbidly obese patients undergoing cesarean delivery. <i>Anesthesiology</i> , 2011 , 114, 529-35 | 4.3 | 66 |
| 43 | Labor room setting compared with the operating room for simulated perimortem cesarean delivery: a randomized controlled trial. <i>Obstetrics and Gynecology</i> , 2011 , 118, 1090-1094 | 4.9 | 32 |
| 42 | Prior epidural lidocaine alters the pharmacokinetics and drug effects of extended-release epidural morphine (DepoDur□) after cesarean delivery. <i>Anesthesia and Analgesia</i> , 2011 , 113, 251-8 | 3.9 | 9 |
| 41 | A sensitive assay for the quantification of morphine and its active metabolites in human plasma and dried blood spots using high-performance liquid chromatography-tandem mass spectrometry. <i>Analytical and Bioanalytical Chemistry</i> , 2011 , 400, 715-28 | 4.4 | 65 |
| 40 | Minimum effective bolus dose of oxytocin during elective Caesarean delivery. <i>British Journal of Anaesthesia</i> , 2010 , 104, 338-43 | 5.4 | 95 |
| 39 | Survey of the Factors Associated with a Woman's Choice to Have an Epidural for Labor Analgesia. <i>Anesthesiology Research and Practice</i> , 2010 , 2010, | 1.1 | 27 |
| 38 | Continuous subcutaneous instillation of bupivacaine compared to saline reduces interleukin 10 and increases substance P in surgical wounds after cesarean delivery. <i>Anesthesia and Analgesia</i> , 2010 , 111, 1452-9 | 3.9 | 46 |
| 37 | Analgesic requirements and postoperative recovery after scheduled compared to unplanned cesarean delivery: a retrospective chart review. <i>International Journal of Obstetric Anesthesia</i> , 2010 , 19, 10-5 | 2.1 | 17 |
| 36 | Neuraxial anesthesia in obstetric patients receiving anticoagulant and antithrombotic drugs. <i>International Journal of Obstetric Anesthesia</i> , 2010 , 19, 193-201 | 2.1 | 24 |
| 35 | Retrospective analysis of anesthetic interventions for obese patients undergoing elective cesarean delivery. <i>Journal of Clinical Anesthesia</i> , 2010 , 22, 519-26 | 1.9 | 21 |
| 34 | The role of interleukin-1 in wound biology. Part II: In vivo and human translational studies. Anesthesia and Analgesia, 2010 , 111, 1534-42 | 3.9 | 67 |

| 33 | The role of interleukin-1 in wound biology. Part I: Murine in silico and in vitro experimental analysis. <i>Anesthesia and Analgesia</i> , 2010 , 111, 1525-33 | 3.9 | 29 |
|----|--|-----|-----|
| 32 | Effect of a preemptive femoral nerve block on cytokine release and hyperalgesia in experimentally inflamed skin of human volunteers. <i>Regional Anesthesia and Pain Medicine</i> , 2010 , 35, 514-9 | 3.4 | 11 |
| 31 | Cost-effectiveness of external cephalic version for term breech presentation. <i>BMC Pregnancy and Childbirth</i> , 2010 , 10, 3 | 3.2 | 45 |
| 30 | Deficits in the provision of cardiopulmonary resuscitation during simulated obstetric crises. <i>American Journal of Obstetrics and Gynecology</i> , 2010 , 203, 179.e1-5 | 6.4 | 65 |
| 29 | A survey of perioperative and postoperative anesthetic practices for cesarean delivery. <i>Anesthesiology Research and Practice</i> , 2009 , 2009, 510642 | 1.1 | 35 |
| 28 | Hetastarch co-loading is as effective as pre-loading for the prevention of hypotension following spinal anesthesia for cesarean delivery. <i>International Journal of Obstetric Anesthesia</i> , 2009 , 18, 150-5 | 2.1 | 37 |
| 27 | A randomized trial of maximum cephalad sensory blockade with single-shot spinal compared with combined spinal-epidural techniques for cesarean delivery. <i>Anesthesia and Analgesia</i> , 2009 , 108, 240-5 | 3.9 | 15 |
| 26 | Patient-controlled epidural analgesia for labor. Anesthesia and Analgesia, 2009, 108, 921-8 | 3.9 | 116 |
| 25 | Postoperative Analgesia: Epidural and Spinal Techniques 2009 , 593-630 | | 1 |
| 24 | Vaginal twin delivery: a survey and review of location, anesthesia coverage and interventions. <i>International Journal of Obstetric Anesthesia</i> , 2008 , 17, 212-6 | 2.1 | 6 |
| 23 | Assessment of knowledge regarding cardiopulmonary resuscitation of pregnant women. <i>International Journal of Obstetric Anesthesia</i> , 2008 , 17, 20-5 | 2.1 | 66 |
| 22 | Cytokine profile in human skin in response to experimental inflammation, noxious stimulation, and administration of a COX-inhibitor: a microdialysis study. <i>Pain</i> , 2008 , 139, 15-27 | 8 | 84 |
| 21 | Local and systemic release of cytokines, nerve growth factor, prostaglandin E2, and substance P in incisional wounds and serum following cesarean delivery. <i>Journal of Pain</i> , 2008 , 9, 650-7 | 5.2 | 47 |
| 20 | Collecting and measuring nociceptive and inflammatory mediators in surgical wounds. <i>Journal of Visualized Experiments</i> , 2008 , | 1.6 | 2 |
| 19 | Respiratory depression after neuraxial opioids in the obstetric setting. <i>Anesthesia and Analgesia</i> , 2008 , 107, 956-61 | 3.9 | 108 |
| 18 | Single-dose, extended-release epidural morphine (DepoDur) compared to conventional epidural morphine for post-cesarean pain. <i>Anesthesia and Analgesia</i> , 2007 , 105, 176-83 | 3.9 | 72 |
| 17 | The effect of colloid and crystalloid preloading on thromboelastography prior to Cesarean delivery. <i>Canadian Journal of Anaesthesia</i> , 2007 , 54, 190-5 | 3 | 42 |
| 16 | Intraoperative forced air-warming during cesarean delivery under spinal anesthesia does not prevent maternal hypothermia. <i>Anesthesia and Analgesia</i> , 2007 , 105, 1413-9, table of contents | 3.9 | 77 |

LIST OF PUBLICATIONS

| 15 | Extended Release Epidural Morphine, Far from Ideal for Postcesarean Delivery Pain Control. <i>Anesthesia and Analgesia</i> , 2007 , 105, 1864-1865 | 3.9 | |
|----|---|-----|-----|
| 14 | The Episure syringe: a novel loss of resistance syringe for locating the epidural space. <i>Anesthesia and Analgesia</i> , 2007 , 105, 1164-6, table of contents | 3.9 | 23 |
| 13 | Fetal oxygen saturation after combined spinal-epidural labor analgesia: a case series. <i>Journal of Clinical Anesthesia</i> , 2007 , 19, 476-8 | 1.9 | 9 |
| 12 | Local infiltration of epinephrine-containing lidocaine with bicarbonate reduces superficial bleeding and pain during labor epidural catheter insertion: a randomized trial. <i>International Journal of Obstetric Anesthesia</i> , 2007 , 16, 116-21 | 2.1 | 16 |
| 11 | A survey of labor patient-controlled epidural anesthesia practice in California hospitals. <i>International Journal of Obstetric Anesthesia</i> , 2006 , 15, 217-22 | 2.1 | 25 |
| 10 | Epidural anesthesia for elective cesarean delivery with intraoperative arterial occlusion balloon catheter placement. <i>Anesthesia and Analgesia</i> , 2006 , 102, 585-7 | 3.9 | 26 |
| 9 | Experimental heat pain for detecting pregnancy-induced analgesia in humans. <i>Anesthesia and Analgesia</i> , 2006 , 103, 1283-7 | 3.9 | 31 |
| 8 | Valdecoxib for postoperative pain management after cesarean delivery: a randomized, double-blind, placebo-controlled study. <i>Anesthesia and Analgesia</i> , 2006 , 103, 664-70 | 3.9 | 32 |
| 7 | "Ultra-light" patient-controlled epidural analgesia during labor: effects of varying regimens on analgesia and physician workload. <i>International Journal of Obstetric Anesthesia</i> , 2005 , 14, 223-9 | 2.1 | 34 |
| 6 | Single-dose, sustained-release epidural morphine in the management of postoperative pain after elective cesarean delivery: results of a multicenter randomized controlled study. <i>Anesthesia and Analgesia</i> , 2005 , 100, 1150-1158 | 3.9 | 93 |
| 5 | Patient preferences for anesthesia outcomes associated with cesarean delivery. <i>Anesthesia and Analgesia</i> , 2005 , 101, 1182-1187 | 3.9 | 115 |
| 4 | The ED50 and ED95 of intrathecal isobaric bupivacaine with opioids for cesarean delivery. <i>Anesthesiology</i> , 2005 , 103, 606-12 | 4.3 | 60 |
| 3 | Neonatal chest wall rigidity following the use of remifentanil for cesarean delivery in a patient with autoimmune hepatitis and thrombocytopenia. <i>International Journal of Obstetric Anesthesia</i> , 2004 , 13, 53-6 | 2.1 | 28 |
| 2 | 'Near-miss' hyperkalaemic cardiac arrest associated with rapid blood transfusion. <i>Anaesthesia</i> , 1999 54, 1094-6 | 6.6 | 19 |

Structural heart disease in pregnant women1-28