Marson Putra

List of Publications by Year in descending order

Source: https://exaly.com/author-pdf/4755409/publications.pdf

Version: 2024-02-01

1162367 1372195 10 272 8 10 citations h-index g-index papers 10 10 10 206 citing authors docs citations times ranked all docs

#	Article	IF	CITATIONS
1	Saracatinib, a Src Tyrosine Kinase Inhibitor, as a Disease Modifier in the Rat DFP Model: Sex Differences, Neurobehavior, Gliosis, Neurodegeneration, and Nitro-Oxidative Stress. Antioxidants, 2022, 11, 61.	2.2	15
2	DFP-Induced Status Epilepticus Severity in Mixed-Sex Cohorts of Adult Rats Housed in the Same Room: Behavioral and EEG Comparisons. Frontiers in Cell and Developmental Biology, 2022, 10, .	1.8	12
3	Differential Impact of Severity and Duration of Status Epilepticus, Medical Countermeasures, and a Disease-Modifier, Saracatinib, on Brain Regions in the Rat Diisopropylfluorophosphate Model. Frontiers in Cellular Neuroscience, 2021, 15, 772868.	1.8	15
4	Soman (GD) Rat Model to Mimic Civilian Exposure to Nerve Agent: Mortality, Video-EEG Based Status Epilepticus Severity, Sex Differences, Spontaneously Recurring Seizures, and Brain Pathology. Frontiers in Cellular Neuroscience, 2021, 15, 798247.	1.8	10
5	Inducible nitric oxide synthase inhibitor, 1400W, mitigates DFP-induced long-term neurotoxicity in the rat model. Neurobiology of Disease, 2020, 133, 104443.	2.1	39
6	Fyn-tau Ablation Modifies PTZ-Induced Seizures and Post-seizure Hallmarks of Early Epileptogenesis. Frontiers in Cellular Neuroscience, 2020, 14, 592374.	1.8	24
7	Sex as a biological variable in the rat model of diisopropylfluorophosphateâ€induced longâ€ŧerm neurotoxicity. Annals of the New York Academy of Sciences, 2020, 1479, 44-64.	1.8	23
8	Diapocynin, an NADPH oxidase inhibitor, counteracts diisopropylfluorophosphateâ€induced longâ€term neurotoxicity in the rat model. Annals of the New York Academy of Sciences, 2020, 1479, 75-93.	1.8	25
9	Role of the Fyn-PKCδ signaling in SE-induced neuroinflammation and epileptogenesis in experimental models of temporal lobe epilepsy. Neurobiology of Disease, 2018, 110, 102-121.	2.1	50
10	1400W, a highly selective inducible nitric oxide synthase inhibitor is a potential disease modifier in the rat kainate model of temporal lobe epilepsy. Neurobiology of Disease, 2016, 93, 184-200.	2.1	59