

Adam J Popchak

List of Publications by Year in descending order

Source: <https://exaly.com/author-pdf/4756982/publications.pdf>

Version: 2024-02-01

36
papers

683
citations

567281

15
h-index

580821

25
g-index

37
all docs

37
docs citations

37
times ranked

568
citing authors

#	ARTICLE	IF	CITATIONS
1	An Increased Lateral Femoral Condyle Ratio Is a Risk Factor for Anterior Cruciate Ligament Injury. <i>Journal of Bone and Joint Surgery - Series A</i> , 2018, 100, 857-864.	3.0	80
2	MRI can accurately detect meniscal ramp lesions of the knee. <i>Knee Surgery, Sports Traumatology, Arthroscopy</i> , 2017, 25, 3955-3960.	4.2	62
3	Distal femur morphology affects rotatory knee instability in patients with anterior cruciate ligament ruptures. <i>Knee Surgery, Sports Traumatology, Arthroscopy</i> , 2019, 27, 1514-1519.	4.2	40
4	Static Lateral Tibial Plateau Subluxation Predicts High-Grade Rotatory Knee Laxity in Anterior Cruciate Ligament-Deficient Knees. <i>American Journal of Sports Medicine</i> , 2019, 47, 277-284.	4.2	37
5	Non-anatomic tunnel position increases the risk of revision anterior cruciate ligament reconstruction. <i>Knee Surgery, Sports Traumatology, Arthroscopy</i> , 2022, 30, 1388-1395.	4.2	35
6	Factors Related to Injury in Youth and Adolescent Baseball Pitching, with an Eye Toward Prevention. <i>American Journal of Physical Medicine and Rehabilitation</i> , 2015, 94, 395-409.	1.4	31
7	Complications After Arthroscopic Shoulder Surgery: A Review of the American Board of Orthopaedic Surgery Database. <i>Journal of the American Academy of Orthopaedic Surgeons Global Research and Reviews</i> , 2018, 2, e093.	0.7	27
8	High-grade rotatory knee laxity may be predictable in ACL injuries. <i>Knee Surgery, Sports Traumatology, Arthroscopy</i> , 2018, 26, 3762-3769.	4.2	24
9	Return to sport testing at 6 months after arthroscopic shoulder stabilization reveals residual strength and functional deficits. <i>Journal of Shoulder and Elbow Surgery</i> , 2020, 29, S107-S114.	2.6	24
10	A Portable System for Remote Rehabilitation Following a Total Knee Replacement: A Pilot Randomized Controlled Clinical Study. <i>Sensors</i> , 2020, 20, 6118.	3.8	23
11	Effects of exercise therapy for the treatment of asymptomatic full-thickness supraspinatus tears on in vivo glenohumeral kinematics. <i>Journal of Shoulder and Elbow Surgery</i> , 2016, 25, 641-649.	2.6	22
12	Female sex is associated with greater rotatory knee laxity in collegiate athletes. <i>Knee Surgery, Sports Traumatology, Arthroscopy</i> , 2018, 26, 1319-1325.	4.2	22
13	Lateral femoral notch depth is not associated with increased rotatory instability in ACL-injured knees: a quantitative pivot shift analysis. <i>Knee Surgery, Sports Traumatology, Arthroscopy</i> , 2018, 26, 1399-1405.	4.2	21
14	Rehabilitation and Return to Sports: Proceedings of the International Consensus Meeting on Cartilage Repair of the Ankle. <i>Foot and Ankle International</i> , 2018, 39, 61S-67S.	2.3	21
15	Rotatory Knee Laxity Exists on a Continuum in Anterior Cruciate Ligament Injury. <i>Journal of Bone and Joint Surgery - Series A</i> , 2020, 102, 213-220.	3.0	19
16	Reliability and validity of a return to sports testing battery for the shoulder. <i>Physical Therapy in Sport</i> , 2021, 48, 1-11.	1.9	17
17	Prognostic Scoring System for Patients Undergoing Reconstructive Foot and Ankle Surgery for Charcot Neuroarthropathy: The Charcot Reconstruction Preoperative Prognostic Score. <i>Journal of Foot and Ankle Surgery</i> , 2018, 57, 451-455.	1.0	15
18	Sports Participation Is an Appropriate Expectation for Recreational Athletes Undergoing Shoulder Arthroplasty. <i>Orthopaedic Journal of Sports Medicine</i> , 2018, 6, 232596711880066.	1.7	15

#	ARTICLE	IF	CITATIONS
19	Criteria-based return-to-sport testing is associated with lower recurrence rates following arthroscopic Bankart repair. <i>Journal of Shoulder and Elbow Surgery</i> , 2021, 30, S14-S20.	2.6	15
20	Anatomic Anterior Cruciate Ligament Reconstruction Using Hamstring Tendons Restores Quantitative Pivot Shift. <i>Orthopaedic Journal of Sports Medicine</i> , 2018, 6, 232596711881236.	1.7	14
21	Return to preinjury sports after anterior cruciate ligament reconstruction is predicted by five independent factors. <i>Knee Surgery, Sports Traumatology, Arthroscopy</i> , 2022, 30, 84-92.	4.2	14
22	CRITERION-BASED REHABILITATION PROGRAM WITH RETURN TO SPORT TESTING FOLLOWING ACL RECONSTRUCTION: A CASE SERIES. <i>International Journal of Sports Physical Therapy</i> , 2020, 15, 1151-1173.	1.3	14
23	Improved Outcomes Following Arthroscopic Superior Capsular Reconstruction May Not Be Associated With Changes in Shoulder Kinematics: An In Vivo Study. <i>Arthroscopy - Journal of Arthroscopic and Related Surgery</i> , 2022, 38, 267-275.	2.7	12
24	Acute Response of the Infraspinatus and Biceps Tendons to Pitching in Youth Baseball. <i>Medicine and Science in Sports and Exercise</i> , 2017, 49, 1168-1175.	0.4	11
25	Rehabilitation and return to sports after anterior shoulder stabilization. <i>Annals of Joint</i> , 0, 2, 62-62.	1.0	10
26	Anterior cruciate ligament reconstruction with lateral extraarticular tenodesis better restores native knee kinematics in combined ACL and meniscal injury. <i>Knee Surgery, Sports Traumatology, Arthroscopy</i> , 2022, 30, 131-138.	4.2	10
27	Graft healing does not influence subjective outcomes and shoulder kinematics after superior capsule reconstruction: a prospective in vivo kinematic study. <i>Journal of Shoulder and Elbow Surgery</i> , 2021, 30, S48-S56.	2.6	9
28	Patient-Reported and Quantitative Outcomes of Anatomic Anterior Cruciate Ligament Reconstruction With Hamstring Tendon Autografts. <i>Orthopaedic Journal of Sports Medicine</i> , 2020, 8, 232596712092615.	1.7	8
29	Repair of high-grade partial thickness supraspinatus tears after surgical completion of the tear have a lower retear rate when compared to full-thickness tear repair. <i>Knee Surgery, Sports Traumatology, Arthroscopy</i> , 2021, 29, 2370-2375.	4.2	8
30	Task-Specific Training for Adults With Chronic Knee Pain: A Case Series. <i>Journal of Orthopaedic and Sports Physical Therapy</i> , 2017, 47, 548-556.	3.5	6
31	Failure rates and clinical outcomes after treatment for long-head biceps brachii tendon pathology: a comparison of three treatment types. <i>JSES International</i> , 2021, 5, 630-635.	1.6	5
32	In a small retrospective cohort of patients with syndesmotic injury, only athletes benefited from placement of a suture button device: a pilot study. <i>Journal of ISAKOS</i> , 2019, 4, 21-25.	2.3	4
33	Associations between range of motion, strength, tear size, patient-reported outcomes, and glenohumeral kinematics in individuals with symptomatic isolated supraspinatus tears. <i>Journal of Shoulder and Elbow Surgery</i> , 2022, 31, 1261-1271.	2.6	4
34	A comparison of 6-month outcomes between periacetabular osteotomy with concomitant hip arthroscopy to isolated hip arthroscopy for femoroacetabular impingement. <i>Archives of Orthopaedic and Trauma Surgery</i> , 2022, 142, 471-480.	2.4	3
35	Postoperative Rehabilitation and Return to Play Criteria After Anterior Shoulder Stabilization Procedures. <i>Operative Techniques in Orthopaedics</i> , 2021, 31, 100863.	0.1	1
36	ACL: Time-Based Return to Play. "Role of Patient Reporting in Return to Play", 2018, , 263-271.		0