

Augustine Rajakumar

List of Publications by Year in Descending Order

Source: <https://exaly.com/author-pdf/4777631/augustine-rajakumar-publications-by-year.pdf>

Version: 2024-04-23

This document has been generated based on the publications and citations recorded by exaly.com. For the latest version of this publication list, visit the link given above.

The third column is the impact factor (IF) of the journal, and the fourth column is the number of citations of the article.

40
papers

1,963
citations

20
h-index

44
g-index

44
ext. papers

2,253
ext. citations

5.2
avg, IF

4.21
L-index

#	Paper	IF	Citations
40	Reduced angiogenic and increased inflammatory profiles of cord blood cells in severe but not mild preeclampsia. <i>Scientific Reports</i> , 2021 , 11, 3630	4.9	0
39	Transcription factor AP2A affects sFLT1 expression and decidualization in decidual stromal cells: Implications to preeclampsia pathology. <i>Pregnancy Hypertension</i> , 2020 , 21, 152-158	2.6	2
38	Transcription factor ID1 is involved in decidualization of stromal cells: Implications in preeclampsia. <i>Pregnancy Hypertension</i> , 2020 , 21, 7-13	2.6	2
37	Alternatively Activated Macrophages Are the Primary Retinoic Acid-Producing Cells in Human Decidua. <i>Reproductive Sciences</i> , 2020 , 27, 334-341	3	2
36	Organotin(IV) derivatives of amide-based carboxylates: Synthesis, spectroscopic characterization, single crystal studies and antimicrobial, antioxidant, cytotoxic, anti-leishmanial, hemolytic, noncancerous, anticancer activities. <i>Inorganica Chimica Acta</i> , 2020 , 505, 119433	2.7	6
35	Aberrant retinoic acid production in the decidua: Implications for pre-eclampsia. <i>Journal of Obstetrics and Gynaecology Research</i> , 2020 , 46, 1007-1016	1.9	1
34	Human Endometrial Stromal Cell Differentiation is Stimulated by PPAR α Activation: New Targets for Infertility?. <i>Journal of Clinical Endocrinology and Metabolism</i> , 2020 , 105,	5.6	2
33	Potency Analysis of Mesenchymal Stromal Cells Using a Phospho-STAT Matrix Loop Analytical Approach. <i>Stem Cells</i> , 2019 , 37, 1119-1125	5.8	11
32	Retinoic Acid Is a Negative Regulator of sFLT1 Expression in Decidual Stromal Cells, and Its Levels Are Reduced in Preeclamptic Decidua. <i>Hypertension</i> , 2019 , 73, 1104-1111	8.5	13
31	Effects of Supraphysiologic Levels of Estradiol on Endometrial Decidualization, sFlt1, and HOXA10 Expression. <i>Reproductive Sciences</i> , 2019 , 26, 1626-1632	3	4
30	Decidual cells from women with preeclampsia exhibit inadequate decidualization and reduced sFlt1 suppression. <i>Pregnancy Hypertension</i> , 2019 , 15, 64-71	2.6	16
29	Associations Between the Features of Gross Placental Morphology and Birthweight. <i>Pediatric and Developmental Pathology</i> , 2019 , 22, 194-204	2.2	8
28	Associations Between Features of Placental Morphology and Birth Weight in Dichorionic Twins. <i>American Journal of Epidemiology</i> , 2019 , 188, 518-526	3.8	5
27	Villous explants from preeclamptic placentas induce sFlt1 in PBMCs: An ex vivo co-culture study. <i>Pregnancy Hypertension</i> , 2018 , 12, 40-46	2.6	6
26	Comparison of diameter-based and image-based measures of surface area from gross placental pathology for use in epidemiologic studies. <i>Placenta</i> , 2018 , 69, 82-85	3.4	0
25	Relationship between hypoxia and downstream pathogenic pathways in preeclampsia. <i>Hypertension in Pregnancy</i> , 2017 , 36, 145-150	2	29
24	Human endometrial stromal cell plasticity: Reversible sFlt1 expression negatively coincides with decidualization. <i>Hypertension in Pregnancy</i> , 2017 , 36, 204-211	2	14

23	FLT1 and transcriptome-wide polyadenylation site (PAS) analysis in preeclampsia. <i>Scientific Reports</i> , 2017 , 7, 12139	4.9	23
22	Induced Human Decidual NK-Like Cells Improve Utero-Placental Perfusion in Mice. <i>PLoS ONE</i> , 2016 , 11, e0164353	3.7	13
21	Soluble fms-like tyrosine kinase 1 promotes angiotensin II sensitivity in preeclampsia. <i>Journal of Clinical Investigation</i> , 2016 , 126, 2561-74	15.9	76
20	Zika Virus Infects Human Placental Macrophages. <i>Cell Host and Microbe</i> , 2016 , 20, 83-90	23.4	315
19	Gelsolin is an endogenous inhibitor of syncytiotrophoblast extracellular vesicle shedding in pregnancy. <i>Pregnancy Hypertension</i> , 2016 , 6, 333-339	2.6	7
18	Vaccine-Derived Neutralizing Antibodies to the Human Cytomegalovirus gH/gL Pentamer Potently Block Primary Cytotrophoblast Infection. <i>Journal of Virology</i> , 2015 , 89, 11884-98	6.6	62
17	Drug Repurposing Screen Identifies Foxo1-Dependent Angiopoietin-2 Regulation in Sepsis. <i>Critical Care Medicine</i> , 2015 , 43, e230-40	1.4	27
16	Excess placental secreted frizzled-related protein 1 in maternal smokers impairs fetal growth. <i>Journal of Clinical Investigation</i> , 2015 , 125, 4021-5	15.9	15
15	Vascular pool of releasable soluble VEGF receptor-1 (sFLT1) in women with previous preeclampsia and uncomplicated pregnancy. <i>Journal of Clinical Endocrinology and Metabolism</i> , 2014 , 99, 978-87	5.6	12
14	Ouabain inhibits placental sFlt1 production by repressing HSP27-dependent HIF-1 α pathway. <i>FASEB Journal</i> , 2014 , 28, 4324-34	0.9	37
13	Preeclampsia is associated with the presence of transcriptionally active placental fragments in the maternal lung. <i>Hypertension</i> , 2013 , 62, 608-13	8.5	32
12	Conversion of peripheral blood NK cells to a decidual NK-like phenotype by a cocktail of defined factors. <i>Journal of Immunology</i> , 2013 , 190, 3939-48	5.3	115
11	Transcriptionally active syncytial aggregates in the maternal circulation may contribute to circulating soluble fms-like tyrosine kinase 1 in preeclampsia. <i>Hypertension</i> , 2012 , 59, 256-64	8.5	127
10	Angiotensin II type 1 receptor antibodies and increased angiotensin II sensitivity in pregnant rats. <i>Hypertension</i> , 2011 , 58, 77-84	8.5	107
9	Preeclampsia and soluble fms-like tyrosine kinase 1. <i>Journal of Clinical Endocrinology and Metabolism</i> , 2009 , 94, 2252-4	5.6	31
8	Maternal endothelial progenitor colony-forming units with macrophage characteristics are reduced in preeclampsia. <i>American Journal of Hypertension</i> , 2009 , 22, 1014-9	2.3	31
7	Increased myeloperoxidase in the placenta and circulation of women with preeclampsia. <i>Hypertension</i> , 2008 , 52, 387-93	8.5	49
6	Agonistic angiotensin II type 1 receptor autoantibodies in postpartum women with a history of preeclampsia. <i>Hypertension</i> , 2007 , 49, 612-7	8.5	98

5	Soluble fms-like tyrosine kinase 1 is increased in preeclampsia but not in normotensive pregnancies with small-for-gestational-age neonates: relationship to circulating placental growth factor. <i>Journal of Clinical Endocrinology and Metabolism</i> , 2005 , 90, 4895-903	5.6	202
4	Trans-activators regulating neuronal glucose transporter isoform-3 gene expression in mammalian neurons. <i>Journal of Biological Chemistry</i> , 2004 , 279, 26768-79	5.4	38
3	Preeclampsia and future cardiovascular disease: potential role of altered angiogenesis and insulin resistance. <i>Journal of Clinical Endocrinology and Metabolism</i> , 2004 , 89, 6239-43	5.6	165
2	Selective overexpression of the hypoxia-inducible transcription factor, HIF-2alpha, in placentas from women with preeclampsia. <i>Biology of Reproduction</i> , 2001 , 64, 499-506	3.9	120
1	Expression, ontogeny, and regulation of hypoxia-inducible transcription factors in the human placenta. <i>Biology of Reproduction</i> , 2000 , 63, 559-69	3.9	138