

# Willem Evert Van Spil

## List of Publications by Year in descending order

Source: <https://exaly.com/author-pdf/4779952/publications.pdf>

Version: 2024-02-01

17  
papers

635  
citations

759233

12  
h-index

888059

17  
g-index

17  
all docs

17  
docs citations

17  
times ranked

838  
citing authors

#	ARTICLE	IF	CITATIONS
1	An automated workflow based on hip shape improves personalized risk prediction for hip osteoarthritis in the CHECK study. <i>Osteoarthritis and Cartilage</i> , 2020, 28, 62-70.	1.3	15
2	A consensus-based framework for conducting and reporting osteoarthritis phenotype research. <i>Arthritis Research and Therapy</i> , 2020, 22, 54.	3.5	28
3	Osteoarthritis phenotypes and novel therapeutic targets. <i>Biochemical Pharmacology</i> , 2019, 165, 41-48.	4.4	135
4	Bone texture analysis for prediction of incident radiographic hip osteoarthritis using machine learning: data from the Cohort Hip and Cohort Knee (CHECK) study. <i>Osteoarthritis and Cartilage</i> , 2019, 27, 906-914.	1.3	26
5	Trajectories of femorotibial cartilage thickness among persons with or at risk of knee osteoarthritis: development of a prediction model to identify progressors. <i>Osteoarthritis and Cartilage</i> , 2019, 27, 257-265.	1.3	16
6	Molecular taxonomy of osteoarthritis for patient stratification, disease management and drug development: biochemical markers associated with emerging clinical phenotypes and molecular endotypes. <i>Current Opinion in Rheumatology</i> , 2019, 31, 80-89.	4.3	76
7	Disease burden of knee osteoarthritis patients with a joint replacement compared to matched controls: a population-based analysis of a Dutch medical claims database. <i>Osteoarthritis and Cartilage</i> , 2018, 26, 202-210.	1.3	6
8	Efficacy of bisphosphonates in specific knee osteoarthritis subpopulations: protocol for an OA Trial Bank systematic review and individual patient data meta-analysis. <i>BMJ Open</i> , 2018, 8, e023889.	1.9	12
9	A sex-specific association between incident radiographic osteoarthritis of hip or knee and incident peripheral arterial calcifications: 8-year prospective data from Cohort Hip and Cohort Knee (CHECK). <i>Osteoarthritis and Cartilage</i> , 2017, 25, 1814-1821.	1.3	12
10	Six weeks of continuous joint distraction appears sufficient for clinical benefit and cartilaginous tissue repair in the treatment of knee osteoarthritis. <i>Knee</i> , 2016, 23, 785-791.	1.6	21
11	Associations of markers of matrix metabolism, inflammation markers, and adipokines with superior cam deformity of the hip and their relation with future hip osteoarthritis. <i>Osteoarthritis and Cartilage</i> , 2015, 23, 1897-1905.	1.3	5
12	The ability of systemic biochemical markers to reflect presence, incidence, and progression of early-stage radiographic knee and hip osteoarthritis: data from CHECK. <i>Osteoarthritis and Cartilage</i> , 2015, 23, 1388-1397.	1.3	39
13	Systemic biochemical markers of joint metabolism and inflammation in relation to radiographic parameters and pain of the knee: data from CHECK, a cohort of early-osteoarthritis subjects. <i>Osteoarthritis and Cartilage</i> , 2015, 23, 48-56.	1.3	32
14	Associations of CTX-II with biochemical markers of bone turnover raise questions on its tissue origin: data from CHECK, a cohort study of early osteoarthritis. <i>Annals of the Rheumatic Diseases</i> , 2013, 72, 29-36.	0.9	68
15	Cross-sectional and predictive associations between plasma adipokines and radiographic signs of early-stage knee osteoarthritis: data from CHECK. <i>Osteoarthritis and Cartilage</i> , 2012, 20, 1278-1285.	1.3	51
16	Clusters within a wide spectrum of biochemical markers for osteoarthritis: data from CHECK, a large cohort of individuals with very early symptomatic osteoarthritis. <i>Osteoarthritis and Cartilage</i> , 2012, 20, 745-754.	1.3	58
17	Lack of evidence of a beneficial effect of azathioprine in dogs treated with prednisolone for idiopathic immune-mediated hemolytic anemia: a retrospective cohort study. <i>BMC Veterinary Research</i> , 2011, 7, 15.	1.9	35