
List of Publications by Year in descending order

Source: https://exaly.com/author-pdf/478418/publications.pdf Version: 2024-02-01



Ι ΙΝΙΔΑ ΡΙΝΙΑΜΑΝ

#	Article	IF	CITATIONS
1	Stress & the gut-brain axis: Regulation by the microbiome. Neurobiology of Stress, 2017, 7, 124-136.	1.9	736
2	Ascending projections from the caudal visceral nucleus of the solitary tract to brain regions involved in food intake and energy expenditure. Brain Research, 2010, 1350, 18-34.	1.1	234
3	Characterization of the central nervous system innervation of the rat spleen using viral transneuronal tracing. Journal of Comparative Neurology, 2001, 439, 1-18.	0.9	203
4	Hindbrain noradrenergic A2 neurons: diverse roles in autonomic, endocrine, cognitive, and behavioral functions. American Journal of Physiology - Regulatory Integrative and Comparative Physiology, 2011, 300, R222-R235.	0.9	167
5	Distribution and neurochemical phenotypes of caudal medullary neurons activated to express cFos following peripheral administration of cholecystokinin. Journal of Comparative Neurology, 1993, 338, 475-490.	0.9	157
6	Cholecystokinin activates catecholaminergic neurons in the caudal medulla that innervate the paraventricular nucleus of the hypothalamus in rats. Journal of Comparative Neurology, 1995, 360, 246-256.	0.9	152
7	Interoceptive stress activates glucagon-like peptide-1 neurons that project to the hypothalamus. American Journal of Physiology - Regulatory Integrative and Comparative Physiology, 1999, 277, R582-R590.	0.9	150
8	Hindbrain Noradrenergic Lesions Attenuate Anorexia and Alter Central cFos Expression in Rats after Gastric Viscerosensory Stimulation. Journal of Neuroscience, 2003, 23, 10084-10092.	1.7	142
9	Stress exposure, food intake and emotional state. Stress, 2015, 18, 381-99.	0.8	128
10	Oxytocinergic inputs to the nucleus of the solitary tract and dorsal motor nucleus of the vagus in neonatal rats. Journal of Comparative Neurology, 1998, 399, 101-109.	0.9	123
11	Central and peripheral GLP-1 systems independently suppress eating. Nature Metabolism, 2021, 3, 258-273.	5.1	107
12	Oxytocin knockout mice demonstrate enhanced intake of sweet and nonsweet carbohydrate solutions. American Journal of Physiology - Regulatory Integrative and Comparative Physiology, 2007, 292, R1828-R1833.	0.9	106
13	Anterograde Transneuronal Viral Tracing of Central Viscerosensory Pathways in Rats. Journal of Neuroscience, 2004, 24, 2782-2786.	1.7	101
14	Progressive Postnatal Assembly of Limbic–Autonomic Circuits Revealed by Central Transneuronal Transport of Pseudorabies Virus. Journal of Neuroscience, 2000, 20, 2731-2741.	1.7	99
15	Enhanced initial and sustained intake of sucrose solution in mice with an oxytocin gene deletion. American Journal of Physiology - Regulatory Integrative and Comparative Physiology, 2005, 289, R1798-R1806.	0.9	99
16	Early Experience Modifies the Postnatal Assembly of Autonomic Emotional Motor Circuits in Rats. Journal of Neuroscience, 2005, 25, 9102-9111.	1.7	89
17	GLP-1 receptor signaling contributes to anorexigenic effect of centrally administered oxytocin in rats. American Journal of Physiology - Regulatory Integrative and Comparative Physiology, 2002, 283, R99-R106.	0.9	85
18	Central c-Fos expression in neonatal and adult rats after subcutaneous injection of hypertonic saline. Neuroscience, 1997, 79, 1165-1175.	1.1	83

#	Article	IF	CITATIONS
19	The Role of Central Glucagon-Like Peptide-1 in Mediating the Effects of Visceral Illness: Differential Effects in Rats and Mice. Endocrinology, 2005, 146, 458-462.	1.4	83
20	Visceral sensory inputs to the endocrine hypothalamus. Frontiers in Neuroendocrinology, 2007, 28, 50-60.	2.5	80
21	Establishment of vagal sensorimotor circuits during fetal development in rats. Journal of Neurobiology, 1993, 24, 641-659.	3.7	77
22	The organization of vagal innervation of rat pancreas using cholera toxin—horseradish peroxidase conjugate. Journal of the Autonomic Nervous System, 1987, 21, 109-125.	1.9	76
23	Persistence of fluoro-gold following degeneration of labeled motoneurons is due to phagocytosis by microglia and macrophages. Neuroscience, 1991, 44, 765-776.	1.1	76
24	Oxytocin gene deletion mice overconsume palatable sucrose solution but not palatable lipid emulsions. American Journal of Physiology - Regulatory Integrative and Comparative Physiology, 2007, 293, R1063-R1068.	0.9	76
25	Postnatal development of catecholamine inputs to the paraventricular nucleus of the hypothalamus in rats. Journal of Comparative Neurology, 2001, 438, 411-422.	0.9	75
26	Negative Energy Balance Blocks Neural and Behavioral Responses to Acute Stress by "Silencing" Central Glucagon-Like Peptide 1 Signaling in Rats. Journal of Neuroscience, 2015, 35, 10701-10714.	1.7	73
27	Thyrotropin-releasing hormone-immunoreactive nerve terminals synapse on the dendrites of gastric vagal motoneurons in the rat. Journal of Comparative Neurology, 1990, 294, 235-251.	0.9	72
28	A functional role for central glucagon-like peptide-1 receptors in lithium chloride-induced anorexia. American Journal of Physiology - Regulatory Integrative and Comparative Physiology, 1999, 277, R1537-R1540.	0.9	72
29	Noradrenergic Inputs to the Bed Nucleus of the Stria Terminalis and Paraventricular Nucleus of the Hypothalamus Underlie Hypothalamic-Pituitary-Adrenal Axis But Not Hypophagic or Conditioned Avoidance Responses to Systemic Yohimbine. Journal of Neuroscience, 2006, 26, 11442-11453.	1.7	66
30	Noradrenergic inputs to the paraventricular hypothalamus contribute to hypothalamic–pituitary–adrenal axis and central Fos activation in rats after acute systemic endotoxin exposure. Neuroscience, 2008, 156, 1093-1102.	1.1	65
31	Vagal Interoceptive Modulation of Motivated Behavior. Physiology, 2018, 33, 151-167.	1.6	65
32	Common and distinct neural inputs to the medial central nucleus of the amygdala and anterior ventrolateral bed nucleus of stria terminalis in rats. Brain Structure and Function, 2013, 218, 187-208.	1.2	64
33	Connections of Barrington's nucleus to the sympathetic nervous system in rats. Journal of the Autonomic Nervous System, 2000, 79, 117-128.	1.9	63
34	Glutamatergic phenotype of glucagon-like peptide 1 neurons in the caudal nucleus of the solitary tract in rats. Brain Structure and Function, 2015, 220, 3011-3022.	1.2	60
35	Ultrastructural localization of thyrotropin-releasing hormone immunoreactivity in the dorsal vagal complex in rat. Neuroscience Letters, 1989, 104, 7-12.	1.0	57
36	Identification of lingual motor control circuits using two strains of pseudorabies virus. Neuroscience, 2002, 115, 1139-1151.	1,1	56

#	Article	IF	CITATIONS
37	Enhanced corticosterone concentrations and attenuated Fos expression in the medial amygdala of female oxytocin knockout mice exposed to psychogenic stress. American Journal of Physiology - Regulatory Integrative and Comparative Physiology, 2004, 287, R1494-R1504.	0.9	56
38	Exogenous cholecystokinin activates cFos expression in medullary but not hypothalamic neurons in neonatal rats. Developmental Brain Research, 1994, 77, 140-145.	2.1	53
39	Overnight food deprivation markedly attenuates hindbrain noradrenergic, glucagon-like peptide-1, and hypothalamic neural responses to exogenous cholecystokinin in male rats. Physiology and Behavior, 2013, 121, 35-42.	1.0	53
40	Postnatal development of hypothalamic inputs to the dorsal vagal complex in rats. Physiology and Behavior, 2003, 79, 65-70.	1.0	51
41	Satiation and Stress-Induced Hypophagia: Examining the Role of Hindbrain Neurons Expressing Prolactin-Releasing Peptide or Glucagon-Like Peptide 1. Frontiers in Neuroscience, 2012, 6, 199.	1.4	51
42	Lesions of the C1 catecholaminergic neurons of the ventrolateral medulla in rats using anti-DβH-saporin. American Journal of Physiology - Regulatory Integrative and Comparative Physiology, 1999, 277, R1063-R1075.	0.9	50
43	Experimental dissociation of neural circuits underlying conditioned avoidance and hypophagic responses to lithium chloride. American Journal of Physiology - Regulatory Integrative and Comparative Physiology, 2007, 293, R1495-R1503.	0.9	49
44	Differential activation of chemically identified neurons in the caudal nucleus of the solitary tract in non-entrained rats after intake of satiating vs. non-satiating meals. Physiology and Behavior, 2014, 136, 47-54.	1.0	47
45	Central Fos expression and conditioned flavor avoidance in rats following intragastric administration of bitter taste receptor ligands. American Journal of Physiology - Regulatory Integrative and Comparative Physiology, 2009, 296, R528-R536.	0.9	45
46	Simplified CLARITY for visualizing immunofluorescence labeling in the developing rat brain. Brain Structure and Function, 2016, 221, 2375-2383.	1.2	44
47	Interoceptive modulation of neuroendocrine, emotional, and hypophagic responses to stress. Physiology and Behavior, 2017, 176, 195-206.	1.0	44
48	Dehydration anorexia is attenuated in oxytocin-deficient mice. American Journal of Physiology - Regulatory Integrative and Comparative Physiology, 2005, 288, R1791-R1799.	0.9	42
49	Retrograde transynaptic pseudorabies virus infection of central autonomic circuits in neonatal rats. Developmental Brain Research, 1999, 114, 207-216.	2.1	39
50	Cholecystokinin and D-fenfluramine inhibit food intake in oxytocin-deficient mice. American Journal of Physiology - Regulatory Integrative and Comparative Physiology, 2003, 285, R1037-R1045.	0.9	39
51	The anxiogenic drug yohimbine activates central viscerosensory circuits in rats. Journal of Comparative Neurology, 2005, 492, 426-441.	0.9	38
52	Burst activation of dopamine neurons produces prolonged post-burst availability of actively released dopamine. Neuropsychopharmacology, 2018, 43, 2083-2092.	2.8	36
53	Excitatory Hindbrain–Forebrain Communication Is Required for Cisplatin-Induced Anorexia and Weight Loss. Journal of Neuroscience, 2017, 37, 362-370.	1.7	35
54	Viscerosensory activation of noradrenergic inputs to the amygdala in rats. Physiology and Behavior, 2002, 77, 723-729.	1.0	34

#	Article	IF	CITATIONS
55	Trimethylthiazoline supports conditioned flavor avoidance and activates viscerosensory, hypothalamic, and limbic circuits in rats. American Journal of Physiology - Regulatory Integrative and Comparative Physiology, 2005, 288, R1716-R1726.	0.9	30
56	Early life experience shapes the functional organization of stress-responsive visceral circuits. Physiology and Behavior, 2011, 104, 632-640.	1.0	30
57	Synaptic Inputs to the Mouse Dorsal Vagal Complex and Its Resident Preproglucagon Neurons. Journal of Neuroscience, 2019, 39, 9767-9781.	1.7	30
58	Chronic Suppression of Glucagon-Like Peptide-1 Receptor (GLP1R) mRNA Translation in the Rat Bed Nucleus of the Stria Terminalis Reduces Anxiety-Like Behavior and Stress-Induced Hypophagia, But Prolongs Stress-Induced Elevation of Plasma Corticosterone. Journal of Neuroscience, 2019, 39, 2649-2663.	1.7	29
59	Distribution of glucagon-like peptide 1-immunopositive neurons in human caudal medulla. Brain Structure and Function, 2015, 220, 1213-1219.	1.2	28
60	Repeated brief postnatal maternal separation enhances hypothalamic gastric autonomic circuits in juvenile rats. Neuroscience, 2010, 165, 265-277.	1.1	27
61	Systemic leptin dose-dependently increases STAT3 phosphorylation within hypothalamic and hindbrain nuclei. American Journal of Physiology - Regulatory Integrative and Comparative Physiology, 2014, 306, R576-R585.	0.9	27
62	Delineation of vagal emetic pathways: intragastric copper sulfate-induced emesis and viral tract tracing in musk shrews. American Journal of Physiology - Regulatory Integrative and Comparative Physiology, 2014, 306, R341-R351.	0.9	27
63	GLPâ€l neurons form a local synaptic circuit within the rodent nucleus of the solitary tract. Journal of Comparative Neurology, 2018, 526, 2149-2164.	0.9	27
64	Ontogeny of hypothalamic-hindbrain feeding control circuits. Developmental Psychobiology, 2006, 48, 389-396.	0.9	24
65	Yohimbine anxiogenesis in the elevated plus maze requires hindbrain noradrenergic neurons that target the anterior ventrolateral bed nucleus of the stria terminalis. European Journal of Neuroscience, 2013, 37, 1340-1349.	1.2	23
66	Distribution of glucose transporter isoform-3 and hexokinase I in the postnatal murine brain1This work was presented in an abstract form at the Pediatric Academic Societies 1998, held in New Orleans, LA.1. Brain Research, 1999, 846, 260-264.	1.1	21
67	Antagonism of central glucagon-like peptide-1 receptors enhances lipopolysaccharide-induced fever. Autonomic Neuroscience: Basic and Clinical, 2000, 85, 98-101.	1.4	21
68	Central neural responses to restraint stress are altered in rats with an early life history of repeated brief maternal separation. Neuroscience, 2011, 192, 413-428.	1.1	21
69	Hindbrain glucagon-like peptide-1 neurons track intake volume and contribute to injection stress-induced hypophagia in meal-entrained rats. American Journal of Physiology - Regulatory Integrative and Comparative Physiology, 2016, 310, R906-R916.	0.9	21
70	Cholecystokinin induces Fos expression in catecholaminergic neurons of the macaque monkey caudal medulla. Brain Research, 1997, 770, 37-44.	1.1	20
71	The postnatal emergence of dehydration anorexia in rats is temporally associated with the emergence of dehydration-induced inhibition of gastric emptying. Physiology and Behavior, 1998, 64, 683-687.	1.0	19
72	Noradrenergic axon terminals contact gastric preautonomic neurons in the paraventricular nucleus of the hypothalamus in rats. Journal of Comparative Neurology, 2007, 501, 608-618.	0.9	19

#	Article	IF	CITATIONS
73	The role of nucleus of the solitary tract glucagonâ€like peptideâ€1 and prolactinâ€releasing peptide neurons in stress: anatomy, physiology and cellular interactions. British Journal of Pharmacology, 2022, 179, 642-658.	2.7	19
74	Yohimbine anxiogenesis in the elevated plus maze is disrupted by bilaterally disconnecting the bed nucleus of the stria terminalis from the central nucleus of the amygdala. Neuroscience, 2012, 223, 200-208.	1.1	18
75	Early experience alters limbic forebrain Fos responses to a stressful interoceptive stimulus in young adult rats. Physiology and Behavior, 2010, 100, 105-115.	1.0	16
76	Immune challenge activates neural inputs to the ventrolateral bed nucleus of the stria terminalis. Physiology and Behavior, 2011, 104, 257-265.	1.0	16
77	Organization of multisynaptic circuits within and between the medial and the central extended amygdala. Journal of Comparative Neurology, 2013, 521, 3406-3431.	0.9	14
78	Ectopic sympathetic preganglionic neurons maintain proper connectivity in the reeler mutant mouse. Neuroscience, 2003, 118, 439-450.	1.1	13
79	Progressive postnatal increases in Fos immunoreactivity in the forebrain and brain stem of rats after viscerosensory stimulation with lithium chloride. American Journal of Physiology - Regulatory Integrative and Comparative Physiology, 2007, 292, R1212-R1223.	0.9	13
80	A potential gastrointestinal link between enhanced postnatal maternal care and reduced anxiety-like behavior in adolescent rats Behavioral Neuroscience, 2009, 123, 1178-1184.	0.6	13
81	Maternal high-fat diet increases independent feeding in pre-weanling rat pups. Physiology and Behavior, 2016, 157, 237-245.	1.0	13
82	Ondansetron blocks LiCl-induced conditioned place avoidance but not conditioned taste/flavor avoidance in rats. Physiology and Behavior, 2009, 98, 381-385.	1.0	12
83	Ghrelin signaling contributes to fasting-induced attenuation of hindbrain neural activation and hypophagic responses to systemic cholecystokinin in rats. American Journal of Physiology - Regulatory Integrative and Comparative Physiology, 2020, 318, R1014-R1023.	0.9	11
84	Psychogenic Stress Activates C-Fos in Nucleus Accumbens-Projecting Neurons of the Hippocampal Ventral Subiculum. International Journal of Neuropsychopharmacology, 2017, 20, 855-860.	1.0	10
85	Postnatal Development of Central Feeding Circuits. , 2004, , 159-194.		9
86	Characterization of the neuroinvasive profile of a pseudorabies virus recombinant expressing the mTurquoise2 reporter in single and multiple injection experiments. Journal of Neuroscience Methods, 2018, 308, 228-239.	1.3	9
87	Hindbrain contributions to anorexia. American Journal of Physiology - Regulatory Integrative and Comparative Physiology, 2004, 287, R1035-R1036.	0.9	8
88	Cocaine self-administration and extinction alter medullary noradrenergic and limbic forebrain cFos responses to acute, noncontingent cocaine injections in adult rats. Neuroscience, 2014, 281, 241-250.	1.1	8
89	High Fat Diet Attenuates Cholecystokinin-Induced cFos Activation of Prolactin-Releasing Peptide-Expressing A2 Noradrenergic Neurons in the Caudal Nucleus of the Solitary Tract. Neuroscience, 2020, 447, 113-121.	1.1	8
90	Nicotine Enhances Footshock- and Lithium Chloride-Conditioned Place Avoidance in Male Rats. Nicotine and Tobacco Research, 2016, 18, 1920-1923.	1.4	7

#	ARTICLE	IF	CITATIONS
91	Access to gastric tissue promotes the survival of axotomized neurons in the dorsal motor nucleus of the vagus in neonatal rats. Journal of Comparative Neurology, 1991, 313, 213-226.	0.9	6
92	Amphetamine-induced activation of neurons within the rat nucleus of the solitary tract. Physiology and Behavior, 2019, 204, 355-363.	1.0	6
93	Sex and metabolic state interact to influence expression of passive avoidance memory in rats: Potential contribution of A2 noradrenergic neurons. Physiology and Behavior, 2021, 239, 113511.	1.0	5
94	Glucagon-like peptide 1 receptor-mediated stimulation of a GABAergic projection from the bed nucleus of the stria terminalis to the hypothalamic paraventricular nucleus. Neurobiology of Stress, 2021, 15, 100363.	1.9	5
95	New horizons for future research – Critical issues to consider for maximizing research excellence and impact. Molecular Metabolism, 2018, 14, 53-59.	3.0	3
96	Development of Central Visceral Circuits. , 2009, , .		2
97	Excitatory Hindbrain–Forebrain Communication Is Required for Cisplatin-Induced Anorexia and Weight Loss. Journal of Neuroscience, 2017, 37, 362-370.	1.7	1
98	Noradrenergic pathways contribute to hypothalamic activation in rats after LPS. FASEB Journal, 2007, 21, .	0.2	1
99	Organization and Postnatal Development of Visceral Sensory Inputs to the Neuroendocrine Hypothalamus. Masterclass in Neuroendocrinology, 2020, , 345-366.	0.1	1
100	M1659 Agonists of Bitter Taste Receptors Activate the Gut-Brain Axis and Influence Food Intake and Gastrointestinal Function in Mice. Gastroenterology, 2008, 134, A-392.	0.6	0
101	Characterization of Autonomic Emotional Motor Circuits in Young Rats. FASEB Journal, 2007, 21, A475.	0.2	0
102	Early life experience alters the functional assembly of viscerosensory neural circuits in the brainstem and forebrain of the rat. FASEB Journal, 2007, 21, A462.	0.2	0
103	Gastrointestinal (GI) infusion of bitter tastants supports conditioned flavor avoidance (CFA) and activates central neural Fos expression. FASEB Journal, 2008, 22, 1185.5.	0.2	0
104	Transneuronal viral tracing of sensory pathways from the stomach to the brain in the musk shrew, a small animal model for vomiting research. FASEB Journal, 2011, 25, 1075.11.	0.2	0