# RaphaÃ«l Rousset

List of Publications by Year
in descending order

Source: https://exaly.com/author-pdf/4788951/publications.pdf

Version: 2024-02-01

| | | | |
|---|---|---|---|
| 16 papers | 1,359 citations | 686830 13 h-index | 940134 16 g-index |
| 19 all docs | 19 docs citations | 19 times ranked | 1403 citing authors |

| # | Article | IF | Citations |
|---|---|---|---|
| 1 | Prickle Mediates Feedback Amplification to Generate Asymmetric Planar Cell Polarity Signaling. Cell, 2002, 109, 371-381. | 13.5 | 393 |
| 2 | The C-terminus of the HTLV-1 Tax oncoprotein mediates interaction with the PDZ domain of cellular proteins. Oncogene, 1998, 16, 643-654. | 2.6 | 177 |
| 3 | naked cuticle targets dishevelled to antagonize Wnt signal transduction. Genes and Development, 2001, 15, 658-671. | 2.7 | 146 |
| 4 | Effects on NF-Î°B1/p105 processing of the interaction between the HTLV-1 transactivator Tax and the proteasome. Nature, 1996, 381, 328-331. | 13.7 | 145 |
| 5 | Exclusion of Int-6 from PML Nuclear Bodies by Binding to the HTLV-I Tax Oncoprotein. Science, 1996, 273, 951-953. | 6.0 | 142 |
| 6 | Vertebrate Proteins Related to Drosophila Naked Cuticle Bind Dishevelled and Antagonize Wnt Signaling. Developmental Biology, 2001, 234, 93-106. | 0.9 | 123 |
| 7 | JNK signalling influences intracellular trafficking during Drosophila morphogenesis through regulation of the novel target gene Rab30. Developmental Biology, 2009, 331, 250-260. | 0.9 | 43 |
| 8 | JNK Signalling Controls Remodelling of the Segment Boundary through Cell Reprogramming during Drosophila Morphogenesis. PLoS Biology, 2010, 8, e1000390. | 2.6 | 38 |
| 9 | The Drosophila serine protease homologue Scarface regulates JNK signalling in a negative-feedback loop during epithelial morphogenesis. Development (Cambridge), 2010, 137, 2177-2186. | 1.2 | 35 |
| 10 | Zinc-dependent Interaction between Dishevelled and the Drosophila Wnt Antagonist Naked Cuticle. Journal of Biological Chemistry, 2002, 277, 49019-49026. | 1.6 | 32 |
| 11 | An Unconventional Nuclear Localization Motif Is Crucial for Function of the Drosophila Wnt/Wingless Antagonist Naked Cuticle. Genetics, 2006, 174, 331-348. | 1.2 | 21 |
| 12 | Tissue Adaptation to Environmental Cues by Symmetric and Asymmetric Division Modes of Intestinal Stem Cells. International Journal of Molecular Sciences, 2020, 21, 6362. | 1.8 | 21 |
| 13 | Drosophila Naked cuticle (Nkd) engages the nuclear import adaptor Importin-Î±3 to antagonize Wnt/Î²-catenin signaling. Developmental Biology, 2008, 318, 17-28. | 0.9 | 19 |
| 14 | Signalling crosstalk at the leading edge controls tissue closure dynamics in the Drosophila embryo. PLoS Genetics, 2017, 13, e1006640. | 1.5 | 10 |
| 15 | Polycomb and Hox Genes Control JNK-Induced Remodeling of the Segment Boundary during Drosophila Morphogenesis. Cell Reports, 2017, 19, 60-71. | 2.9 | 8 |
| 16 | Drosophila Morphogenesis: The Newtonian Revolution. Current Biology, 2003, 13, R494-R495. | 1.8 | 6 |