

Keith Waters

List of Publications by Citations

Source: <https://exaly.com/author-pdf/4804161/keith-waters-publications-by-citations.pdf>

Version: 2024-04-24

This document has been generated based on the publications and citations recorded by exaly.com. For the latest version of this publication list, visit the link given above.

The third column is the impact factor (IF) of the journal, and the fourth column is the number of citations of the article.

50
papers

2,253
citations

26
h-index

47
g-index

53
ext. papers

2,745
ext. citations

6.9
avg, IF

4.66
L-index

#	Paper	IF	Citations
50	Premature death after self-harm: a multicentre cohort study. <i>Lancet, The</i> , 2012 , 380, 1568-74	4.0	194
49	Suicide following self-harm: findings from the Multicentre Study of self-harm in England, 2000-2012. <i>Journal of Affective Disorders</i> , 2015 , 175, 147-51	6.6	175
48	UK legislation on analgesic packs: before and after study of long term effect on poisonings. <i>BMJ, The</i> , 2004 , 329, 1076	5.9	166
47	Repetition of self-harm and suicide following self-harm in children and adolescents: findings from the Multicentre Study of Self-harm in England. <i>Journal of Child Psychology and Psychiatry and Allied Disciplines</i> , 2012 , 53, 1212-9	7.9	164
46	Toxicity of antidepressants: rates of suicide relative to prescribing and non-fatal overdose. <i>British Journal of Psychiatry</i> , 2010 , 196, 354-8	5.4	127
45	Epidemiology and trends in non-fatal self-harm in three centres in England: 2000-2007. <i>British Journal of Psychiatry</i> , 2010 , 197, 493-8	5.4	121
44	Epidemiology and trends in non-fatal self-harm in three centres in England, 2000-2012: findings from the Multicentre Study of Self-harm in England. <i>BMJ Open</i> , 2016 , 6, e010538	3	120
43	Incidence of suicide, hospital-presenting non-fatal self-harm, and community-occurring non-fatal self-harm in adolescents in England (the iceberg model of self-harm): a retrospective study. <i>Lancet Psychiatry, the</i> , 2018 , 5, 167-174	23.3	107
42	How do methods of non-fatal self-harm relate to eventual suicide?. <i>Journal of Affective Disorders</i> , 2012 , 136, 526-33	6.6	84
41	Does clinical management improve outcomes following self-harm? Results from the multicentre study of self-harm in England. <i>PLoS ONE</i> , 2013 , 8, e70434	3.7	80
40	Risk factors for repetition and suicide following self-harm in older adults: multicentre cohort study. <i>British Journal of Psychiatry</i> , 2012 , 200, 399-404	5.4	71
39	Epidemiology and nature of self-harm in children and adolescents: findings from the multicentre study of self-harm in England. <i>European Child and Adolescent Psychiatry</i> , 2012 , 21, 369-77	5.5	64
38	Deliberate self-harm and antidepressant drugs. Investigation of a possible link. <i>British Journal of Psychiatry</i> , 2000 , 177, 551-6	5.4	63
37	Psychosocial assessment and repetition of self-harm: the significance of single and multiple repeat episode analyses. <i>Journal of Affective Disorders</i> , 2010 , 127, 257-65	6.6	62
36	Ethnic differences in self-harm, rates, characteristics and service provision: three-city cohort study. <i>British Journal of Psychiatry</i> , 2010 , 197, 212-8	5.4	58
35	The development of a population-level clinical screening tool for self-harm repetition and suicide: the ReACT Self-Harm Rule. <i>Psychological Medicine</i> , 2012 , 42, 2383-94	6.9	53
34	Suicide following presentation to hospital for non-fatal self-harm in the Multicentre Study of Self-harm: a long-term follow-up study. <i>Lancet Psychiatry, the</i> , 2019 , 6, 1021-1030	23.3	49

33	Rates of self-harm presenting to general hospitals: a comparison of data from the Multicentre Study of Self-Harm in England and Hospital Episode Statistics. <i>BMJ Open</i> , 2016 , 6, e009749	3	48
32	Hospital management of suicidal behaviour and subsequent mortality: a prospective cohort study. <i>Lancet Psychiatry</i> , 2015 , 2, 809-16	23.3	39
31	Alcohol use and misuse, self-harm and subsequent mortality: an epidemiological and longitudinal study from the multicentre study of self-harm in England. <i>Emergency Medicine Journal</i> , 2015 , 32, 793-9	1.5	39
30	Shared characteristics of suicides and other unnatural deaths following non-fatal self-harm? A multicentre study of risk factors. <i>Psychological Medicine</i> , 2012 , 42, 727-41	6.9	39
29	Mortality in children and adolescents following presentation to hospital after non-fatal self-harm in the Multicentre Study of Self-harm: a prospective observational cohort study. <i>The Lancet Child and Adolescent Health</i> , 2020 , 4, 111-120	14.5	31
28	Switching methods of self-harm at repeat episodes: Findings from a multicentre cohort study. <i>Journal of Affective Disorders</i> , 2015 , 180, 44-51	6.6	29
27	Impact of the recent recession on self-harm: Longitudinal ecological and patient-level investigation from the Multicentre Study of Self-harm in England. <i>Journal of Affective Disorders</i> , 2016 , 191, 132-8	6.6	29
26	Trends in prescribing and self-poisoning in relation to UK regulatory authority warnings against use of SSRI antidepressants in under-18-year-olds. <i>British Journal of Clinical Pharmacology</i> , 2009 , 68, 618-29	3.8	29
25	Impact of different pack sizes of paracetamol in the United Kingdom and Ireland on intentional overdoses: a comparative study. <i>BMC Public Health</i> , 2011 , 11, 460	4.1	27
24	Self-harm during the early period of the COVID-19 pandemic in England: Comparative trend analysis of hospital presentations. <i>Journal of Affective Disorders</i> , 2021 , 282, 991-995	6.6	24
23	Self-harm and life problems: findings from the Multicentre Study of Self-harm in England. <i>Social Psychiatry and Psychiatric Epidemiology</i> , 2016 , 51, 183-92	4.5	18
22	Impact of withdrawal of the analgesic Co-proxamol on nonfatal self-poisoning in the UK. <i>Crisis</i> , 2011 , 32, 81-7	2.8	16
21	Relative toxicity of benzodiazepines and hypnotics commonly used for self-poisoning: An epidemiological study of fatal toxicity and case fatality. <i>Journal of Psychopharmacology</i> , 2018 , 32, 654-662	4.6	15
20	Risk factors associated with repetition of self-harm in black and minority ethnic (BME) groups: a multi-centre cohort study. <i>Journal of Affective Disorders</i> , 2013 , 148, 435-9	6.6	13
19	Evaluation of a resource for people bereaved by suicide. <i>Crisis</i> , 2012 , 33, 254-64	2.8	13
18	Incidence and general hospital costs of self-harm across England: estimates based on the multicentre study of self-harm. <i>Epidemiology and Psychiatric Sciences</i> , 2020 , 29, e108	5.1	10
17	A comparative study of non-fatal self-poisoning with antidepressants relative to prescribing in three centres in England. <i>Journal of Affective Disorders</i> , 2010 , 123, 95-101	6.6	9
16	Paracetamol self-poisoning: Epidemiological study of trends and patient characteristics from the multicentre study of self-harm in England. <i>Journal of Affective Disorders</i> , 2020 , 276, 699-706	6.6	9

15	Self-harm and the COVID-19 pandemic: A study of factors contributing to self-harm during lockdown restrictions. <i>Journal of Psychiatric Research</i> , 2021 , 137, 437-443	5.2	9
14	Suicide and all-cause mortality following routine hospital management of self-harm: Propensity score analysis using multicentre cohort data. <i>PLoS ONE</i> , 2018 , 13, e0204670	3.7	9
13	Self-harm in midlife: analysis using data from the Multicentre Study of Self-harm in England. <i>British Journal of Psychiatry</i> , 2019 , 1-8	5.4	8
12	Variation by ethnic group in premature mortality risk following self-harm: a multicentre cohort study in England. <i>BMC Psychiatry</i> , 2015 , 15, 254	4.2	7
11	Alcohol-related mortality following self-harm: a multicentre cohort study. <i>JRSM Open</i> , 2014 , 5, 2054270414533326	4.5	7
10	High-Volume Repeaters of Self-Harm. <i>Crisis</i> , 2016 , 37, 427-437	2.8	6
9	Trends and characteristics of deliberate self-harm hospital presentations in an English County. <i>International Journal of Psychiatry in Clinical Practice</i> , 2006 , 10, 59-63	2.4	4
8	A Descriptive Study of Feelings of Arrested Escape (Entrapment) and Arrested Anger in People Presenting to an Emergency Department Following an Episode of Self-Harm. <i>Frontiers in Psychiatry</i> , 2016 , 7, 155	5	4
7	Self-harm during the early period of the COVID-19 Pandemic in England: comparative trend analysis of hospital presentations		1
6	Self-harm and the COVID-19 pandemic: a study of factors contributing to self-harm during lockdown restrictions		1
5	Risk of suicide in patients who present to hospital after self-cutting according to site of injury: findings from the Multicentre Study of Self-harm in England. <i>Psychological Medicine</i> , 2021 , 1-9	6.9	1
4	Self-harm in children and adolescents by ethnic group: an observational cohort study from the Multicentre Study of Self-Harm in England. <i>The Lancet Child and Adolescent Health</i> , 2021 , 5, 782-791	14.5	1
3	Cost-effectiveness of psychosocial assessment for individuals who present to hospital following self-harm in England: a model-based retrospective analysis.. <i>European Psychiatry</i> , 2022 , 1-32	6	0
2	Self-harm in children 12 years and younger: characteristics and outcomes based on the Multicentre Study of Self-harm in England. <i>Social Psychiatry and Psychiatric Epidemiology</i> , 2021 , 1	4.5	0
1	Self-harm in people experiencing homelessness: investigation of incidence, characteristics and outcomes using data from the Multicentre Study of Self-Harm in England.. <i>BJPsych Open</i> , 2022 , 8, e74	5	0