Onur Kaypakli

List of Publications by Year in descending order

Source: https://exaly.com/author-pdf/4820607/publications.pdf

Version: 2024-02-01

758635 839053 40 377 12 18 h-index citations g-index papers 42 42 42 632 all docs docs citations times ranked citing authors

#	Article	IF	CITATIONS
1	Oxidative Stress and Paraoxonase 1 Activity Predict Contrast-Induced Nephropathy in Patients With ST-Segment Elevation Myocardial Infarction Undergoing Primary Percutaneous Coronary Intervention. Angiology, 2015, 66, 339-345.	0.8	37
2	Fibrosis Marker Soluble ST2 Predicts Atrial Fibrillation Recurrence after Cryoballoon Catheter Ablation of Nonvalvular Paroxysmal Atrial Fibrillation. Korean Circulation Journal, 2018, 48, 920.	0.7	35
3	Mean platelet volume is independently associated with renal dysfunction in stable coronary artery disease. Platelets, 2014, 25, 274-278.	1.1	29
4	Paraoxonase-1 activity and oxidative stress in patients with anterior ST elevation myocardial infarction undergoing primary percutaneous coronary intervention with and without no-reflow. Atherosclerosis, 2014, 234, 415-420.	0.4	27
5	High-sensitivity cardiac troponin T levels in newly diagnosed hypertensive patients with different left ventricle geometry. Blood Pressure, 2014, 23, 240-247.	0.7	26
6	Decreased left atrial global longitudinal strain predicts the risk of atrial fibrillation recurrence after cryoablation in paroxysmal atrial fibrillation. Journal of Interventional Cardiac Electrophysiology, 2020, 58, 51-59.	0.6	21
7	Sâ \in R difference in V1â \in V2 is a novel criterion for differentiating the left from right ventricular outflow tract arrhythmias. Annals of Noninvasive Electrocardiology, 2018, 23, e12516.	0.5	17
8	Gamma glutamyl transferase activity is Independently associated with oxidative stress rather than SYNTAX score. Scandinavian Journal of Clinical and Laboratory Investigation, 2015, 75, 7-12.	0.6	16
9	Lower serum 25-hydroxyvitamin D level is associated with impaired myocardial performance and left ventricle hypertrophy in newly diagnosed hypertensive patients. Anatolian Journal of Cardiology, 2015, 15, 744-750.	0.5	16
10	Serum Bilirubin Level and Aortic Intima–Media Thickness in Patients Without Clinical Manifestation of Atherosclerotic Cardiovascular Disease. Angiology, 2014, 65, 308-312.	0.8	15
11	Relationship between platelet indices and international normalized ratio in patients with non-valvular atrial fibrillation. Platelets, 2014, 25, 311-316.	1.1	15
12	Association of P wave duration index with atrial fibrillation recurrence after cryoballoon catheter ablation. Journal of Electrocardiology, 2018, 51, 182-187.	0.4	15
13	Association between high-sensitivity troponin T, left ventricular hypertrophy, and myocardial performance index. Herz, 2015, 40, 1004-1010.	0.4	13
14	Highâ€Sensitivity Cardiac Troponin T Predicts Nondipper Hypertension in Newly Diagnosed Hypertensive Patients. Journal of Clinical Hypertension, 2013, 15, 731-736.	1.0	11
15	Increased Aortic Stiffness Predicts Contrast-Induced Nephropathy in Patients With Stable Coronary Artery Disease Undergoing Percutaneous Coronary Intervention. Angiology, 2014, 65, 806-811.	0.8	10
16	Serum 25â€Hydroxyvitamin D Level and Aortic Intimaâ€Media Thickness in Patients Without Clinical Manifestation of Atherosclerotic Cardiovascular Disease. Journal of Clinical Laboratory Analysis, 2015, 29, 305-311.	0.9	10
17	High morning blood pressure surge is associated with oxidative stress and paraoxonase 1 activity in newly diagnosed hypertensive patients. Clinical and Experimental Hypertension, 2016, 38, 680-685.	0.5	10
18	Paraoxonase (PON1) activity in patients with subclinical thoracic aortic atherosclerosis. International Journal of Cardiovascular Imaging, 2014, 30, 889-895.	0.7	8

#	Article	IF	CITATIONS
19	Relationship between mean platelet volume and morning blood pressure surge in newly diagnosed hypertensive patients. Anatolian Journal of Cardiology, 2015, 15, 107-112.	0.4	8
20	Coronary sinus lead delay index for optimization of coronary sinus lead placement. Annals of Noninvasive Electrocardiology, 2018, 23, .	0.5	4
21	Morning blood pressure surge is associated with the severity of stable coronary artery disease in hypertensive patients. Clinical and Experimental Hypertension, 2021, 43, 334-340.	0.5	4
22	Increased Rates of Coronary Artery Calcium Score in Patients with Non - Functioning Adrenal Incidentaloma. Endocrine, Metabolic and Immune Disorders - Drug Targets, 2021, 21, 1319-1325.	0.6	4
23	Is cystatin C an evaluative marker for right heart functions in systemic sclerosis?. International Journal of Cardiology, 2016, 221, 478-483.	0.8	3
24	High Left Ventricular Lead Sensing Delay Predicts QRS Narrowing and Good Response to Cardiac Resynchronization Therapy. PACE - Pacing and Clinical Electrophysiology, 2016, 39, 1317-1326.	0.5	3
25	Silent atrial fibrillation is associated with P-wave duration index in patients with cardiac resynchronisation therapy. Archives of Medical Sciences Atherosclerotic Diseases, 2019, 4, 74-81.	0.5	3
26	Impaired kidney function is associated with SYNTAX score in patients with stable coronary artery disease. Turk Kardiyoloji Dernegi Arsivi, 2014, 42, 621-628.	0.6	3
27	Estimated glomerular filtration rate is associated with both arterial stiffness and N-terminal pro-brain natriuretic peptide in newly diagnosed hypertensive patients. Clinical and Experimental Hypertension, 2014, 36, 374-379.	0.5	2
28	The Possible Effects of Dapagliflozin on 12-derived Electrocardiogram in Patients with Type 2 Diabetes Mellitus. Endocrine, Metabolic and Immune Disorders - Drug Targets, 2019, 19, 207-213.	0.6	2
29	Prevalence of cardiac arrhythmia and risk factors in chronic kidney disease patients. Saudi Journal of Kidney Diseases and Transplantation: an Official Publication of the Saudi Center for Organ Transplantation, Saudi Arabia, 2018, 29, 567.	0.4	2
30	Gamma Glutamyl Transferase Activity is Associated With Both Paraoxonase Activity and Aortic Stiffness in Hypertensive Patients. Journal of Clinical Laboratory Analysis, 2015, 29, 390-396.	0.9	1
31	Coronary Flow Velocity Reserve in Patients with Ascending Aorta Aneurysm. Echocardiography, 2015, 32, 975-982.	0.3	1
32	Increased hs-CRP and decreased 1,25-dihydroxyvitamin D are associated with increased left ventricle lead threshold. Journal of Interventional Cardiac Electrophysiology, 2016, 47, 177-183.	0.6	1
33	Subclinical atrial fibrillation frequency and associated parameters in patients with cardiac resynchronization therapy. Journal of Interventional Cardiac Electrophysiology, 2018, 52, 217-223.	0.6	1
34	Morning blood pressure surge is associated independently with orthostatic hypotension in hypertensive patients under treatment. Blood Pressure Monitoring, 2018, 23, 191-197.	0.4	1
35	Delta wave notching time is associated with accessory pathway localization in patients with Wolff-Parkinson-White syndrome. Journal of Interventional Cardiac Electrophysiology, 2018, 53, 73-79.	0.6	1
36	The effect of heart rate and pulse pressure on mean arterial pressure: the combined formula for calculation of mean arterial pressure. Blood Pressure Monitoring, 2021, 26, 373-379.	0.4	1

3

#	Article	IF	CITATIONS
37	NT-proBNP predicts impaired myocardial function in newly diagnosed hypertensive patients with preserved ejection fraction. Clinical and Experimental Hypertension, 2014, 36, 289-294.	0.5	0
38	A new use for maximum deflection index: Detection of intraventricular dyssynchrony. Journal of Electrocardiology, 2017, 50, 301-306.	0.4	0
39	Calculation of the Correct Angle of Bifurcation Predicts the Atherosclerotic Lesion Severity More Accurately. Journal of the Saudi Heart Association, 2021, 33, 147-148.	0.2	0
40	V1rÂ+ÂDIIq is a novel and accurate criterion to predict right vs. left paraseptal accessory pathways. Journal of Electrocardiology, 2022, 70, 30-34.	0.4	0