Onur Kaypakli

List of Publications by Year in descending order

Source: https://exaly.com/author-pdf/4820607/publications.pdf

Version: 2024-02-01

758635 839053 40 377 12 18 h-index citations g-index papers 42 42 42 632 all docs docs citations times ranked citing authors

#	Article	IF	Citations
1	V1rÂ+ÂDIIq is a novel and accurate criterion to predict right vs. left paraseptal accessory pathways. Journal of Electrocardiology, 2022, 70, 30-34.	0.4	O
2	Morning blood pressure surge is associated with the severity of stable coronary artery disease in hypertensive patients. Clinical and Experimental Hypertension, 2021, 43, 334-340.	0.5	4
3	Calculation of the Correct Angle of Bifurcation Predicts the Atherosclerotic Lesion Severity More Accurately. Journal of the Saudi Heart Association, 2021, 33, 147-148.	0.2	O
4	The effect of heart rate and pulse pressure on mean arterial pressure: the combined formula for calculation of mean arterial pressure. Blood Pressure Monitoring, 2021, 26, 373-379.	0.4	1
5	Increased Rates of Coronary Artery Calcium Score in Patients with Non - Functioning Adrenal Incidentaloma. Endocrine, Metabolic and Immune Disorders - Drug Targets, 2021, 21, 1319-1325.	0.6	4
6	Decreased left atrial global longitudinal strain predicts the risk of atrial fibrillation recurrence after cryoablation in paroxysmal atrial fibrillation. Journal of Interventional Cardiac Electrophysiology, 2020, 58, 51-59.	0.6	21
7	Silent atrial fibrillation is associated with P-wave duration index in patients with cardiac resynchronisation therapy. Archives of Medical Sciences Atherosclerotic Diseases, 2019, 4, 74-81.	0.5	3
8	The Possible Effects of Dapagliflozin on 12-derived Electrocardiogram in Patients with Type 2 Diabetes Mellitus. Endocrine, Metabolic and Immune Disorders - Drug Targets, 2019, 19, 207-213.	0.6	2
9	Coronary sinus lead delay index for optimization of coronary sinus lead placement. Annals of Noninvasive Electrocardiology, 2018, 23, .	0.5	4
10	Association of P wave duration index with atrial fibrillation recurrence after cryoballoon catheter ablation. Journal of Electrocardiology, 2018, 51, 182-187.	0.4	15
11	Sâ€R difference in V1â€V2 is a novel criterion for differentiating the left from right ventricular outflow tract arrhythmias. Annals of Noninvasive Electrocardiology, 2018, 23, e12516.	0.5	17
12	Fibrosis Marker Soluble ST2 Predicts Atrial Fibrillation Recurrence after Cryoballoon Catheter Ablation of Nonvalvular Paroxysmal Atrial Fibrillation. Korean Circulation Journal, 2018, 48, 920.	0.7	35
13	Subclinical atrial fibrillation frequency and associated parameters in patients with cardiac resynchronization therapy. Journal of Interventional Cardiac Electrophysiology, 2018, 52, 217-223.	0.6	1
14	Morning blood pressure surge is associated independently with orthostatic hypotension in hypertensive patients under treatment. Blood Pressure Monitoring, 2018, 23, 191-197.	0.4	1
15	Delta wave notching time is associated with accessory pathway localization in patients with Wolff-Parkinson-White syndrome. Journal of Interventional Cardiac Electrophysiology, 2018, 53, 73-79.	0.6	1
16	Prevalence of cardiac arrhythmia and risk factors in chronic kidney disease patients. Saudi Journal of Kidney Diseases and Transplantation: an Official Publication of the Saudi Center for Organ Transplantation, Saudi Arabia, 2018, 29, 567.	0.4	2
17	A new use for maximum deflection index: Detection of intraventricular dyssynchrony. Journal of Electrocardiology, 2017, 50, 301-306.	0.4	O
18	Increased hs-CRP and decreased 1,25-dihydroxyvitamin D are associated with increased left ventricle lead threshold. Journal of Interventional Cardiac Electrophysiology, 2016, 47, 177-183.	0.6	1

#	Article	IF	Citations
19	High morning blood pressure surge is associated with oxidative stress and paraoxonase 1 activity in newly diagnosed hypertensive patients. Clinical and Experimental Hypertension, 2016, 38, 680-685.	0.5	10
20	Is cystatin C an evaluative marker for right heart functions in systemic sclerosis?. International Journal of Cardiology, 2016, 221, 478-483.	0.8	3
21	High Left Ventricular Lead Sensing Delay Predicts QRS Narrowing and Good Response to Cardiac Resynchronization Therapy. PACE - Pacing and Clinical Electrophysiology, 2016, 39, 1317-1326.	0.5	3
22	Gamma Glutamyl Transferase Activity is Associated With Both Paraoxonase Activity and Aortic Stiffness in Hypertensive Patients. Journal of Clinical Laboratory Analysis, 2015, 29, 390-396.	0.9	1
23	Relationship between mean platelet volume and morning blood pressure surge in newly diagnosed hypertensive patients. Anatolian Journal of Cardiology, 2015, 15, 107-112.	0.4	8
24	Coronary Flow Velocity Reserve in Patients with Ascending Aorta Aneurysm. Echocardiography, 2015, 32, 975-982.	0.3	1
25	Association between high-sensitivity troponin T, left ventricular hypertrophy, and myocardial performance index. Herz, 2015, 40, 1004-1010.	0.4	13
26	Gamma glutamyl transferase activity is Independently associated with oxidative stress rather than SYNTAX score. Scandinavian Journal of Clinical and Laboratory Investigation, 2015, 75, 7-12.	0.6	16
27	Oxidative Stress and Paraoxonase 1 Activity Predict Contrast-Induced Nephropathy in Patients With ST-Segment Elevation Myocardial Infarction Undergoing Primary Percutaneous Coronary Intervention. Angiology, 2015, 66, 339-345.	0.8	37
28	Serum 25â€Hydroxyvitamin D Level and Aortic Intimaâ€Media Thickness in Patients Without Clinical Manifestation of Atherosclerotic Cardiovascular Disease. Journal of Clinical Laboratory Analysis, 2015, 29, 305-311.	0.9	10
29	Lower serum 25-hydroxyvitamin D level is associated with impaired myocardial performance and left ventricle hypertrophy in newly diagnosed hypertensive patients. Anatolian Journal of Cardiology, 2015, 15, 744-750.	0.5	16
30	NT-proBNP predicts impaired myocardial function in newly diagnosed hypertensive patients with preserved ejection fraction. Clinical and Experimental Hypertension, 2014, 36, 289-294.	0.5	0
31	Estimated glomerular filtration rate is associated with both arterial stiffness and N-terminal pro-brain natriuretic peptide in newly diagnosed hypertensive patients. Clinical and Experimental Hypertension, 2014, 36, 374-379.	0.5	2
32	Mean platelet volume is independently associated with renal dysfunction in stable coronary artery disease. Platelets, 2014, 25, 274-278.	1.1	29
33	Increased Aortic Stiffness Predicts Contrast-Induced Nephropathy in Patients With Stable Coronary Artery Disease Undergoing Percutaneous Coronary Intervention. Angiology, 2014, 65, 806-811.	0.8	10
34	Serum Bilirubin Level and Aortic Intima–Media Thickness in Patients Without Clinical Manifestation of Atherosclerotic Cardiovascular Disease. Angiology, 2014, 65, 308-312.	0.8	15
35	Relationship between platelet indices and international normalized ratio in patients with non-valvular atrial fibrillation. Platelets, 2014, 25, 311-316.	1.1	15
36	Paraoxonase (PON1) activity in patients with subclinical thoracic aortic atherosclerosis. International Journal of Cardiovascular Imaging, 2014, 30, 889-895.	0.7	8

#	Article	IF	CITATION
37	High-sensitivity cardiac troponin T levels in newly diagnosed hypertensive patients with different left ventricle geometry. Blood Pressure, 2014, 23, 240-247.	0.7	26
38	Paraoxonase-1 activity and oxidative stress in patients with anterior ST elevation myocardial infarction undergoing primary percutaneous coronary intervention with and without no-reflow. Atherosclerosis, 2014, 234, 415-420.	0.4	27
39	Impaired kidney function is associated with SYNTAX score in patients with stable coronary artery disease. Turk Kardiyoloji Dernegi Arsivi, 2014, 42, 621-628.	0.6	3
40	Highâ€Sensitivity Cardiac Troponin T Predicts Nondipper Hypertension in Newly Diagnosed Hypertensive Patients. Journal of Clinical Hypertension, 2013, 15, 731-736.	1.0	11