Laetitia Koppe

List of Publications by Year in Descending Order

Source: https://exaly.com/author-pdf/4825928/laetitia-koppe-publications-by-year.pdf

Version: 2024-04-20

This document has been generated based on the publications and citations recorded by exaly.com. For the latest version of this publication list, visit the link given above.

The third column is the impact factor (IF) of the journal, and the fourth column is the number of citations of the article.

1,194 19 34 g-index

53 1,653 5.7 4.94 ext. papers ext. citations avg, IF L-index

#	Paper	IF	Citations
43	Therapeutic strategies to limit tryptophan metabolites toxicity during chronic kidney disease 2022 , 28	1-295	
42	Probiotic Intake and Inflammation in Patients With Chronic Kidney Disease: An Analysis of the CKD-REIN Cohort <i>Frontiers in Nutrition</i> , 2022 , 9, 772596	6.2	1
41	A prospective observational study for justification, safety, and efficacy of a third dose of mRNA vaccine in patients receiving maintenance hemodialysis. <i>Kidney International</i> , 2021 ,	9.9	17
40	Myostatin and muscle atrophy during chronic kidney disease. <i>Nephrology Dialysis Transplantation</i> , 2021 , 36, 1986-1993	4.3	10
39	COVID-19 vaccine acceptance among haemodialysis patients: a French survey. <i>CKJ: Clinical Kidney Journal</i> , 2021 , 14, 1985-1986	4.5	O
38	Evolution of renal function in patients with severe intestinal failure on home parenteral nutrition. <i>CKJ: Clinical Kidney Journal</i> , 2021 , 14, 925-932	4.5	2
37	New clinical evidences for urea toxicity. Nephrology Dialysis Transplantation, 2021,	4.3	1
36	A low aromatic amino-acid diet improves renal function and prevent kidney fibrosis in mice with chronic kidney disease. <i>Scientific Reports</i> , 2021 , 11, 19184	4.9	2
35	The ROMANOV study found impaired humoral and cellular immune responses to SARS-CoV-2 mRNA vaccine in virus-unexposed patients receiving maintenance hemodialysis. <i>Kidney International</i> , 2021 , 100, 928-936	9.9	18
34	The protein-bound uremic toxin p-cresyl-sulfate promotes intracellular ROS production and lipid peroxidation in 3T3-L1 adipose cells. <i>Biochimie</i> , 2021 , 189, 137-143	4.6	2
33	P0922A LOW AROMATIC AMINO-ACID DIET IMPROVES RENAL FUNCTION AND PREVENTS KIDNEY FIBROSIS IN MICE WITH CHRONIC KIDNEY DISEASE. <i>Nephrology Dialysis Transplantation</i> , 2020 , 35,	4.3	1
32	Chronic Kidney Disease-Associated Immune Dysfunctions: Impact of Protein-Bound Uremic Retention Solutes on Immune Cells. <i>Toxins</i> , 2020 , 12,	4.9	23
31	Effects of Fecal Microbiota Transplantation on Composition in Mice with CKD. <i>Toxins</i> , 2020 , 12,	4.9	19
30	Accumulation of natriuretic peptides is associated with protein energy wasting and activation of browning in white adipose tissue in chronic kidney disease. <i>Kidney International</i> , 2020 , 98, 663-672	9.9	2
29	Source and Composition in Amino Acid of Dietary Proteins in the Primary Prevention and Treatment of CKD. <i>Nutrients</i> , 2020 , 12,	6.7	3
28	Ketoacid Analogues Supplementation in Chronic Kidney Disease and Future Perspectives. <i>Nutrients</i> , 2019 , 11,	6.7	31
27	Vegetarian diets and chronic kidney disease. <i>Nephrology Dialysis Transplantation</i> , 2019 , 34, 199-207	4.3	36

(2014-2019)

26	The Role for Protein Restriction in Addition to Renin-Angiotensin-Aldosterone System Inhibitors in the Management of CKD. <i>American Journal of Kidney Diseases</i> , 2019 , 73, 248-257	7.4	36
25	Is there still a place for prebiotics in chronic kidney disease?. <i>Nephrology Dialysis Transplantation</i> , 2019 , 34, 1812-1816	4.3	3
24	Emerging role of myostatin and its inhibition in the setting of chronic kidney disease. <i>Kidney International</i> , 2019 , 95, 506-517	9.9	33
23	Is 3-Carboxy-4-methyl-5-propyl-2-furanpropionate (CMPF) a Clinically Relevant Uremic Toxin in Haemodialysis Patients?. <i>Toxins</i> , 2018 , 10,	4.9	8
22	The Role of Gut Microbiota and Diet on Uremic Retention Solutes Production in the Context of Chronic Kidney Disease. <i>Toxins</i> , 2018 , 10,	4.9	34
21	Distal Colon Motor Dysfunction in Mice with Chronic Kidney Disease: Putative Role of Uremic Toxins. <i>Toxins</i> , 2018 , 10,	4.9	17
20	SaO045ACTIVATION OF BROWNING IN WHITE ADIPOSE TISSUE DURING CHRONIC KIDNEY DISEASE. <i>Nephrology Dialysis Transplantation</i> , 2018 , 33, i334-i334	4.3	
19	Serum levels of the adipokine zinc-alpha2-glycoprotein (ZAG) predict mortality in hemodialysis patients. <i>Kidney International</i> , 2018 , 94, 983-992	9.9	8
18	Metabolic Abnormalities in Diabetes and Kidney Disease: Role of Uremic Toxins. <i>Current Diabetes Reports</i> , 2018 , 18, 97	5.6	20
17	SP351INTEREST OF FREE VITAMIN D IN CKD. Nephrology Dialysis Transplantation, 2017 , 32, iii228-iii228	4.3	O
16	p-Cresyl glucuronide is a major metabolite of p-cresol in mouse: in contrast to p-cresyl sulphate, p-cresyl glucuronide fails to promote insulin resistance. <i>Nephrology Dialysis Transplantation</i> , 2017 , 32, 2000-2009	4.3	11
15	Microbiota and prebiotics modulation of uremic toxin generation. <i>Panminerva Medica</i> , 2017 , 59, 173-18	72	17
14	Urea impairs Itell glycolysis and insulin secretion in chronic kidney disease. <i>Journal of Clinical Investigation</i> , 2016 , 126, 3598-612	15.9	53
13	CMPF: A Biomarker for Type 2 Diabetes Mellitus Progression?. <i>Trends in Endocrinology and Metabolism</i> , 2016 , 27, 439-440	8.8	15
12	Probiotics and chronic kidney disease. <i>Kidney International</i> , 2015 , 88, 958-66	9.9	118
11	Ozone exposure triggers insulin resistance through muscle c-Jun N-terminal kinase activation. <i>Diabetes</i> , 2015 , 64, 1011-24	0.9	53
10	Nutrition: Intradialytic oral nutritionthe ultimate conviction. <i>Nature Reviews Nephrology</i> , 2014 , 10, 11-	214.9	1
9	Role of altered intestinal microbiota in systemic inflammation and cardiovascular disease in chronic kidney disease. <i>Future Microbiology</i> , 2014 , 9, 399-410	2.9	100

8	Insulin resistance in chronic kidney disease: new lessons from experimental models. <i>Nephrology Dialysis Transplantation</i> , 2014 , 29, 1666-74	4.3	50
7	The relationship between renal function and plasma concentration of the cachectic factor zinc-alpha2-glycoprotein (ZAG) in adult patients with chronic kidney disease. <i>PLoS ONE</i> , 2014 , 9, e10347	73 ^{.7}	17
6	Protein-bound uremic toxinsliew targets to prevent insulin resistance and dysmetabolism in patients with chronic kidney disease. <i>Journal of Renal Nutrition</i> , 2013 , 23, 464-6	3	23
5	Ectopic lipid accumulation: A potential cause for metabolic disturbances and a contributor to the alteration of kidney function. <i>Biochimie</i> , 2013 , 95, 1971-9	4.6	85
4	White adipose tissue overproduces the lipid-mobilizing factor zinc 🛭-glycoprotein in chronic kidney disease. <i>Kidney International</i> , 2013 , 83, 878-86	9.9	35
3	p-Cresyl sulfate promotes insulin resistance associated with CKD. <i>Journal of the American Society of Nephrology: JASN</i> , 2013 , 24, 88-99	12.7	161
2	Performance of creatinine-based equations compared in older patients. <i>Journal of Nephrology</i> , 2013 , 26, 716-23	4.8	55
1	Human uremic plasma and not urea induces exuberant secretion of leptin in 3T3-L1 adipocytes. Journal of Renal Nutrition, 2011, 21, 72-5	3	15