Yi-Pu Chen

List of Publications by Year in descending order

Source: https://exaly.com/author-pdf/4827469/publications.pdf

Version: 2024-02-01

		1039406	996533
17	223	9	15
papers	citations	h-index	g-index
18	18	18	326
all docs	docs citations	times ranked	citing authors

#	Article	IF	CITATIONS
1	Retrospective Study of Phospholipase A2 Receptor and IgG Subclasses in Glomerular Deposits in Chinese Patients with Membranous Nephropathy. PLoS ONE, 2016, 11, e0156263.	1.1	40
2	Artificially Cultivated <i>Ophiocordyceps sinensis</i> Alleviates Diabetic Nephropathy and Its Podocyte Injury via Inhibiting P2X7R Expression and NLRP3 Inflammasome Activation. Journal of Diabetes Research, 2018, 2018, 1-16.	1.0	33
3	JNK-dependent AP-1 activation is required for aristolochic acid-induced TGF- $\hat{1}^2$ 1 synthesis in human renal proximal epithelial cells. American Journal of Physiology - Renal Physiology, 2012, 302, F1569-F1575.	1.3	25
4	The Protective Effects of Curcumin on Obesity-Related Glomerulopathy Are Associated with Inhibition of Wnt/ <i<math>\hat{l}^2-Catenin Signaling Activation in Podocytes. Evidence-based Complementary and Alternative Medicine, 2015, 2015, 1-12.</i<math>	0.5	17
5	Cryoglobulinemic vasculitis and glomerulonephritis. Chinese Medical Journal, 2019, 132, 1723-1732.	0.9	15
6	AJKD Atlas of Renal Pathology: Fibronectin Glomerulopathy. American Journal of Kidney Diseases, 2017, 70, e21-e22.	2.1	14
7	Aldosterone is involved in the pathogenesis of obesity-related glomerulopathy through activation of Wnt/ \hat{l}^2 -catenin signaling in podocytes. Molecular Medicine Reports, 2018, 17, 4589-4598.	1.1	13
8	Purinergic 2X7 Receptor is Involved in the Podocyte Damage of Obesity-Related Glomerulopathy via Activating Nucleotide-Binding and Oligomerization Domain-Like Receptor Protein 3 Inflammasome. Chinese Medical Journal, 2018, 131, 2713-2725.	0.9	13
9	Inhibition of local aldosterone by eplerenone reduces renal structural damage in a novel model of chronic cyclosporine A nephrotoxicity. JRAAS - Journal of the Renin-Angiotensin-Aldosterone System, 2015, 16, 301-310.	1.0	11
10	Derivation and validation of a prediction score for acute kidney injury secondary to acute myocardial infarction in Chinese patients. BMC Nephrology, 2019, 20, 195.	0.8	10
11	Clinical Prediction Scores for Type 1 Cardiorenal Syndrome Derived and Validated in Chinese Cohorts. CardioRenal Medicine, 2015, 5, 12-19.	0.7	8
12	Hirsutella sinensis Attenuates Aristolochic Acid-Induced Renal Tubular Epithelial-Mesenchymal Transition by Inhibiting TGF- \hat{l}^21 and Snail Expression. PLoS ONE, 2016, 11, e0149242.	1.1	6
13	SMRT sequencing revealed to be an effective method for ADTKD-MUC1 diagnosis through follow-up analysis of a Chinese family. Scientific Reports, 2020, 10, 8616.	1.6	6
14	Aristolochic Acid-Induced Autophagy Promotes Epithelial-to-Myofibroblast Transition in Human Renal Proximal Tubule Epithelial Cells. Evidence-based Complementary and Alternative Medicine, 2017, 2017, 1-13.	0.5	5
15	Comparative efficacy and safety of oral anticoagulants for the treatment of venous thromboembolism in the patients with different renal functions: a systematic review, pairwise and network meta-analysis. BMJ Open, 2022, 12, e048619.	0.8	3
16	Two kinds of rare light chain cast nephropathy caused by multiple myeloma: case reports and literature review. BMC Nephrology, 2021, 22, 42.	0.8	2
17	Renal Parenchymal Malakoplakia Presenting as Acute Renal Failure in a Young Woman. Chinese Medical Journal, 2016, 129, 1880-1881.	0.9	1