

Robert N Luben

List of Publications by Year in Descending Order

Source: <https://exaly.com/author-pdf/4831885/robert-n-luben-publications-by-year.pdf>

Version: 2024-04-10

This document has been generated based on the publications and citations recorded by exaly.com. For the latest version of this publication list, visit the link given above.

The third column is the impact factor (IF) of the journal, and the fourth column is the number of citations of the article.

523 papers	47,480 citations	102 h-index	201 g-index
555 ext. papers	53,448 ext. citations	7.6 avg, IF	6.71 L-index

#	Paper	IF	Citations
523	Is there a dose-response relationship between musical instrument playing and later-life cognition? A cohort study using EPIC-Norfolk data. <i>Age and Ageing</i> , 2021 , 50, 220-226	3	1
522	Germline variants and breast cancer survival in patients with distant metastases at primary breast cancer diagnosis. <i>Scientific Reports</i> , 2021 , 11, 19787	4.9	0
521	Calcium intake, calcium supplementation and cardiovascular disease and mortality in the British population: EPIC-norfolk prospective cohort study and meta-analysis. <i>European Journal of Epidemiology</i> , 2021 , 36, 669-683	12.1	0
520	The Relationship Between Cognitive Performance Using Tests Assessing a Range of Cognitive Domains and Future Dementia Diagnosis in a British Cohort: A Ten-Year Prospective Study. <i>Journal of Alzheimer's Disease</i> , 2021 , 81, 123-135	4.3	1
519	Alcohol Consumption and Incident Cataract Surgery in Two Large UK Cohorts. <i>Ophthalmology</i> , 2021 , 128, 837-847	7.3	4
518	The relationship between alcohol intake and falls hospitalization: Results from the EPIC-Norfolk. <i>Geriatrics and Gerontology International</i> , 2021 , 21, 657-663	2.9	3
517	Association between serum secretory phospholipase A2 and risk of ischaemic stroke. <i>European Journal of Neurology</i> , 2021 , 28, 3650-3655	6	1
516	Long Term Prognostic Impact of Sex-specific Longitudinal Changes in Blood Pressure. The EPIC-Norfolk Prospective Population Cohort Study. <i>European Journal of Preventive Cardiology</i> , 2021 , ,	3.9	2
515	Obesity is Associated With Increased Risk of Crohn's disease, but not Ulcerative Colitis: A Pooled Analysis of Five Prospective Cohort Studies. <i>Clinical Gastroenterology and Hepatology</i> , 2021 ,	6.9	4
514	Baseline anticholinergic burden from medications predicts poorer baseline and long-term health-related quality of life in 16 675 men and women of EPIC-Norfolk prospective population-based cohort study. <i>Pharmacoepidemiology and Drug Safety</i> , 2021 , 30, 135-143	2.6	1
513	Long-term effects of gestational diabetes on bone mineral density and fracture risk: Analysis of the Norfolk cohort of the European Prospective Investigation into Cancer (EPIC-Norfolk) population-based study. <i>Maturitas</i> , 2021 , 144, 68-73	5	2
512	Cross-sectional and prospective associations between active living environments and accelerometer-assessed physical activity in the EPIC-Norfolk cohort. <i>Health and Place</i> , 2021 , 67, 102490	4.6	1
511	Positive Associations of Dietary Intake and Plasma Concentrations of Vitamin E with Skeletal Muscle Mass, Heel Bone Ultrasound Attenuation and Fracture Risk in the EPIC-Norfolk Cohort. <i>Antioxidants</i> , 2021 , 10,	7.1	3
510	Genome-wide meta-analysis identifies 127 open-angle glaucoma loci with consistent effect across ancestries. <i>Nature Communications</i> , 2021 , 12, 1258	17.4	47
509	Association of germline genetic variants with breast cancer-specific survival in patient subgroups defined by clinic-pathological variables related to tumor biology and type of systemic treatment. <i>Breast Cancer Research</i> , 2021 , 23, 86	8.3	1
508	Usual physical activity and subsequent hospital usage over 20 years in a general population: the EPIC-Norfolk cohort. <i>BMC Geriatrics</i> , 2020 , 20, 165	4.1	5
507	Germline HOXB13 mutations p.G84E and p.R217C do not confer an increased breast cancer risk. <i>Scientific Reports</i> , 2020 , 10, 9688	4.9	2

506	Cross-sectional and prospective relationship between occupational and leisure-time inactivity and cognitive function in an ageing population: the European Prospective Investigation into Cancer and Nutrition in Norfolk (EPIC-Norfolk) study. <i>International Journal of Epidemiology</i> , 2020 , 49, 1338-1352	7.8	
505	A Mediterranean Diet Is Positively Associated with Bone and Muscle Health in a Non-Mediterranean Region in 25,450 Men and Women from EPIC-Norfolk. <i>Nutrients</i> , 2020 , 12,	6.7	7
504	The associations of longitudinal changes in consumption of total and types of dairy products and markers of metabolic risk and adiposity: findings from the European Investigation into Cancer and Nutrition (EPIC)-Norfolk study, United Kingdom. <i>American Journal of Clinical Nutrition</i> , 2020 , 111, 1018-1026	7	16
503	Retinal vasculometric characteristics and their associations with polymyalgia rheumatica and giant cell arteritis in a prospective cohort: EPIC-Norfolk Eye Study. <i>Annals of the Rheumatic Diseases</i> , 2020 , 79, 547-549	2.4	
502	Use of Medications with Anticholinergic Properties and the Long-Term Risk of Hospitalization for Falls and Fractures in the EPIC-Norfolk Longitudinal Cohort Study. <i>Drugs and Aging</i> , 2020 , 37, 105-114	4.7	7
501	Biomarker-estimated flavan-3-ol intake is associated with lower blood pressure in cross-sectional analysis in EPIC Norfolk. <i>Scientific Reports</i> , 2020 , 10, 17964	4.9	11
500	Anticholinergic medication exposure predicts poor physical capability: Findings from a large prospective cohort study in England. <i>Maturitas</i> , 2020 , 142, 55-63	5	2
499	Retinal Vasculometry Associations With Glaucoma: Findings From the European Prospective Investigation of Cancer-Norfolk Eye Study. <i>American Journal of Ophthalmology</i> , 2020 , 220, 140-151	4.9	2
498	Lower Dietary and Circulating Vitamin C in Middle- and Older-Aged Men and Women Are Associated with Lower Estimated Skeletal Muscle Mass. <i>Journal of Nutrition</i> , 2020 , 150, 2789-2798	4.1	12
497	Sociodemographic and lifestyle predictors of incident hospital admissions with multimorbidity in a general population, 1999-2019: the EPIC-Norfolk cohort. <i>BMJ Open</i> , 2020 , 10, e042115	3	3
496	Dietary acid-base load and its association with risk of osteoporotic fractures and low estimated skeletal muscle mass. <i>European Journal of Clinical Nutrition</i> , 2020 , 74, 33-42	5.2	3
495	Evaluation of (-)-epicatechin metabolites as recovery biomarker of dietary flavan-3-ol intake. <i>Scientific Reports</i> , 2019 , 9, 13108	4.9	9
494	Plasma vitamin C concentrations and risk of incident respiratory diseases and mortality in the European Prospective Investigation into Cancer-Norfolk population-based cohort study. <i>European Journal of Clinical Nutrition</i> , 2019 , 73, 1492-1500	5.2	11
493	Joint association of mammographic density adjusted for age and body mass index and polygenic risk score with breast cancer risk. <i>Breast Cancer Research</i> , 2019 , 21, 68	8.3	18
492	FEV1 and total Cardiovascular mortality and morbidity over an 18 years follow-up Population-Based Prospective EPIC-NORFOLK Study. <i>BMC Public Health</i> , 2019 , 19, 501	4.1	5
491	Risk factors for herpes simplex virus type-1 infection and reactivation: Cross-sectional studies among EPIC-Norfolk participants. <i>PLoS ONE</i> , 2019 , 14, e0215553	3.7	8
490	Changes in plasma phospholipid fatty acid profiles over 13 years and correlates of change: European Prospective Investigation into Cancer and Nutrition-Norfolk Study. <i>American Journal of Clinical Nutrition</i> , 2019 , 109, 1527-1534	7	7
489	Biomarkers of Dietary Omega-6 Fatty Acids and Incident Cardiovascular Disease and Mortality. <i>Circulation</i> , 2019 , 139, 2422-2436	16.7	118

488	Plasma Vitamin C Levels: Risk Factors for Deficiency and Association with Self-Reported Functional Health in the European Prospective Investigation into Cancer-Norfolk. <i>Nutrients</i> , 2019 , 11,	6.7	15
487	Physical activity trajectories and mortality: population based cohort study. <i>BMJ, The</i> , 2019 , 365, l2323	5.9	86
486	Disentangling the genetics of lean mass. <i>American Journal of Clinical Nutrition</i> , 2019 , 109, 276-287	7	24
485	Changes in waist circumference and risk of all-cause and CVD mortality: results from the European Prospective Investigation into Cancer in Norfolk (EPIC-Norfolk) cohort study. <i>BMC Cardiovascular Disorders</i> , 2019 , 19, 238	2.3	16
484	The :p.Arg658* truncating variant is associated with risk of triple-negative breast cancer. <i>Npj Breast Cancer</i> , 2019 , 5, 38	7.8	12
483	Low-dose methotrexate: potential clinical impact on haematological and constitutional symptoms in myeloproliferative neoplasms. <i>British Journal of Haematology</i> , 2019 , 187, e69-e72	4.5	1
482	Residential area deprivation and risk of subsequent hospital admission in a British population: the EPIC-Norfolk cohort. <i>BMJ Open</i> , 2019 , 9, e031251	3	2
481	Development and Validation of Lifestyle-Based Models to Predict Incidence of the Most Common Potentially Preventable Cancers. <i>Cancer Epidemiology Biomarkers and Prevention</i> , 2019 , 28, 67-75	4	8
480	Retinal Vasculometry Associations with Cardiometabolic Risk Factors in the European Prospective Investigation of Cancer-Norfolk Study. <i>Ophthalmology</i> , 2019 , 126, 96-106	7.3	20
479	Dietary Fiber and the Risk of Pancreatic Cancer. <i>Pancreas</i> , 2019 , 48, 121-125	2.6	5
478	Targeted Resequencing of the Coding Sequence of 38 Genes Near Breast Cancer GWAS Loci in a Large Case-Control Study. <i>Cancer Epidemiology Biomarkers and Prevention</i> , 2019 , 28, 822-825	4	4
477	Incidence of inflammatory polyarthritis in polymyalgia rheumatica: a population-based cohort study. <i>Annals of the Rheumatic Diseases</i> , 2019 , 78, 704-705	2.4	1
476	Cross-sectional associations of dietary and circulating magnesium with skeletal muscle mass in the EPIC-Norfolk cohort. <i>Clinical Nutrition</i> , 2019 , 38, 317-323	5.9	15
475	Low thyroid function is not associated with an accelerated deterioration in renal function. <i>Nephrology Dialysis Transplantation</i> , 2019 , 34, 650-659	4.3	20
474	Baseline anticholinergic burden from medications predicts incident fatal and non-fatal stroke in the EPIC-Norfolk general population. <i>International Journal of Epidemiology</i> , 2018 , 47, 625-633	7.8	15
473	Randomised trial of coconut oil, olive oil or butter on blood lipids and other cardiovascular risk factors in healthy men and women. <i>BMJ Open</i> , 2018 , 8, e020167	3	81
472	Fibre intake and the development of inflammatory bowel disease: A European prospective multi-centre cohort study (EPIC-IBD). <i>Journal of Crohns and Colitis</i> , 2018 , 12, 129-136	1.5	48
471	Weight change and 15-year mortality: results from the European Prospective Investigation into Cancer in Norfolk (EPIC-Norfolk) cohort study. <i>European Journal of Epidemiology</i> , 2018 , 33, 37-53	12.1	18

470	Prediagnostic Serum Vitamin D Levels and the Risk of Crohn's Disease and Ulcerative Colitis in European Populations: A Nested Case-Control Study. <i>Inflammatory Bowel Diseases</i> , 2018 , 24, 633-640	4.5	28
469	Physical capability predicts mortality in late mid-life as well as in old age: Findings from a large British cohort study. <i>Archives of Gerontology and Geriatrics</i> , 2018 , 74, 77-82	4	15
468	The association between physical activity and the risk of symptomatic Barrett's oesophagus: a UK prospective cohort study. <i>European Journal of Gastroenterology and Hepatology</i> , 2018 , 30, 71-75	2.2	4
467	Low Free Testosterone and Prostate Cancer Risk: A Collaborative Analysis of 20 Prospective Studies. <i>European Urology</i> , 2018 , 74, 585-594	10.2	36
466	Dietary oleic acid is inversely associated with pancreatic cancer - Data from food diaries in a cohort study. <i>Pancreatology</i> , 2018 , 18, 655-660	3.8	12
465	Prediction of acute myeloid leukaemia risk in healthy individuals. <i>Nature</i> , 2018 , 559, 400-404	50.4	368
464	Effect of Long-Term Low Lipoproteins on Neurocognitive Function. <i>Journal of the American College of Cardiology</i> , 2018 , 72, 1176-1177	15.1	1
463	Individual and combined impact of lifestyle factors on atrial fibrillation in apparently healthy men and women: The EPIC-Norfolk prospective population study. <i>European Journal of Preventive Cardiology</i> , 2018 , 25, 1374-1383	3.9	10
462	Polyclonal human antibodies against glycans bearing red meat-derived non-human sialic acid N-glycolylneuraminic acid are stable, reproducible, complex and vary between individuals: Total antibody levels are associated with colorectal cancer risk. <i>PLoS ONE</i> , 2018 , 13, e0197464	3.7	32
461	Association between intake of less-healthy foods defined by the United Kingdom's nutrient profile model and cardiovascular disease: A population-based cohort study. <i>PLoS Medicine</i> , 2018 , 15, e1002484	11.6	18
460	Impact of physical activity on the risk of cardiovascular disease in middle-aged and older adults: EPIC Norfolk prospective population study. <i>European Journal of Preventive Cardiology</i> , 2018 , 25, 200-208	3.9	54
459	Dimension of pain-related quality of life and self-reported mental health in men and women of the European Prospective Investigation into Cancer-Norfolk cohort: a population-based cross-sectional study. <i>British Journal of Pain</i> , 2018 , 12, 35-46	2.1	
458	A Common Glaucoma-risk Variant of SIX6 Alters Retinal Nerve Fiber Layer and Optic Disc Measures in a European Population: The EPIC-Norfolk Eye Study. <i>Journal of Glaucoma</i> , 2018 , 27, 743-749	2.1	10
457	Descriptive epidemiology of changes in objectively measured sedentary behaviour and physical activity: six-year follow-up of the EPIC-Norfolk cohort. <i>International Journal of Behavioral Nutrition and Physical Activity</i> , 2018 , 15, 122	8.4	8
456	The Relation Between Thyroid Function and Anemia: A Pooled Analysis of Individual Participant Data. <i>Journal of Clinical Endocrinology and Metabolism</i> , 2018 , 103, 3658-3667	5.6	19
455	Understanding the relationship between cognition and death: a within cohort examination of cognitive measures and mortality. <i>European Journal of Epidemiology</i> , 2018 , 33, 1049-1062	12.1	18
454	Alcohol consumption and future hospital usage: The EPIC-Norfolk prospective population study. <i>PLoS ONE</i> , 2018 , 13, e0200747	3.7	2
453	Dietary antioxidant intake and the risk of developing Barrett's oesophagus and oesophageal adenocarcinoma. <i>British Journal of Cancer</i> , 2018 , 118, 1658-1661	8.7	4

452	Circulating isoflavone and lignan concentrations and prostate cancer risk: a meta-analysis of individual participant data from seven prospective studies including 2,828 cases and 5,593 controls. <i>International Journal of Cancer</i> , 2018 , 143, 2677-2686	7.5	13
451	Evaluation at scale of microbiome-derived metabolites as biomarker of flavan-3-ol intake in epidemiological studies. <i>Scientific Reports</i> , 2018 , 8, 9859	4.9	30
450	New insights into the genetics of primary open-angle glaucoma based on meta-analyses of intraocular pressure and optic disc characteristics. <i>Human Molecular Genetics</i> , 2017 , 26, 438-453	5.6	80
449	No association of alcohol use and the risk of ulcerative colitis or Crohn's disease: data from a European Prospective cohort study (EPIC). <i>European Journal of Clinical Nutrition</i> , 2017 , 71, 512-518	5.2	33
448	Thyroid Function Tests in the Reference Range and Fracture: Individual Participant Analysis of Prospective Cohorts. <i>Journal of Clinical Endocrinology and Metabolism</i> , 2017 , 102, 2719-2728	5.6	27
447	Genome-wide meta-analysis of 241,258 adults accounting for smoking behaviour identifies novel loci for obesity traits. <i>Nature Communications</i> , 2017 , 8, 14977	17.4	105
446	Higher Meat Intake Is Positively Associated With Higher Risk of Developing Pancreatic Cancer in an Age-Dependent Manner and Are Modified by Plasma Antioxidants: A Prospective Cohort Study (EPIC-Norfolk) Using Data From Food Diaries. <i>Pancreas</i> , 2017 , 46, 672-678	2.6	10
445	Changes in physical activity following total hip or knee arthroplasty: a matched case-control study from the EPIC-Norfolk cohort. <i>Clinical Rehabilitation</i> , 2017 , 31, 1548-1557	3.3	7
444	Carotenoid dietary intakes and plasma concentrations are associated with heel bone ultrasound attenuation and osteoporotic fracture risk in the European Prospective Investigation into Cancer and Nutrition (EPIC)-Norfolk cohort. <i>British Journal of Nutrition</i> , 2017 , 117, 1439-1453	3.6	27
443	Sulfate, nitrate and blood pressure - An EPIC interaction between sulfur and nitrogen. <i>Pharmacological Research</i> , 2017 , 122, 127-129	10.2	10
442	Longitudinal associations between marine omega-3 supplement users and coronary heart disease in a UK population-based cohort. <i>BMJ Open</i> , 2017 , 7, e017471	3	6
441	Omega-6 fatty acid biomarkers and incident type 2 diabetes: pooled analysis of individual-level data for 39 740 adults from 20 prospective cohort studies. <i>Lancet Diabetes and Endocrinology</i> , 2017 , 5, 965-974	18.1	150
440	Thyroid Function Within the Normal Range, Subclinical Hypothyroidism, and the Risk of Atrial Fibrillation. <i>Circulation</i> , 2017 , 136, 2100-2116	16.7	95
439	Weather, day length and physical activity in older adults: Cross-sectional results from the European Prospective Investigation into Cancer and Nutrition (EPIC) Norfolk Cohort. <i>PLoS ONE</i> , 2017 , 12, e0177767	3.7	35
438	Dietary Polyphenols in the Aetiology of Crohn's Disease and Ulcerative Colitis-A Multicenter European Prospective Cohort Study (EPIC). <i>Inflammatory Bowel Diseases</i> , 2017 , 23, 2072-2082	4.5	18
437	Dog ownership supports the maintenance of physical activity during poor weather in older English adults: cross-sectional results from the EPIC Norfolk cohort. <i>Journal of Epidemiology and Community Health</i> , 2017 ,	5.1	29
436	Prediagnostic Serum Vitamin D Levels and Risk of Inflammatory Bowel Disease: A Pan-European, Nested Case-control Study. <i>Gastroenterology</i> , 2017 , 152, S59	13.3	2
435	Large meta-analysis of genome-wide association studies identifies five loci for lean body mass. <i>Nature Communications</i> , 2017 , 8, 80	17.4	88

434	Rare, protein-truncating variants in , and , but not , are associated with increased breast cancer risks. <i>Journal of Medical Genetics</i> , 2017 , 54, 732-741	5.8	47
433	Genetic modifiers of CHEK2*1100delC-associated breast cancer risk. <i>Genetics in Medicine</i> , 2017 , 19, 599-603		51
432	Glaucoma and intraocular pressure in EPIC-Norfolk Eye Study: cross sectional study. <i>BMJ, The</i> , 2017 , 358, j3889	5.9	47
431	Genome-wide physical activity interactions in adiposity - A meta-analysis of 200,452 adults. <i>PLoS Genetics</i> , 2017 , 13, e1006528	6	103
430	Ideal cardiovascular health and risk of cardiovascular events in the EPIC-Norfolk prospective population study. <i>European Journal of Preventive Cardiology</i> , 2016 , 23, 986-94	3.9	36
429	A principal component meta-analysis on multiple anthropometric traits identifies novel loci for body shape. <i>Nature Communications</i> , 2016 , 7, 13357	17.4	46
428	Long-term mortality of hospitalized pneumonia in the EPIC-Norfolk cohort. <i>Epidemiology and Infection</i> , 2016 , 144, 803-9	4.3	6
427	Daytime napping, sleep duration and increased 8-year risk of type 2 diabetes in a British population. <i>Nutrition, Metabolism and Cardiovascular Diseases</i> , 2016 , 26, 996-1003	4.5	32
426	Meta-analysis of gene-environment-wide association scans accounting for education level identifies additional loci for refractive error. <i>Nature Communications</i> , 2016 , 7, 11008	17.4	79
425	Genome-wide analysis identifies 12 loci influencing human reproductive behavior. <i>Nature Genetics</i> , 2016 , 48, 1462-1472	36.3	198
424	Predicting admissions and time spent in hospital over a decade in a population-based record linkage study: the EPIC-Norfolk cohort. <i>BMJ Open</i> , 2016 , 6, e009461	3	12
423	Fatigue is associated with excess mortality in the general population: results from the EPIC-Norfolk study. <i>BMC Medicine</i> , 2016 , 14, 122	11.4	28
422	Dairy Products, Dietary Calcium, and Risk of Inflammatory Bowel Disease: Results From a European Prospective Cohort Investigation. <i>Inflammatory Bowel Diseases</i> , 2016 , 22, 1403-11	4.5	55
421	The descriptive epidemiology of accelerometer-measured physical activity in older adults. <i>International Journal of Behavioral Nutrition and Physical Activity</i> , 2016 , 13, 2	8.4	65
420	EB Polyunsaturated Fatty Acid Biomarkers and Coronary Heart Disease: Pooling Project of 19 Cohort Studies. <i>JAMA Internal Medicine</i> , 2016 , 176, 1155-66	11.5	238
419	Investigating Physical Activity in the Etiology of Pancreatic Cancer: The Age at Which This Is Measured Is Important and Is Independent of Body Mass Index. <i>Pancreas</i> , 2016 , 45, 388-93	2.6	8
418	CYP19A1 fine-mapping and Mendelian randomization: estradiol is causal for endometrial cancer. <i>Endocrine-Related Cancer</i> , 2016 , 23, 77-91	5.7	41
417	Reply to W Lin and R Wang. <i>American Journal of Clinical Nutrition</i> , 2016 , 103, 290-1	7	

416	Habitual chocolate consumption and the risk of incident heart failure among healthy men and women. <i>Nutrition, Metabolism and Cardiovascular Diseases</i> , 2016 , 26, 722-34	4.5	10
415	No evidence that protein truncating variants in BRIP1 are associated with breast cancer risk: implications for gene panel testing. <i>Journal of Medical Genetics</i> , 2016 , 53, 298-309	5.8	83
414	Breast cancer risk variants at 6q25 display different phenotype associations and regulate ESR1, RMND1 and CCDC170. <i>Nature Genetics</i> , 2016 , 48, 374-86	36.3	93
413	Heterogeneous impact of classic atherosclerotic risk factors on different arterial territories: the EPIC-Norfolk prospective population study. <i>European Heart Journal</i> , 2016 , 37, 880-9	9.5	24
412	RAD51B in Familial Breast Cancer. <i>PLoS ONE</i> , 2016 , 11, e0153788	3.7	18
411	Cross Sectional Associations between Socio-Demographic Factors and Cognitive Performance in an Older British Population: The European Investigation of Cancer in Norfolk (EPIC-Norfolk) Study. <i>PLoS ONE</i> , 2016 , 11, e0166779	3.7	6
410	Genome-wide association reveals that common genetic variation in the kallikrein-kinin system is associated with serum L-arginine levels. <i>Thrombosis and Haemostasis</i> , 2016 , 116, 1041-1049	7	5
409	Retinal Nerve Fiber Layer Measures and Cognitive Function in the EPIC-Norfolk Cohort Study 2016 , 57, 1921-6		17
408	Genetically Predicted Body Mass Index and Breast Cancer Risk: Mendelian Randomization Analyses of Data from 145,000 Women of European Descent. <i>PLoS Medicine</i> , 2016 , 13, e1002105	11.6	80
407	Fracture Risk in Relation to Serum 25-Hydroxyvitamin D and Physical Activity: Results from the EPIC-Norfolk Cohort Study. <i>PLoS ONE</i> , 2016 , 11, e0164160	3.7	9
406	Dietary Patterns and Risk of Inflammatory Bowel Disease in Europe: Results from the EPIC Study. <i>Inflammatory Bowel Diseases</i> , 2016 , 22, 345-54	4.5	127
405	Objective Sedentary Time, Moderate-to-Vigorous Physical Activity, and Physical Capability in a British Cohort. <i>Medicine and Science in Sports and Exercise</i> , 2016 , 48, 421-9	1.2	29
404	Thyroid dysfunction and anaemia in a large population-based study. <i>Clinical Endocrinology</i> , 2016 , 84, 627-31	3.4	24
403	Associations with intraocular pressure across Europe: The European Eye Epidemiology (E) Consortium. <i>European Journal of Epidemiology</i> , 2016 , 31, 1101-1111	12.1	16
402	Association of genetic susceptibility variants for type 2 diabetes with breast cancer risk in women of European ancestry. <i>Cancer Causes and Control</i> , 2016 , 27, 679-93	2.8	15
401	Daytime napping and increased risk of incident respiratory diseases: symptom, marker, or risk factor?. <i>Sleep Medicine</i> , 2016 , 23, 12-15	4.6	15
400	Thyroid Function Within the Reference Range and the Risk of Stroke: An Individual Participant Data Analysis. <i>Journal of Clinical Endocrinology and Metabolism</i> , 2016 , 101, 4270-4282	5.6	50
399	Topical Beta-Blockers and Cardiovascular Mortality: Systematic Review and Meta-Analysis with Data from the EPIC-Norfolk Cohort Study. <i>Ophthalmic Epidemiology</i> , 2016 , 23, 277-84	1.9	3

398	Exposure to Ambient Air Pollution and the Risk of Inflammatory Bowel Disease: A European Nested Case-Control Study. <i>Digestive Diseases and Sciences</i> , 2016 , 61, 2963-2971	4	30
397	Cross-sectional associations between different measures of obesity and muscle strength in men and women in a British cohort study. <i>Journal of Nutrition, Health and Aging</i> , 2015 , 19, 3-11	5.2	48
396	Sleep duration and risk of fatal and nonfatal stroke: a prospective study and meta-analysis. <i>Neurology</i> , 2015 , 84, 1072-9	6.5	134
395	Area deprivation and age related macular degeneration in the EPIC-Norfolk Eye Study. <i>Public Health</i> , 2015 , 129, 103-9	4	9
394	Habitual chocolate consumption and risk of cardiovascular disease among healthy men and women. <i>Heart</i> , 2015 , 101, 1279-87	5.1	53
393	Associations between flavan-3-ol intake and CVD risk in the Norfolk cohort of the European Prospective Investigation into Cancer (EPIC-Norfolk). <i>Free Radical Biology and Medicine</i> , 2015 , 84, 1-10	7.8	28
392	Subclinical Hypothyroidism and the Risk of Stroke Events and Fatal Stroke: An Individual Participant Data Analysis. <i>Journal of Clinical Endocrinology and Metabolism</i> , 2015 , 100, 2181-91	5.6	112
391	Increasing Prevalence of Myopia in Europe and the Impact of Education. <i>Ophthalmology</i> , 2015 , 122, 1489-97	9.7	220
390	Polymorphisms in a Putative Enhancer at the 10q21.2 Breast Cancer Risk Locus Regulate NRBF2 Expression. <i>American Journal of Human Genetics</i> , 2015 , 97, 22-34	11	26
389	Prevalence of refractive error in Europe: the European Eye Epidemiology (E(3)) Consortium. <i>European Journal of Epidemiology</i> , 2015 , 30, 305-15	12.1	193
388	Performance of the CHARGE-AF risk model for incident atrial fibrillation in the EPIC Norfolk cohort. <i>European Journal of Preventive Cardiology</i> , 2015 , 22, 932-9	3.9	44
387	Assessment of variation in immunosuppressive pathway genes reveals TGFBR2 to be associated with prognosis of estrogen receptor-negative breast cancer after chemotherapy. <i>Breast Cancer Research</i> , 2015 , 17, 18	8.3	17
386	Total anticholinergic burden and risk of mortality and cardiovascular disease over 10 years in 21,636 middle-aged and older men and women of EPIC-Norfolk prospective population study. <i>Age and Ageing</i> , 2015 , 44, 219-25	3	58
385	Novel Associations between Common Breast Cancer Susceptibility Variants and Risk-Predicting Mammographic Density Measures. <i>Cancer Research</i> , 2015 , 75, 2457-67	10.1	45
384	Dietary magnesium and potassium intakes and circulating magnesium are associated with heel bone ultrasound attenuation and osteoporotic fracture risk in the EPIC-Norfolk cohort study. <i>American Journal of Clinical Nutrition</i> , 2015 , 102, 376-84	7	44
383	Reply to WB Grant. <i>American Journal of Clinical Nutrition</i> , 2015 , 102, 230-1	7	
382	Inverse association between bone mineral density and risk of aortic stenosis in men and women in EPIC-Norfolk prospective study. <i>International Journal of Cardiology</i> , 2015 , 178, 29-30	3.2	11
381	Genome-wide meta-analysis identifies six novel loci associated with habitual coffee consumption. <i>Molecular Psychiatry</i> , 2015 , 20, 647-656	15.1	167

380	Association between sucrose intake and risk of overweight and obesity in a prospective sub-cohort of the European Prospective Investigation into Cancer in Norfolk (EPIC-Norfolk). <i>Public Health Nutrition</i> , 2015 , 18, 2815-24	3.3	31
379	Longitudinal association of C-reactive protein and Haemoglobin A1c over 13 years: the European Prospective Investigation into Cancer--Norfolk study. <i>Cardiovascular Diabetology</i> , 2015 , 14, 61	8.7	7
378	A comprehensive evaluation of interaction between genetic variants and use of menopausal hormone therapy on mammographic density. <i>Breast Cancer Research</i> , 2015 , 17, 110	8.3	13
377	The Influence of Age and Sex on Genetic Associations with Adult Body Size and Shape: A Large-Scale Genome-Wide Interaction Study. <i>PLoS Genetics</i> , 2015 , 11, e1005378	6	220
376	Opposites don't attract: high spouse concordance for dietary supplement use in the European Prospective Investigation into Cancer in Norfolk (EPIC-Norfolk) cohort study. <i>Public Health Nutrition</i> , 2015 , 18, 1060-6	3.3	5
375	The association between a biomarker score for fruit and vegetable intake and incident type 2 diabetes: the EPIC-Norfolk study. <i>European Journal of Clinical Nutrition</i> , 2015 , 69, 449-54	5.2	35
374	Prediction of breast cancer risk based on profiling with common genetic variants. <i>Journal of the National Cancer Institute</i> , 2015 , 107,	9.7	324
373	The relationship between dietary magnesium intake, stroke and its major risk factors, blood pressure and cholesterol, in the EPIC-Norfolk cohort. <i>International Journal of Cardiology</i> , 2015 , 196, 108-14	3.2	29
372	Television viewing, walking speed, and grip strength in a prospective cohort study. <i>Medicine and Science in Sports and Exercise</i> , 2015 , 47, 735-42	1.2	16
371	Subclinical thyroid dysfunction and fracture risk: a meta-analysis. <i>JAMA - Journal of the American Medical Association</i> , 2015 , 313, 2055-65	27.4	190
370	Association of HDL cholesterol efflux capacity with incident coronary heart disease events: a prospective case-control study. <i>Lancet Diabetes and Endocrinology</i> , 2015 , 3, 507-13	18.1	300
369	Parkinson's Disease Case Ascertainment in the EPIC Cohort: The NeuroEPIC4PD Study. <i>Neurodegenerative Diseases</i> , 2015 , 15, 331-8	2.3	8
368	Cross-sectional and prospective associations between dietary and plasma vitamin C, heel bone ultrasound, and fracture risk in men and women in the European Prospective Investigation into Cancer in Norfolk cohort. <i>American Journal of Clinical Nutrition</i> , 2015 , 102, 1416-24	7	15
367	Greater accordance with the Dietary Approaches to Stop Hypertension dietary pattern is associated with lower diet-related greenhouse gas production but higher dietary costs in the United Kingdom. <i>American Journal of Clinical Nutrition</i> , 2015 , 102, 138-45	7	60
366	Reply: The importance of ventricular-vascular uncoupling. <i>JACC: Heart Failure</i> , 2015 , 3, 95-96	7.9	
365	Contribution of cod liver oil-related nutrients (vitamins A, D, E and eicosapentaenoic acid and docosahexaenoic acid) to daily nutrient intake and their associations with plasma concentrations in the EPIC-Norfolk cohort. <i>Journal of Human Nutrition and Dietetics</i> , 2015 , 28, 568-82	3.1	11
364	Flavonoid intake in European adults (18 to 64 years). <i>PLoS ONE</i> , 2015 , 10, e0128132	3.7	113
363	Cross Sectional and Longitudinal Associations between Cardiovascular Risk Factors and Age Related Macular Degeneration in the EPIC-Norfolk Eye Study. <i>PLoS ONE</i> , 2015 , 10, e0132565	3.7	23

362	The SNP rs6500843 in 16p13.3 is associated with survival specifically among chemotherapy-treated breast cancer patients. <i>Oncotarget</i> , 2015 , 6, 7390-407	3.3	14
361	Estimated dietary intakes and sources of flavanols in the German population (German National Nutrition Survey II). <i>European Journal of Nutrition</i> , 2014 , 53, 635-43	5.2	15
360	Dietary dairy product intake and incident type 2 diabetes: a prospective study using dietary data from a 7-day food diary. <i>Diabetologia</i> , 2014 , 57, 909-17	10.3	126
359	Body mass index, smoking, and alcohol and risks of Barrett's esophagus and esophageal adenocarcinoma: a UK prospective cohort study. <i>Digestive Diseases and Sciences</i> , 2014 , 59, 1552-9	4	34
358	Daytime napping and the risk of all-cause and cause-specific mortality: a 13-year follow-up of a British population. <i>American Journal of Epidemiology</i> , 2014 , 179, 1115-24	3.8	75
357	Assessment of the dietary intake of total flavan-3-ols, monomeric flavan-3-ols, proanthocyanidins and theaflavins in the European Union. <i>British Journal of Nutrition</i> , 2014 , 111, 1463-73	3.6	77
356	Visual acuity, self-reported vision and falls in the EPIC-Norfolk Eye study. <i>British Journal of Ophthalmology</i> , 2014 , 98, 377-82	5.5	49
355	Body fat percentage, body mass index and waist-to-hip ratio as predictors of mortality and cardiovascular disease. <i>Heart</i> , 2014 , 100, 1613-9	5.1	79
354	Genome-wide association study identifies multiple loci associated with both mammographic density and breast cancer risk. <i>Nature Communications</i> , 2014 , 5, 5303	17.4	84
353	A systematic review and meta-analysis of 130,000 individuals shows smoking does not modify the association of APOE genotype on risk of coronary heart disease. <i>Atherosclerosis</i> , 2014 , 237, 5-12	3.1	24
352	Estimated urinary sodium excretion and risk of heart failure in men and women in the EPIC-Norfolk study. <i>European Journal of Heart Failure</i> , 2014 , 16, 394-402	12.3	66
351	Using lifestyle factors to identify individuals at higher risk of inflammatory polyarthritis (results from the European Prospective Investigation of Cancer-Norfolk and the Norfolk Arthritis Register--the EPIC-2-NOAR Study). <i>Annals of the Rheumatic Diseases</i> , 2014 , 73, 219-26	2.4	64
350	Low bone mineral density predicts incident heart failure in men and women: the EPIC (European Prospective Investigation into Cancer and Nutrition)-Norfolk prospective study. <i>JACC: Heart Failure</i> , 2014 , 2, 380-9	7.9	25
349	Distribution of lipid parameters according to different socio-economic indicators- the EPIC-Norfolk prospective population study. <i>BMC Public Health</i> , 2014 , 14, 782	4.1	20
348	Evidence that breast cancer risk at the 2q35 locus is mediated through IGFBP5 regulation. <i>Nature Communications</i> , 2014 , 4, 4999	17.4	87
347	Bone mineral density and incidence of stroke: European prospective investigation into cancer-norfolk population-based study, systematic review, and meta-analysis. <i>Stroke</i> , 2014 , 45, 373-82	6.7	26
346	Systemic medication and intraocular pressure in a British population: the EPIC-Norfolk Eye Study. <i>Ophthalmology</i> , 2014 , 121, 1501-7	7.3	45
345	MicroRNA related polymorphisms and breast cancer risk. <i>PLoS ONE</i> , 2014 , 9, e109973	3.7	37

344	Cod liver oil supplement consumption and health: cross-sectional results from the EPIC-Norfolk cohort study. <i>Nutrients</i> , 2014 , 6, 4320-37	6.7	5
343	Daytime napping, sleep duration and serum C reactive protein: a population-based cohort study. <i>BMJ Open</i> , 2014 , 4, e006071	3	38
342	A genome-wide copy number association study of osteoporotic fractures points to the 6p25.1 locus. <i>Journal of Medical Genetics</i> , 2014 , 51, 122-31	5.8	32
341	Dietary intake measurement using 7 d diet diaries in British men and women in the European Prospective Investigation into Cancer-Norfolk study: a focus on methodological issues. <i>British Journal of Nutrition</i> , 2014 , 111, 516-26	3.6	33
340	Cohort profile: A prospective cohort study of objective physical and cognitive capability and visual health in an ageing population of men and women in Norfolk (EPIC-Norfolk 3). <i>International Journal of Epidemiology</i> , 2014 , 43, 1063-72	7.8	60
339	Area deprivation, individual socioeconomic status and low vision in the EPIC-Norfolk Eye Study. <i>Journal of Epidemiology and Community Health</i> , 2014 , 68, 204-10	5.1	23
338	Serum 25-hydroxyvitamin D, mortality, and incident cardiovascular disease, respiratory disease, cancers, and fractures: a 13-y prospective population study. <i>American Journal of Clinical Nutrition</i> , 2014 , 100, 1361-70	7	60
337	A new tool for converting food frequency questionnaire data into nutrient and food group values: FETA research methods and availability. <i>BMJ Open</i> , 2014 , 4, e004503	3	95
336	Longitudinal association of C-reactive protein and lung function over 13 years: The EPIC-Norfolk study. <i>American Journal of Epidemiology</i> , 2014 , 179, 48-56	3.8	49
335	2q36.3 is associated with prognosis for oestrogen receptor-negative breast cancer patients treated with chemotherapy. <i>Nature Communications</i> , 2014 , 5, 4051	17.4	13
334	Serum carbon and nitrogen stable isotopes as potential biomarkers of dietary intake and their relation with incident type 2 diabetes: the EPIC-Norfolk study. <i>American Journal of Clinical Nutrition</i> , 2014 , 100, 708-18	7	29
333	Association between high dietary intake of the n-3 polyunsaturated fatty acid docosahexaenoic acid and reduced risk of Crohn's disease. <i>Alimentary Pharmacology and Therapeutics</i> , 2014 , 39, 834-42	6.1	49
332	Plasma vitamin C and risk of hospitalisation with diagnosis of atrial fibrillation in men and women in EPIC-Norfolk prospective study. <i>International Journal of Cardiology</i> , 2014 , 177, 830-5	3.2	12
331	FTO genetic variants, dietary intake and body mass index: insights from 177,330 individuals. <i>Human Molecular Genetics</i> , 2014 , 23, 6961-72	5.6	120
330	Genetic determinants of heel bone properties: genome-wide association meta-analysis and replication in the GEFOS/GENOMOS consortium. <i>Human Molecular Genetics</i> , 2014 , 23, 3054-68	5.6	78
329	Dietary intake of carbohydrates and risk of type 2 diabetes: the European Prospective Investigation into Cancer-Norfolk study. <i>British Journal of Nutrition</i> , 2014 , 111, 342-52	3.6	20
328	Cognitive function in a general population of men and women: a cross sectional study in the European Investigation of Cancer-Norfolk cohort (EPIC-Norfolk). <i>BMC Geriatrics</i> , 2014 , 14, 142	4.1	20
327	Carbohydrate intake in the etiology of Crohn's disease and ulcerative colitis. <i>Inflammatory Bowel Diseases</i> , 2014 , 20, 2013-21	4.5	58

326	Dietary arachidonic and oleic acid intake in ulcerative colitis etiology: a prospective cohort study using 7-day food diaries. <i>European Journal of Gastroenterology and Hepatology</i> , 2014 , 26, 11-8	2.2	46
325	The association between Mediterranean Diet Score and glucokinase regulatory protein gene variation on the markers of cardiometabolic risk: an analysis in the European Prospective Investigation into Cancer (EPIC)-Norfolk study. <i>British Journal of Nutrition</i> , 2014 , 112, 122-31	3.6	17
324	Corneal biomechanical properties and glaucoma-related quantitative traits in the EPIC-Norfolk Eye Study 2014 , 55, 117-24		22
323	Self-reported sleep patterns in a British population cohort. <i>Sleep Medicine</i> , 2014 , 15, 295-302	4.6	40
322	Social adversity experience and blood pressure control following antihypertensive medication use in a community sample of older adults. <i>International Journal of Behavioral Medicine</i> , 2014 , 21, 456-63	2.6	
321	Abstract 19753: Hdl Cholesterol Efflux Capacity is Inversely Associated With Incident Chd Events Independent of Hdl-c and APOA-I Concentrations. <i>Circulation</i> , 2014 , 130,	16.7	2
320	Seventeen year risk of all-cause and cause-specific mortality associated with C-reactive protein, fibrinogen and leukocyte count in men and women: the EPIC-Norfolk study. <i>European Journal of Epidemiology</i> , 2013 , 28, 541-50	12.1	54
319	Multiple independent variants at the TERT locus are associated with telomere length and risks of breast and ovarian cancer. <i>Nature Genetics</i> , 2013 , 45, 371-84, 384e1-2	36.3	422
318	The physical capability of community-based men and women from a British cohort: the European Prospective Investigation into Cancer (EPIC)-Norfolk study. <i>BMC Geriatrics</i> , 2013 , 13, 93	4.1	22
317	Prediction of measured weight from self-reported weight was not improved after stratification by body mass index. <i>Obesity</i> , 2013 , 21, E137-42	8	13
316	C-reactive protein and fracture risk: European prospective investigation into Cancer Norfolk Study. <i>Bone</i> , 2013 , 56, 67-72	4.7	20
315	Does ICD-10 hospital discharge code I50 identify people with heart failure? A validation study within the EPIC-Norfolk study. <i>International Journal of Cardiology</i> , 2013 , 168, 4413-4	3.2	9
314	Rate of weight gain predicts change in physical activity levels: a longitudinal analysis of the EPIC-Norfolk cohort. <i>International Journal of Obesity</i> , 2013 , 37, 404-9	5.5	51
313	Large-scale genotyping identifies 41 new loci associated with breast cancer risk. <i>Nature Genetics</i> , 2013 , 45, 353-61, 361e1-2	36.3	813
312	Body mass index and the risk for Crohn's disease and ulcerative colitis: data from a European Prospective Cohort Study (The IBD in EPIC Study). <i>American Journal of Gastroenterology</i> , 2013 , 108, 575-82	8.7	113
311	Identification of heart rate-associated loci and their effects on cardiac conduction and rhythm disorders. <i>Nature Genetics</i> , 2013 , 45, 621-31	36.3	219
310	Dietary antioxidants and the aetiology of pancreatic cancer: a cohort study using data from food diaries and biomarkers. <i>Gut</i> , 2013 , 62, 1489-96	19.2	38
309	Demographic and disease-related predictors of abnormal lung function in patients with established inflammatory polyarthritis and a comparison with the general population. <i>Annals of the Rheumatic Diseases</i> , 2013 , 72, 1517-23	2.4	5

308	The small eye phenotype in the EPIC-Norfolk eye study: prevalence and visual impairment in microphthalmos and nanophthalmos. <i>BMJ Open</i> , 2013 , 3,	3	6
307	Genome-wide association study of intraocular pressure identifies the GLCCI1/ICA1 region as a glaucoma susceptibility locus. <i>Human Molecular Genetics</i> , 2013 , 22, 4653-60	5.6	24
306	A genome-wide association scan (GWAS) for mean telomere length within the COGS project: identified loci show little association with hormone-related cancer risk. <i>Human Molecular Genetics</i> , 2013 , 22, 5056-64	5.6	107
305	Seropositivity and higher immunoglobulin g antibody levels against cytomegalovirus are associated with mortality in the population-based European prospective investigation of Cancer-Norfolk cohort. <i>Clinical Infectious Diseases</i> , 2013 , 56, 1421-7	11.6	68
304	Laser scanning tomography in the EPIC-Norfolk Eye Study: principal components and associations 2013 , 54, 6638-45		11
303	Genotype-phenotype analysis of SNPs associated with primary angle closure glaucoma (rs1015213, rs3753841 and rs11024102) and ocular biometry in the EPIC-Norfolk Eye Study. <i>British Journal of Ophthalmology</i> , 2013 , 97, 704-7	5.5	20
302	The association between social stress and global cognitive function in a population-based study: the European Prospective Investigation into Cancer (EPIC)-Norfolk study. <i>Psychological Medicine</i> , 2013 , 43, 655-66	6.9	11
301	Differences in dietary supplement use and secular and seasonal trends assessed using three different instruments in the EPIC-Norfolk population study. <i>Journal of Dietary Supplements</i> , 2013 , 10, 142-51	2.3	6
300	Common genetic variants of the natriuretic peptide gene locus are not associated with heart failure risk in participants in the EPIC-Norfolk study. <i>European Journal of Heart Failure</i> , 2013 , 15, 624-7	12.3	7
299	Distribution and determinants of C-reactive protein in the older adult population: European Prospective Investigation into Cancer-Norfolk study. <i>European Journal of Clinical Investigation</i> , 2013 , 43, 899-911	4.6	21
298	The association of cycling with all-cause, cardiovascular and cancer mortality: findings from the population-based EPIC-Norfolk cohort. <i>BMJ Open</i> , 2013 , 3, e003797	3	31
297	The EPIC-Norfolk Eye Study: rationale, methods and a cross-sectional analysis of visual impairment in a population-based cohort. <i>BMJ Open</i> , 2013 , 3,	3	33
296	Associations with retinal nerve fiber layer measures in the EPIC-Norfolk Eye Study 2013 , 54, 5028-34		40
295	Self-rated health and cardiovascular disease incidence: results from a longitudinal population-based cohort in Norfolk, UK. <i>PLoS ONE</i> , 2013 , 8, e65290	3.7	44
294	Body fat mass is a predictor of risk of osteoporotic fractures in women but not in men: a prospective population study. <i>Journal of Internal Medicine</i> , 2012 , 271, 472-80	10.8	35
293	The association between prior infection with five serotypes of Coxsackievirus B and incident type 2 diabetes mellitus in the EPIC-Norfolk study. <i>Diabetologia</i> , 2012 , 55, 967-70	10.3	2
292	Dietary, lifestyle and clinicopathological factors associated with APC mutations and promoter methylation in colorectal cancers from the EPIC-Norfolk study. <i>Journal of Pathology</i> , 2012 , 228, 405-15	9.4	43
291	Resting heart rate and incident heart failure in apparently healthy men and women in the EPIC-Norfolk study. <i>European Journal of Heart Failure</i> , 2012 , 14, 1163-70	12.3	26

290	Genome-wide meta-analysis identifies 56 bone mineral density loci and reveals 14 loci associated with risk of fracture. <i>Nature Genetics</i> , 2012 , 44, 491-501	36.3	866
289	Testosterone, SHBG and differential white blood cell count in middle-aged and older men. <i>Maturitas</i> , 2012 , 71, 274-8	5	12
288	Educational attainment and mean leukocyte telomere length in women in the European Prospective Investigation into Cancer (EPIC)-Norfolk population study. <i>Brain, Behavior, and Immunity</i> , 2012 , 26, 414-8	16.6	22
287	Influence of inflammatory polyarthritis on quantitative heel ultrasound measurements. <i>BMC Musculoskeletal Disorders</i> , 2012 , 13, 133	2.8	1
286	A latent variable partial least squares path modeling approach to regional association and polygenic effect with applications to a human obesity study. <i>PLoS ONE</i> , 2012 , 7, e31927	3.7	9
285	Subclinical thyroid dysfunction and the risk of heart failure events: an individual participant data analysis from 6 prospective cohorts. <i>Circulation</i> , 2012 , 126, 1040-9	16.7	315
284	Circulating 25-hydroxyvitamin D concentration and the risk of type 2 diabetes: results from the European Prospective Investigation into Cancer (EPIC)-Norfolk cohort and updated meta-analysis of prospective studies. <i>Diabetologia</i> , 2012 , 55, 2173-82	10.3	181
283	Self-reported and measured anthropometric data and risk of colorectal cancer in the EPIC-Norfolk study. <i>International Journal of Obesity</i> , 2012 , 36, 107-18	5.5	28
282	Plasma phospholipid fatty acid concentration and incident coronary heart disease in men and women: the EPIC-Norfolk prospective study. <i>PLoS Medicine</i> , 2012 , 9, e1001255	11.6	160
281	Urinary bisphenol A concentration and risk of future coronary artery disease in apparently healthy men and women. <i>Circulation</i> , 2012 , 125, 1482-90	16.7	199
280	Lipoprotein(a) and risk of coronary, cerebrovascular, and peripheral artery disease: the EPIC-Norfolk prospective population study. <i>Arteriosclerosis, Thrombosis, and Vascular Biology</i> , 2012 , 32, 3058-65	9.4	97
279	Uncorrected refractive error in older British adults: the EPIC-Norfolk Eye Study. <i>British Journal of Ophthalmology</i> , 2012 , 96, 991-6	5.5	21
278	A prospective study of the association between quantity and variety of fruit and vegetable intake and incident type 2 diabetes. <i>Diabetes Care</i> , 2012 , 35, 1293-300	14.6	143
277	Common breast cancer susceptibility variants in LSP1 and RAD51L1 are associated with mammographic density measures that predict breast cancer risk. <i>Cancer Epidemiology Biomarkers and Prevention</i> , 2012 , 21, 1156-66	4	92
276	Identification of a novel percent mammographic density locus at 12q24. <i>Human Molecular Genetics</i> , 2012 , 21, 3299-305	5.6	28
275	Higher immunoglobulin G antibody levels against cytomegalovirus are associated with incident ischemic heart disease in the population-based EPIC-Norfolk cohort. <i>Journal of Infectious Diseases</i> , 2012 , 206, 1897-903	7	51
274	Mammographic breast density and breast cancer: evidence of a shared genetic basis. <i>Cancer Research</i> , 2012 , 72, 1478-84	10.1	50
273	Relationship between plasma fibrinogen and fiber intake in the EPIC-Norfolk cohort. <i>European Journal of Clinical Nutrition</i> , 2012 , 66, 443-51	5.2	6

272	Fibre intake in relation to serum total cholesterol levels and CHD risk: a comparison of dietary assessment methods. <i>European Journal of Clinical Nutrition</i> , 2012 , 66, 296-304	5.2	11
271	Intake estimation of total and individual flavan-3-ols, proanthocyanidins and theaflavins, their food sources and determinants in the European Prospective Investigation into Cancer and Nutrition (EPIC) study. <i>British Journal of Nutrition</i> , 2012 , 108, 1095-108	3.6	81
270	Subclinical hyperthyroidism and the risk of coronary heart disease and mortality. <i>Archives of Internal Medicine</i> , 2012 , 172, 799-809		331
269	Differential white blood cell count and incident heart failure in men and women in the EPIC-Norfolk study. <i>European Heart Journal</i> , 2012 , 33, 523-30	9.5	37
268	Genetic variation near IRS1 associates with reduced adiposity and an impaired metabolic profile. <i>Nature Genetics</i> , 2011 , 43, 753-60	36.3	237
267	Plasma vitamin C predicts incident heart failure in men and women in European Prospective Investigation into Cancer and Nutrition-Norfolk prospective study. <i>American Heart Journal</i> , 2011 , 162, 246-53	4.9	67
266	Full-scale scores of the Mini Mental State Examination can be generated from an abbreviated version. <i>Journal of Clinical Epidemiology</i> , 2011 , 64, 1005-13	5.7	13
265	Intraocular pressure and corneal biomechanics in an adult British population: the EPIC-Norfolk eye study 2011 , 52, 8179-85		45
264	Mendelian Randomisation Study of Childhood BMI and Early Menarche. <i>Journal of Obesity</i> , 2011 , 2011, 180729	3.7	48
263	Physical activity and ocular perfusion pressure: the EPIC-Norfolk eye study 2011 , 52, 8186-92		31
262	The aetiology of symptomatic gallstones quantification of the effects of obesity, alcohol and serum lipids on risk. Epidemiological and biomarker data from a UK prospective cohort study (EPIC-Norfolk). <i>European Journal of Gastroenterology and Hepatology</i> , 2011 , 23, 733-40	2.2	38
261	Aspirin in the aetiology of Crohn's disease and ulcerative colitis: a European prospective cohort study. <i>Alimentary Pharmacology and Therapeutics</i> , 2011 , 34, 649-55	6.1	44
260	Associations of endogenous testosterone and SHBG with glycated haemoglobin in middle-aged and older men. <i>Clinical Endocrinology</i> , 2011 , 74, 572-8	3.4	33
259	Effects of body size and sociodemographic characteristics on differences between self-reported and measured anthropometric data in middle-aged men and women: the EPIC-Norfolk study. <i>European Journal of Clinical Nutrition</i> , 2011 , 65, 357-67	5.2	37
258	MLH1 promoter methylation, diet, and lifestyle factors in mismatch repair deficient colorectal cancer patients from EPIC-Norfolk. <i>Nutrition and Cancer</i> , 2011 , 63, 1000-10	2.8	26
257	Genetic predisposition to obesity leads to increased risk of type 2 diabetes. <i>Diabetologia</i> , 2011 , 54, 776-82	2.3	49
256	No evidence of an increased mortality risk associated with low levels of glycated haemoglobin in a non-diabetic UK population. <i>Diabetologia</i> , 2011 , 54, 2025-32	10.3	27
255	Individual and cumulative effect of type 2 diabetes genetic susceptibility variants on risk of coronary heart disease. <i>Diabetologia</i> , 2011 , 54, 2283-7	10.3	17

254	No evidence for a causal link between uric acid and type 2 diabetes: a Mendelian randomisation approach. <i>Diabetologia</i> , 2011 , 54, 2561-9	10.3	73
253	Alterations in PTEN and PIK3CA in colorectal cancers in the EPIC Norfolk study: associations with clinicopathological and dietary factors. <i>BMC Cancer</i> , 2011 , 11, 123	4.8	60
252	SMAD7 and MGMT genotype variants and cancer incidence in the European Prospective Investigation into Cancer and Nutrition (EPIC)-Norfolk Study. <i>Cancer Epidemiology</i> , 2011 , 35, 369-74	2.8	17
251	Cigarette smoking and endogenous sex hormones in postmenopausal women. <i>Journal of Clinical Endocrinology and Metabolism</i> , 2011 , 96, 3184-92	5.6	54
250	Association between dietary phyto-oestrogens and bone density in men and postmenopausal women. <i>British Journal of Nutrition</i> , 2011 , 106, 1063-9	3.6	12
249	N-Nitroso compounds and cancer incidence: the European Prospective Investigation into Cancer and Nutrition (EPIC)-Norfolk Study. <i>American Journal of Clinical Nutrition</i> , 2011 , 93, 1053-61	7	132
248	Television viewing time independently predicts all-cause and cardiovascular mortality: the EPIC Norfolk study. <i>International Journal of Epidemiology</i> , 2011 , 40, 150-9	7.8	222
247	Association between plasma vitamin C concentrations and blood pressure in the European prospective investigation into cancer-Norfolk population-based study. <i>Hypertension</i> , 2011 , 58, 372-9	8.5	25
246	Dietary fat and breast cancer: comparison of results from food diaries and food-frequency questionnaires in the UK Dietary Cohort Consortium. <i>American Journal of Clinical Nutrition</i> , 2011 , 94, 1043-52	7	27
245	CHD risk in relation to alcohol intake from categorical and open-ended dietary instruments. <i>Public Health Nutrition</i> , 2011 , 14, 402-9	3.3	3
244	Incidence of type 2 diabetes using proposed HbA1c diagnostic criteria in the european prospective investigation of cancer-norfolk cohort: implications for preventive strategies. <i>Diabetes Care</i> , 2011 , 34, 950-6	14.6	45
243	Lifestyle behaviours and quality-adjusted life years in middle and older age. <i>Age and Ageing</i> , 2011 , 40, 589-95	3	29
242	Life stress, emotional health, and mean telomere length in the European Prospective Investigation into Cancer (EPIC)-Norfolk population study. <i>Journals of Gerontology - Series A Biological Sciences and Medical Sciences</i> , 2011 , 66, 1152-62	6.4	91
241	Mendelian randomization study of B-type natriuretic peptide and type 2 diabetes: evidence of causal association from population studies. <i>PLoS Medicine</i> , 2011 , 8, e1001112	11.6	79
240	Television viewing and incident cardiovascular disease: prospective associations and mediation analysis in the EPIC Norfolk Study. <i>PLoS ONE</i> , 2011 , 6, e20058	3.7	76
239	Prospective study of insulin-like growth factor-I, insulin-like growth factor-binding protein 3, genetic variants in the IGF1 and IGFBP3 genes and risk of coronary artery disease. <i>International Journal of Molecular Epidemiology and Genetics</i> , 2011 , 2, 261-85	0.9	8
238	Initial thyroid status and cardiovascular risk factors: the EPIC-Norfolk prospective population study. <i>Clinical Endocrinology</i> , 2010 , 72, 404-10	3.4	115
237	Alcohol intake and risk of colorectal cancer: results from the UK Dietary Cohort Consortium. <i>British Journal of Cancer</i> , 2010 , 103, 747-56	8.7	18

236	Biological, clinical and population relevance of 95 loci for blood lipids. <i>Nature</i> , 2010 , 466, 707-13	50.4	2742
235	Meta-analysis identifies 13 new loci associated with waist-hip ratio and reveals sexual dimorphism in the genetic basis of fat distribution. <i>Nature Genetics</i> , 2010 , 42, 949-60	36.3	724
234	Association analyses of 249,796 individuals reveal 18 new loci associated with body mass index. <i>Nature Genetics</i> , 2010 , 42, 937-48	36.3	2267
233	A single nucleotide polymorphism in the 3-hydroxy-3-methylglutaryl-coenzyme A reductase gene (HMGCR) influences the serum triacylglycerol relationship with dietary fat and fibre in the European Prospective Investigation into Cancer and Nutrition in Norfolk (EPIC-Norfolk) study. <i>British Journal of Nutrition</i> , 2010 , 104, 765-72	3.6	5
232	Dietary fiber and colorectal cancer risk: a nested case-control study using food diaries. <i>Journal of the National Cancer Institute</i> , 2010 , 102, 614-26	9.7	172
231	Physical functional health predicts the incidence of coronary heart disease in the European Prospective Investigation into Cancer-Norfolk prospective population-based study. <i>International Journal of Epidemiology</i> , 2010 , 39, 996-1003	7.8	12
230	Genetic variants influencing circulating lipid levels and risk of coronary artery disease. <i>Arteriosclerosis, Thrombosis, and Vascular Biology</i> , 2010 , 30, 2264-76	9.4	318
229	Breast, colorectal, and prostate cancer risk in the European Prospective Investigation into Cancer and Nutrition-Norfolk in relation to phytoestrogen intake derived from an improved database. <i>American Journal of Clinical Nutrition</i> , 2010 , 91, 440-8	7	97
228	Cumulative effects and predictive value of common obesity-susceptibility variants identified by genome-wide association studies. <i>American Journal of Clinical Nutrition</i> , 2010 , 91, 184-90	7	163
227	Telomere length in prospective and retrospective cancer case-control studies. <i>Cancer Research</i> , 2010 , 70, 3170-6	10.1	131
226	Refractive error, axial length and anterior chamber depth of the eye in British adults: the EPIC-Norfolk Eye Study. <i>British Journal of Ophthalmology</i> , 2010 , 94, 827-30	5.5	70
225	Fatty acids measured in plasma and erythrocyte-membrane phospholipids and derived by food-frequency questionnaire and the risk of new-onset type 2 diabetes: a pilot study in the European Prospective Investigation into Cancer and Nutrition (EPIC)-Norfolk cohort. <i>American Journal of Clinical Nutrition</i> , 2010 , 92, 1211-22	7	167
224	Lifestyle factors and p53 mutation patterns in colorectal cancer patients in the EPIC-Norfolk study. <i>Mutagenesis</i> , 2010 , 25, 351-8	2.8	12
223	Physical activity attenuates the genetic predisposition to obesity in 20,000 men and women from EPIC-Norfolk prospective population study. <i>PLoS Medicine</i> , 2010 , 7, e1000332	11.6	201
222	Fine scale mapping of the breast cancer 16q12 locus. <i>Human Molecular Genetics</i> , 2010 , 19, 2507-15	5.6	57
221	The Short-Form Six-Dimension utility index predicted mortality in the European Prospective Investigation into Cancer-Norfolk prospective population-based study. <i>Journal of Clinical Epidemiology</i> , 2010 , 63, 192-8	5.7	5
220	The association between physical activity in different domains of life and risk of osteoporotic fractures. <i>Bone</i> , 2010 , 47, 693-700	4.7	25
219	HMGCR gene polymorphism is associated with stroke risk in the EPIC-Norfolk study. <i>European Journal of Cardiovascular Prevention and Rehabilitation</i> , 2010 , 17, 89-93		7

218	Family history of premature coronary heart disease and risk prediction in the EPIC-Norfolk prospective population study. <i>Heart</i> , 2010 , 96, 1985-9	5.1	79
217	Physical activity reduces the risk of symptomatic gallstones: a prospective cohort study. <i>European Journal of Gastroenterology and Hepatology</i> , 2010 , 22, 983-8	2.2	39
216	Dietary n-3 polyunsaturated fatty acids and the aetiology of ulcerative colitis: a UK prospective cohort study. <i>European Journal of Gastroenterology and Hepatology</i> , 2010 , 22, 602-6	2.2	132
215	Mastery is associated with cardiovascular disease mortality in men and women at apparently low risk. <i>Health Psychology</i> , 2010 , 29, 412-20	5	48
214	Intake of dietary fats and colorectal cancer risk: prospective findings from the UK Dietary Cohort Consortium. <i>Cancer Epidemiology</i> , 2010 , 34, 562-7	2.8	20
213	No evidence that social stress is associated with breast cancer incidence. <i>Breast Cancer Research and Treatment</i> , 2010 , 120, 169-74	4.4	35
212	Meat, poultry and fish and risk of colorectal cancer: pooled analysis of data from the UK dietary cohort consortium. <i>Cancer Causes and Control</i> , 2010 , 21, 1417-25	2.8	44
211	Dietary, lifestyle and clinicopathological factors associated with BRAF and K-ras mutations arising in distinct subsets of colorectal cancers in the EPIC Norfolk study. <i>BMC Cancer</i> , 2010 , 10, 99	4.8	41
210	MGMT Ile143Val polymorphism, dietary factors and the risk of breast, colorectal and prostate cancer in the European Prospective Investigation into Cancer and Nutrition (EPIC)-Norfolk study. <i>DNA Repair</i> , 2010 , 9, 421-8	4.3	23
209	Differential white blood cell count and type 2 diabetes: systematic review and meta-analysis of cross-sectional and prospective studies. <i>PLoS ONE</i> , 2010 , 5, e13405	3.7	91
208	APOE genotype, lipids, and coronary heart disease risk: a prospective population study. <i>Archives of Internal Medicine</i> , 2009 , 169, 1424-9		62
207	Linoleic acid, a dietary n-6 polyunsaturated fatty acid, and the aetiology of ulcerative colitis: a nested case-control study within a European prospective cohort study. <i>Gut</i> , 2009 , 58, 1606-11	19.2	235
206	Social class, risk factors, and stroke incidence in men and women: a prospective study in the European prospective investigation into cancer in Norfolk cohort. <i>Stroke</i> , 2009 , 40, 1070-7	6.7	58
205	Association between type of dietary fish and seafood intake and the risk of incident type 2 diabetes: the European prospective investigation of cancer (EPIC)-Norfolk cohort study. <i>Diabetes Care</i> , 2009 , 32, 1857-63	14.6	109
204	Self-rated health does not explain the socioeconomic differential in mortality: a prospective study in the EPIC-Norfolk cohort. <i>Journal of Epidemiology and Community Health</i> , 2009 , 63, 329-31	5.1	8
203	Author's Response Associations between dietary methods and biomarkers, and between fruits and vegetables and risk of ischaemic heart disease, in the EPIC Norfolk Cohort Study: response to letter by McNeill et al.. <i>International Journal of Epidemiology</i> , 2009 , 38, 885-885	7.8	1
202	Retinol-binding protein 4 and prediction of incident coronary events in healthy men and women. <i>Journal of Clinical Endocrinology and Metabolism</i> , 2009 , 94, 255-60	5.6	51
201	Osteoprotegerin and soluble receptor activator of nuclear factor-kappaB ligand and risk for coronary events: a nested case-control approach in the prospective EPIC-Norfolk population study 1993-2003. <i>Arteriosclerosis, Thrombosis, and Vascular Biology</i> , 2009 , 29, 975-80	9.4	61

200	Early age at menarche associated with cardiovascular disease and mortality. <i>Journal of Clinical Endocrinology and Metabolism</i> , 2009 , 94, 4953-60	5.6	334
199	FGFR2 variants and breast cancer risk: fine-scale mapping using African American studies and analysis of chromatin conformation. <i>Human Molecular Genetics</i> , 2009 , 18, 1692-703	5.6	100
198	No association between APOE and major depressive disorder in a community sample of 17,507 adults. <i>Journal of Psychiatric Research</i> , 2009 , 43, 843-7	5.2	14
197	Does the association between self-rated health and mortality vary by social class?. <i>Social Science and Medicine</i> , 2009 , 68, 275-80	5.1	58
196	How far can we explain the social class differential in respiratory function? A cross-sectional population study of 21,991 men and women from EPIC-Norfolk. <i>European Journal of Epidemiology</i> , 2009 , 24, 193-201	12.1	12
195	Different measures of social class in women and mortality. <i>European Journal of Epidemiology</i> , 2009 , 24, 231-6	12.1	12
194	Estimation of absolute fracture risk among middle-aged and older men and women: the EPIC-Norfolk population cohort study. <i>European Journal of Epidemiology</i> , 2009 , 24, 259-66	12.1	8
193	Association of C-reactive protein with type 2 diabetes: prospective analysis and meta-analysis. <i>Diabetologia</i> , 2009 , 52, 1040-7	10.3	149
192	Factor VII, blood lipids and fat intake: gene-nutrient interaction and risk of coronary heart disease with the factor VII R353Q polymorphism. <i>European Journal of Clinical Nutrition</i> , 2009 , 63, 771-7	5.2	5
191	Six new loci associated with body mass index highlight a neuronal influence on body weight regulation. <i>Nature Genetics</i> , 2009 , 41, 25-34	36.3	1368
190	Newly discovered breast cancer susceptibility loci on 3p24 and 17q23.2. <i>Nature Genetics</i> , 2009 , 41, 585-90	36.3	393
189	Genome-wide association study identifies eight loci associated with blood pressure. <i>Nature Genetics</i> , 2009 , 41, 666-76	36.3	970
188	Genetic variation in LIN28B is associated with the timing of puberty. <i>Nature Genetics</i> , 2009 , 41, 729-33	36.3	258
187	Baseline alcohol consumption, type of alcoholic beverage and risk of colorectal cancer in the European Prospective Investigation into Cancer and Nutrition-Norfolk study. <i>Cancer Epidemiology</i> , 2009 , 33, 347-54	2.8	29
186	Respiratory function as a marker of bone health and fracture risk in an older population. <i>Journal of Bone and Mineral Research</i> , 2009 , 24, 956-63	6.3	9
185	Is QUS or DXA better for predicting the 10-year absolute risk of fracture?. <i>Journal of Bone and Mineral Research</i> , 2009 , 24, 1319-25	6.3	61
184	Is bowel habit linked to colorectal cancer? - Results from the EPIC-Norfolk study. <i>European Journal of Cancer</i> , 2009 , 45, 139-45	7.5	17
183	The effect of including quantitative heel ultrasound in models for estimation of 10-year absolute risk of fracture. <i>Bone</i> , 2009 , 45, 180-4	4.7	23

182	Effect of age and sex on the relationship between different socioeconomic indices and self-reported functional health in the EPIC-Norfolk population-based study. <i>Annals of Epidemiology</i> , 2009 , 19, 289-97	6.4	4
181	A HMGCR polymorphism is associated with relations between blood pressure and urinary sodium and potassium ratio in the Epic-Norfolk Study. <i>Journal of the American Society of Hypertension</i> , 2009 , 3, 238-44		6
180	Combined effect of health behaviours and risk of first ever stroke in 20,040 men and women over 11 years' follow-up in Norfolk cohort of European Prospective Investigation of Cancer (EPIC Norfolk): prospective population study. <i>BMJ, The</i> , 2009 , 338, b349	5.9	102
179	Common variants near MC4R are associated with fat mass, weight and risk of obesity. <i>Nature Genetics</i> , 2008 , 40, 768-75	36.3	1048
178	Prospective association between emotional health and clinical evidence of Parkinson's disease. <i>European Journal of Neurology</i> , 2008 , 15, 1148-54	6	35
177	Accuracy of death certification and hospital record linkage for identification of incident stroke. <i>BMC Medical Research Methodology</i> , 2008 , 8, 74	4.7	36
176	Socioeconomic position and risk of short-term weight gain: prospective study of 14,619 middle-aged men and women. <i>BMC Public Health</i> , 2008 , 8, 112	4.1	16
175	Social inequalities in self-rated health by age: cross-sectional study of 22,457 middle-aged men and women. <i>BMC Public Health</i> , 2008 , 8, 230	4.1	64
174	Cross-sectional association between fish consumption and albuminuria: the European Prospective Investigation of Cancer-Norfolk Study. <i>American Journal of Kidney Diseases</i> , 2008 , 52, 876-86	7.4	14
173	Breast cancer risk in relation to urinary and serum biomarkers of phytoestrogen exposure in the European Prospective into Cancer-Norfolk cohort study. <i>Breast Cancer Research</i> , 2008 , 10, R32	8.3	60
172	Physical activity and fibrinogen concentrations in 23,201 men and women in the EPIC-Norfolk population-based study. <i>Atherosclerosis</i> , 2008 , 198, 419-25	3.1	13
171	LDL-cholesterol concentrations: a genome-wide association study. <i>Lancet, The</i> , 2008 , 371, 483-91	40	292
170	Sense of coherence, lifestyle choices and mortality. <i>Journal of Epidemiology and Community Health</i> , 2008 , 62, 829-31	5.1	35
169	Risk factors for first-ever stroke in the EPIC-Norfolk prospective population-based study. <i>European Journal of Cardiovascular Prevention and Rehabilitation</i> , 2008 , 15, 663-9		27
168	Plasma vitamin C level, fruit and vegetable consumption, and the risk of new-onset type 2 diabetes mellitus: the European prospective investigation of cancer--Norfolk prospective study. <i>Archives of Internal Medicine</i> , 2008 , 168, 1493-9		201
167	Dietary energy density predicts the risk of incident type 2 diabetes: the European Prospective Investigation of Cancer (EPIC)-Norfolk Study. <i>Diabetes Care</i> , 2008 , 31, 2120-5	14.6	57
166	Hormone replacement therapy and symptomatic gallstones - a prospective population study in the EPIC-Norfolk cohort. <i>Digestion</i> , 2008 , 77, 4-9	3.6	15
165	Diet in the aetiology of ulcerative colitis: a European prospective cohort study. <i>Digestion</i> , 2008 , 77, 57-64	3.6	113

164	Identification of common variants in the SHBG gene affecting sex hormone-binding globulin levels and breast cancer risk in postmenopausal women. <i>Cancer Epidemiology Biomarkers and Prevention</i> , 2008 , 17, 3490-8	4	47
163	Lack of prospective associations between plasma and urinary phytoestrogens and risk of prostate or colorectal cancer in the European Prospective into Cancer-Norfolk study. <i>Cancer Epidemiology Biomarkers and Prevention</i> , 2008 , 17, 2891-4	4	53
162	Associations between dietary methods and biomarkers, and between fruits and vegetables and risk of ischaemic heart disease, in the EPIC Norfolk Cohort Study. <i>International Journal of Epidemiology</i> , 2008 , 37, 978-87	7.8	76
161	Psychological distress, major depressive disorder, and risk of stroke. <i>Neurology</i> , 2008 , 70, 788-94	6.5	112
160	Depression and ischemic heart disease mortality: evidence from the EPIC-Norfolk United Kingdom prospective cohort study. <i>American Journal of Psychiatry</i> , 2008 , 165, 515-23	11.9	154
159	Relationship between subdomains of total physical activity and mortality. <i>Medicine and Science in Sports and Exercise</i> , 2008 , 40, 1909-15	1.2	72
158	Combined Impact of Health Behaviours and Mortality in Men and Women: The EPIC-Norfolk Prospective Population Study. <i>Obstetrical and Gynecological Survey</i> , 2008 , 63, 376-377	2.4	11
157	Plasma vitamin C concentrations predict risk of incident stroke over 10 y in 20 649 participants of the European Prospective Investigation into Cancer Norfolk prospective population study. <i>American Journal of Clinical Nutrition</i> , 2008 , 87, 64-9	7	85
156	Blood pressure and interactions between the angiotensin polymorphism AGT M235T and sodium intake: a cross-sectional population study. <i>American Journal of Clinical Nutrition</i> , 2008 , 88, 392-7	7	36
155	Major depression, C-reactive protein, and incident ischemic heart disease in healthy men and women. <i>Psychosomatic Medicine</i> , 2008 , 70, 850-5	3.7	25
154	Combined impact of health behaviours and mortality in men and women: the EPIC-Norfolk prospective population study. <i>PLoS Medicine</i> , 2008 , 5, e12	11.6	512
153	Measured height loss predicts fractures in middle-aged and older men and women: the EPIC-Norfolk prospective population study. <i>Journal of Bone and Mineral Research</i> , 2008 , 23, 425-32	6.3	37
152	Occupational social class, risk factors and cardiovascular disease incidence in men and women: a prospective study in the European Prospective Investigation of Cancer and Nutrition in Norfolk (EPIC-Norfolk) cohort. <i>European Journal of Epidemiology</i> , 2008 , 23, 449-58	12.1	45
151	Occupational social class, educational level, smoking and body mass index, and cause-specific mortality in men and women: a prospective study in the European Prospective Investigation of Cancer and Nutrition in Norfolk (EPIC-Norfolk) cohort. <i>European Journal of Epidemiology</i> , 2008 , 23, 511-22	12.1	48
150	Association between age at menarche and risk of diabetes in adults: results from the EPIC-Norfolk cohort study. <i>Diabetologia</i> , 2008 , 51, 781-6	10.3	144
149	Energy intake at breakfast and weight change: prospective study of 6,764 middle-aged men and women. <i>American Journal of Epidemiology</i> , 2008 , 167, 188-92	3.8	73
148	Physical activity and mammographic breast density in the EPIC-Norfolk cohort study. <i>American Journal of Epidemiology</i> , 2008 , 167, 579-85	3.8	15
147	Past oral contraceptive and hormone therapy use and endogenous hormone concentrations in postmenopausal women. <i>Menopause</i> , 2008 , 15, 332-9	2.5	18

146	Healthy lifestyle choices: could sense of coherence aid health promotion?. <i>Journal of Epidemiology and Community Health</i> , 2007 , 61, 871-6	5.1	104
145	Serum myeloperoxidase levels are associated with the future risk of coronary artery disease in apparently healthy individuals: the EPIC-Norfolk Prospective Population Study. <i>Journal of the American College of Cardiology</i> , 2007 , 50, 159-65	15.1	483
144	Genome-wide association study identifies novel breast cancer susceptibility loci. <i>Nature</i> , 2007 , 447, 1087-93	5.4	1957
143	Differential leucocyte count and the risk of future coronary artery disease in healthy men and women: the EPIC-Norfolk Prospective Population Study. <i>Journal of Internal Medicine</i> , 2007 , 262, 678-89	10.8	61
142	A common variant of the p16(INK4a) genetic region is associated with physical function in older people. <i>Mechanisms of Ageing and Development</i> , 2007 , 128, 370-7	5.6	75
141	No association between the BDNF Val66Met polymorphism and mood status in a non-clinical community sample of 7389 older adults. <i>Journal of Psychiatric Research</i> , 2007 , 41, 404-9	5.2	66
140	The brain-derived neurotrophic factor Val66Met polymorphism is associated with sense of coherence in a non-clinical community sample of 7335 adults. <i>Journal of Psychiatric Research</i> , 2007 , 41, 707-10	5.2	11
139	Elevated serum ferritin levels predict new-onset type 2 diabetes: results from the EPIC-Norfolk prospective study. <i>Diabetologia</i> , 2007 , 50, 949-56	10.3	189
138	Apolipoprotein A-II is inversely associated with risk of future coronary artery disease. <i>Circulation</i> , 2007 , 116, 2029-35	16.7	88
137	Circulating secretory phospholipase A2 activity and risk of incident coronary events in healthy men and women: the EPIC-Norfolk study. <i>Arteriosclerosis, Thrombosis, and Vascular Biology</i> , 2007 , 27, 1177-83	9.4	89
136	Physical health-related quality of life predicts stroke in the EPIC-Norfolk. <i>Neurology</i> , 2007 , 69, 2243-8	6.5	27
135	Cholesterol levels in small LDL particles predict the risk of coronary heart disease in the EPIC-Norfolk prospective population study. <i>European Heart Journal</i> , 2007 , 28, 2770-7	9.5	44
134	Glycated hemoglobin and risk of stroke in people without known diabetes in the European Prospective Investigation into Cancer (EPIC)-Norfolk prospective population study: a threshold relationship?. <i>Stroke</i> , 2007 , 38, 271-5	6.7	52
133	Adaptation to social adversity is associated with stroke incidence: evidence from the EPIC-Norfolk prospective cohort study. <i>Stroke</i> , 2007 , 38, 1447-53	6.7	42
132	Apolipoprotein E polymorphisms, dietary fat and fibre, and serum lipids: the EPIC Norfolk study. <i>European Heart Journal</i> , 2007 , 28, 2930-6	9.5	64
131	Study of association between common variation in the insulin-like growth factor 2 gene and indices of obesity and body size in middle-aged men and women. <i>Journal of Clinical Endocrinology and Metabolism</i> , 2007 , 92, 2734-8	5.6	33
130	Plasma levels of plant sterols and the risk of coronary artery disease: the prospective EPIC-Norfolk Population Study. <i>Journal of Lipid Research</i> , 2007 , 48, 139-44	6.3	86
129	Epidemiologic assessment of sugars consumption using biomarkers: comparisons of obese and nonobese individuals in the European prospective investigation of cancer Norfolk. <i>Cancer Epidemiology Biomarkers and Prevention</i> , 2007 , 16, 1651-4	4	57

128	Body fat distribution and risk of coronary heart disease in men and women in the European Prospective Investigation Into Cancer and Nutrition in Norfolk cohort: a population-based prospective study. <i>Circulation</i> , 2007 , 116, 2933-43	16.7	352
127	Usual physical activity and endogenous sex hormones in postmenopausal women: the European prospective investigation into cancer-norfolk population study. <i>Cancer Epidemiology Biomarkers and Prevention</i> , 2007 , 16, 900-5	4	62
126	Phytoestrogen exposure is associated with circulating sex hormone levels in postmenopausal women and interact with ESR1 and NR1H2 gene variants. <i>Cancer Epidemiology Biomarkers and Prevention</i> , 2007 , 16, 1009-16	4	52
125	Endogenous testosterone and mortality due to all causes, cardiovascular disease, and cancer in men: European prospective investigation into cancer in Norfolk (EPIC-Norfolk) Prospective Population Study. <i>Circulation</i> , 2007 , 116, 2694-701	16.7	596
124	Rat Mcs5a is a compound quantitative trait locus with orthologous human loci that associate with breast cancer risk. <i>Proceedings of the National Academy of Sciences of the United States of America</i> , 2007 , 104, 6299-304	11.5	53
123	Role of the apolipoprotein B-apolipoprotein A-I ratio in cardiovascular risk assessment: a case-control analysis in EPIC-Norfolk. <i>Annals of Internal Medicine</i> , 2007 , 146, 640-8	8	116
122	Self-reported mental health-related quality of life and mortality in men and women in the European Prospective Investigation into Cancer (EPIC-Norfolk): a prospective population study. <i>Psychosomatic Medicine</i> , 2007 , 69, 410-4	3.7	20
121	Fruit and vegetable consumption and self-reported functional health in men and women in the European Prospective Investigation into Cancer-Norfolk (EPIC-Norfolk): a population-based cross-sectional study. <i>Public Health Nutrition</i> , 2007 , 10, 34-41	3.3	66
120	Modifiable lifestyle behaviors and functional health in the European Prospective Investigation into Cancer (EPIC)-Norfolk population study. <i>Preventive Medicine</i> , 2007 , 44, 109-16	4.3	23
119	Common variants in the ATM, BRCA1, BRCA2, CHEK2 and TP53 cancer susceptibility genes are unlikely to increase breast cancer risk. <i>Breast Cancer Research</i> , 2007 , 9, R27	8.3	79
118	Effect of age on the relationship of occupational social class with prevalence of modifiable cardiovascular risk factors and cardiovascular diseases. A population-based cross-sectional study from European Prospective Investigation into Cancer - Norfolk (EPIC-Norfolk). <i>Gerontology</i> , 2006 , 52, 51-8	5.5	12
117	Combined work and leisure physical activity and risk of stroke in men and women in the European prospective investigation into Cancer-Norfolk Prospective Population Study. <i>Neuroepidemiology</i> , 2006 , 27, 122-9	5.4	30
116	How predictive is breast arterial calcification of cardiovascular disease and risk factors when found at screening mammography?. <i>American Journal of Roentgenology</i> , 2006 , 187, 73-80	5.4	75
115	IGF1 and IGFBP3 tagging polymorphisms are associated with circulating levels of IGF1, IGFBP3 and risk of breast cancer. <i>Human Molecular Genetics</i> , 2006 , 15, 1-10	5.6	160
114	Dietary antioxidants and asthma in adults. <i>Thorax</i> , 2006 , 61, 388-93	7.3	81
113	Apolipoprotein A-V, triglycerides and risk of coronary artery disease: the prospective Epic-Norfolk Population Study. <i>Journal of Lipid Research</i> , 2006 , 47, 2064-70	6.3	76
112	Serum levels of mannose-binding lectin and the risk of future coronary artery disease in apparently healthy men and women. <i>Arteriosclerosis, Thrombosis, and Vascular Biology</i> , 2006 , 26, 2345-50	9.4	54
111	Phytoestrogen exposure, polymorphisms in COMT, CYP19, ESR1, and SHBG genes, and their associations with prostate cancer risk. <i>Nutrition and Cancer</i> , 2006 , 56, 31-9	2.8	49

110	Physical activity, C-reactive protein levels and the risk of future coronary artery disease in apparently healthy men and women: the EPIC-Norfolk prospective population study. <i>European Journal of Cardiovascular Prevention and Rehabilitation</i> , 2006 , 13, 970-6		16
109	Serum lipoprotein lipase concentration and risk for future coronary artery disease: the EPIC-Norfolk prospective population study. <i>Arteriosclerosis, Thrombosis, and Vascular Biology</i> , 2006 , 26, 637-42	9.4	54
108	Implications of gene-environment interaction in studies of gene variants in breast cancer: an example of dietary isoflavones and the D356N polymorphism in the sex hormone-binding globulin gene. <i>Cancer Research</i> , 2006 , 66, 8980-3	10.1	31
107	Work and leisure time physical activity assessed using a simple, pragmatic, validated questionnaire and incident cardiovascular disease and all-cause mortality in men and women: The European Prospective Investigation into Cancer in Norfolk prospective population study. <i>International Journal of Epidemiology</i> , 2006 , 35, 1034-43	7.8	126
106	Single-nucleotide polymorphisms in the RB1 gene and association with breast cancer in the British population. <i>British Journal of Cancer</i> , 2006 , 94, 1921-6	8.7	10
105	Genetic variants in epigenetic genes and breast cancer risk. <i>Carcinogenesis</i> , 2006 , 27, 1661-9	4.6	77
104	Tagging single-nucleotide polymorphisms in antioxidant defense enzymes and susceptibility to breast cancer. <i>Cancer Research</i> , 2006 , 66, 1225-33	10.1	69
103	Endogenous versus exogenous exposure to N-nitroso compounds and gastric cancer risk in the European Prospective Investigation into Cancer and Nutrition (EPIC-EURGAST) study. <i>Carcinogenesis</i> , 2006 , 27, 1497-501	4.6	128
102	C-reactive protein levels and coronary artery disease incidence and mortality in apparently healthy men and women: the EPIC-Norfolk prospective population study 1993-2003. <i>Atherosclerosis</i> , 2006 , 187, 415-22	3.1	134
101	Serum lipid concentration in relation to anthropometric indices of central and peripheral fat distribution in 20,021 British men and women: results from the EPIC-Norfolk population-based cohort study. <i>Atherosclerosis</i> , 2006 , 189, 420-7	3.1	38
100	Social adversity, the serotonin transporter (5-HTTLPR) polymorphism and major depressive disorder. <i>Biological Psychiatry</i> , 2006 , 59, 224-9	7.9	213
99	Relation between self-reported physical functional health and chronic disease mortality in men and women in the European Prospective Investigation into Cancer (EPIC-Norfolk): a prospective population study. <i>Annals of Epidemiology</i> , 2006 , 16, 492-500	6.4	55
98	Smoking predicts long-term mortality in stroke: The European Prospective Investigation into Cancer (EPIC)-Norfolk prospective population study. <i>Preventive Medicine</i> , 2006 , 42, 128-31	4.3	15
97	Mastery, sense of coherence, and mortality: evidence of independent associations from the EPIC-Norfolk Prospective Cohort Study. <i>Health Psychology</i> , 2006 , 25, 102-10	5	89
96	Habitual fish consumption and risk of incident stroke: the European Prospective Investigation into Cancer (EPIC)-Norfolk prospective population study. <i>Public Health Nutrition</i> , 2006 , 9, 882-8	3.3	50
95	Associations among mammographic density, circulating sex hormones, and polymorphisms in sex hormone metabolism genes in postmenopausal women. <i>Cancer Epidemiology Biomarkers and Prevention</i> , 2006 , 15, 1502-8	4	68
94	Fibrinogen plasma levels modify the association between the factor XIII Val34Leu variant and risk of coronary artery disease: the EPIC-Norfolk prospective population study. <i>Journal of Thrombosis and Haemostasis</i> , 2006 , 4, 2204-9	15.4	20
93	Tissue factor serum levels and the risk of future coronary artery disease in apparently healthy men and women: the EPIC-Norfolk prospective population study. <i>Journal of Thrombosis and Haemostasis</i> , 2006 , 4, 2391-6	15.4	9

92	Obesity indices and self-reported functional health in men and women in the EPIC-Norfolk. <i>Obesity</i> , 2006 , 14, 884-93	8	13
91	HapMap-based study of the 17q21 ERBB2 amplicon in susceptibility to breast cancer. <i>British Journal of Cancer</i> , 2006 , 95, 1689-95	8.7	31
90	Plasma concentrations of ascorbic acid and C-reactive protein, and risk of future coronary artery disease, in apparently healthy men and women: the EPIC-Norfolk prospective population study. <i>British Journal of Nutrition</i> , 2006 , 96, 516-22	3.6	41
89	Long-term cryoconservation and stability of vitamin C in serum samples of the European prospective investigation into cancer and nutrition. <i>Cancer Epidemiology Biomarkers and Prevention</i> , 2005 , 14, 1837-40	4	14
88	Prospective cohort study of hostility and the risk of cardiovascular disease mortality. <i>International Journal of Cardiology</i> , 2005 , 100, 155-61	3.2	64
87	The CAFE computer program for nutritional analysis of the EPIC-Norfolk Food Frequency questionnaire and identification of extreme nutrient values. <i>Journal of Human Nutrition and Dietetics</i> , 2005 , 18, 99-116	3.1	117
86	Cigarette smoking and fat distribution in 21,828 British men and women: a population-based study. <i>Obesity</i> , 2005 , 13, 1466-75		207
85	Self-reported parkinsonian symptoms in the EPIC-Norfolk cohort. <i>BMC Neurology</i> , 2005 , 5, 15	3.1	9
84	Common variation in EMSY and risk of breast and ovarian cancer: a case-control study using HapMap tagging SNPs. <i>BMC Cancer</i> , 2005 , 5, 81	4.8	14
83	Polymorphisms in the CYP19 gene may affect the positive correlations between serum and urine phytoestrogen metabolites and plasma androgen concentrations in men. <i>Journal of Nutrition</i> , 2005 , 135, 2680-6	4.1	43
82	Plasma ascorbic acid concentrations and fat distribution in 19,068 British men and women in the European Prospective Investigation into Cancer and Nutrition Norfolk cohort study. <i>American Journal of Clinical Nutrition</i> , 2005 , 82, 1203-9	7	92
81	Respiratory function and self-reported functional health: EPIC-Norfolk population study. <i>European Respiratory Journal</i> , 2005 , 26, 494-502	13.6	30
80	RESPONSE: Re: Polymorphisms Associated With Circulating Sex Hormone Levels in Postmenopausal Women. <i>Journal of the National Cancer Institute</i> , 2005 , 97, 153-154	9.7	6
79	Fibrinogen and cigarette smoking in men and women in the European Prospective Investigation into Cancer in Norfolk (EPIC-Norfolk) population. <i>European Journal of Cardiovascular Prevention and Rehabilitation</i> , 2005 , 12, 144-50		9
78	Allelic association of the human homologue of the mouse modifier Ptprij with breast cancer. <i>Human Molecular Genetics</i> , 2005 , 14, 2349-56	5.6	64
77	Polymorphisms in the initiators of RET (rearranged during transfection) signaling pathway and susceptibility to sporadic medullary thyroid carcinoma. <i>Journal of Clinical Endocrinology and Metabolism</i> , 2005 , 90, 6268-74	5.6	67
76	Further evidence for an HLA-related recessive mutation in nasopharyngeal carcinoma among the Chinese. <i>British Journal of Cancer</i> , 2005 , 92, 967-70	8.7	37
75	INS VNTR class genotype and indexes of body size and obesity: population-based studies of 7,999 middle-aged men and women. <i>Diabetes</i> , 2005 , 54, 2812-5	0.9	12

74	Measurement of fruit and vegetable consumption with diet questionnaires and implications for analyses and interpretation. <i>American Journal of Epidemiology</i> , 2005 , 161, 987-94	3.8	66
73	Serum levels of type II secretory phospholipase A2 and the risk of future coronary artery disease in apparently healthy men and women: the EPIC-Norfolk Prospective Population Study. <i>Arteriosclerosis, Thrombosis, and Vascular Biology</i> , 2005 , 25, 839-46	9.4	85
72	Fibrinogen and cigarette smoking in men and women in the European Prospective Investigation into Cancer in Norfolk (EPIC-Norfolk) population. <i>European Journal of Cardiovascular Prevention and Rehabilitation</i> , 2005 , 12, 144-150		17
71	Phytoestrogen exposure correlation with plasma estradiol in postmenopausal women in European Prospective Investigation of Cancer and Nutrition-Norfolk may involve diet-gene interactions. <i>Cancer Epidemiology Biomarkers and Prevention</i> , 2005 , 14, 213-20	4	19
70	Blood pressure and urinary sodium in men and women: the Norfolk Cohort of the European Prospective Investigation into Cancer (EPIC-Norfolk). <i>American Journal of Clinical Nutrition</i> , 2004 , 80, 1397-403	7	117
69	Adjusting for energy intake--what measure to use in nutritional epidemiological studies?. <i>International Journal of Epidemiology</i> , 2004 , 33, 1382-6	7.8	60
68	Dietary fat and the risk of clinical type 2 diabetes: the European prospective investigation of Cancer-Norfolk study. <i>American Journal of Epidemiology</i> , 2004 , 159, 73-82	3.8	78
67	Abdominal obesity and respiratory function in men and women in the EPIC-Norfolk Study, United Kingdom. <i>American Journal of Epidemiology</i> , 2004 , 159, 1140-9	3.8	162
66	Re: "Sense of coherence and mortality in men and women in the EPIC-Norfolk United Kingdom prospective cohort study". <i>American Journal of Epidemiology</i> , 2004 , 159, 1202-3; author reply 1203-4	3.8	4
65	Correlated measurement error--implications for nutritional epidemiology. <i>International Journal of Epidemiology</i> , 2004 , 33, 1373-81	7.8	60
64	Childhood smoking is an independent risk factor for obstructive airways disease in women. <i>Thorax</i> , 2004 , 59, 682-6	7.3	19
63	Traditional cardiovascular risk factors measured prior to the onset of inflammatory polyarthritis. <i>Rheumatology</i> , 2004 , 43, 731-6	3.9	33
62	Microalbuminuria independently predicts all-cause and cardiovascular mortality in a British population: The European Prospective Investigation into Cancer in Norfolk (EPIC-Norfolk) population study. <i>International Journal of Epidemiology</i> , 2004 , 33, 189-98	7.8	138
61	Microalbuminuria, cardiovascular risk factors and cardiovascular morbidity in a British population: the EPIC-Norfolk population-based study. <i>European Journal of Cardiovascular Prevention and Rehabilitation</i> , 2004 , 11, 207-13		38
60	Plasma levels of cholesteryl ester transfer protein and the risk of future coronary artery disease in apparently healthy men and women: the prospective EPIC (European Prospective Investigation into Cancer and nutrition)-Norfolk population study. <i>Circulation</i> , 2004 , 110, 1418-23	16.7	188
59	Residential area deprivation predicts fruit and vegetable consumption independently of individual educational level and occupational social class: a cross sectional population study in the Norfolk cohort of the European Prospective Investigation into Cancer (EPIC-Norfolk). <i>Journal of Epidemiology and Community Health</i> , 2004 , 58, 681-6	5.1	94
58	A prospective study of microalbuminuria and incident coronary heart disease and its prognostic significance in a British population: the EPIC-Norfolk study. <i>American Journal of Epidemiology</i> , 2004 , 159, 284-93	3.8	76
57	Vitamin C and the risk of developing inflammatory polyarthritis: prospective nested case-control study. <i>Annals of the Rheumatic Diseases</i> , 2004 , 63, 843-7	2.4	83

56	The effect of correlated measurement error in multivariate models of diet. <i>American Journal of Epidemiology</i> , 2004 , 160, 59-67	3.8	42
55	Polymorphisms associated with circulating sex hormone levels in postmenopausal women. <i>Journal of the National Cancer Institute</i> , 2004 , 96, 936-45	9.7	291
54	IL-8 plasma concentrations and the risk of future coronary artery disease in apparently healthy men and women: the EPIC-Norfolk prospective population study. <i>Arteriosclerosis, Thrombosis, and Vascular Biology</i> , 2004 , 24, 1503-8	9.4	150
53	Habitual fish consumption and glycated haemoglobin: the EPIC-Norfolk study. <i>European Journal of Clinical Nutrition</i> , 2004 , 58, 277-84	5.2	17
52	Occupational social class, educational level and area deprivation independently predict plasma ascorbic acid concentration: a cross-sectional population based study in the Norfolk cohort of the European Prospective Investigation into Cancer (EPIC-Norfolk). <i>European Journal of Clinical Nutrition</i> , 2004 , 58, 1432-5	5.2	24
51	Microalbuminuria and stroke in a British population: the European Prospective Investigation into Cancer in Norfolk (EPIC-Norfolk) population study. <i>Journal of Internal Medicine</i> , 2004 , 255, 247-56	10.8	81
50	Dietary risk factors for the development of inflammatory polyarthritis: evidence for a role of high level of red meat consumption. <i>Arthritis and Rheumatism</i> , 2004 , 50, 3804-12		108
49	Macrophage migration inhibitory factor and the risk of myocardial infarction or death due to coronary artery disease in adults without prior myocardial infarction or stroke: the EPIC-Norfolk Prospective Population study. <i>American Journal of Medicine</i> , 2004 , 117, 390-7	2.4	73
48	Prediction of total and hip fracture risk in men and women by quantitative ultrasound of the calcaneus: EPIC-Norfolk prospective population study. <i>Lancet, The</i> , 2004 , 363, 197-202	4.0	226
47	Area deprivation predicts lung function independently of education and social class. <i>European Respiratory Journal</i> , 2004 , 24, 157-61	13.6	49
46	Fat distribution, body mass index and blood pressure in 22,090 men and women in the Norfolk cohort of the European Prospective Investigation into Cancer and Nutrition (EPIC-Norfolk) study. <i>Journal of Hypertension</i> , 2004 , 22, 2067-74	1.9	99
45	Association of hemoglobin A1c with cardiovascular disease and mortality in adults: the European prospective investigation into cancer in Norfolk. <i>Annals of Internal Medicine</i> , 2004 , 141, 413-20	8	693
44	Phytoestrogen concentrations in serum and spot urine as biomarkers for dietary phytoestrogen intake and their relation to breast cancer risk in European prospective investigation of cancer and nutrition-norfolk. <i>Cancer Epidemiology Biomarkers and Prevention</i> , 2004 , 13, 698-708	4	76
43	STK15 polymorphisms and association with risk of invasive ovarian cancer. <i>Cancer Epidemiology Biomarkers and Prevention</i> , 2004 , 13, 1589-94	4	24
42	Residential area deprivation predicts smoking habit independently of individual educational level and occupational social class. A cross sectional study in the Norfolk cohort of the European Investigation into Cancer (EPIC-Norfolk). <i>Journal of Epidemiology and Community Health</i> , 2003 , 57, 270-6	5.1	139
41	Adverse experience in childhood as a developmental risk factor for altered immune status in adulthood. <i>International Journal of Behavioral Medicine</i> , 2003 , 10, 251-68	2.6	37
40	Inflammatory dispositions: a population-based study of the association between hostility and peripheral leukocyte counts. <i>Personality and Individual Differences</i> , 2003 , 35, 1271-1284	3.3	10
39	Television viewing and low participation in vigorous recreation are independently associated with obesity and markers of cardiovascular disease risk: EPIC-Norfolk population-based study. <i>European Journal of Clinical Nutrition</i> , 2003 , 57, 1089-96	5.2	241

38	Smoking status and differential white cell count in men and women in the EPIC-Norfolk population. <i>Atherosclerosis</i> , 2003 , 169, 331-7	3.1	106
37	Association of depression with peripheral leukocyte counts in EPIC-Norfolk--role of sex and cigarette smoking. <i>Journal of Psychosomatic Research</i> , 2003 , 54, 303-6	4.1	19
36	Dietary fibre in food and protection against colorectal cancer in the European Prospective Investigation into Cancer and Nutrition (EPIC): an observational study. <i>Lancet, The</i> , 2003 , 361, 1496-501	4.0	809
35	Are imprecise methods obscuring a relation between fat and breast cancer?. <i>Lancet, The</i> , 2003 , 362, 212-4	4.0	346
34	Sense of coherence and mortality in men and women in the EPIC-Norfolk United Kingdom prospective cohort study. <i>American Journal of Epidemiology</i> , 2003 , 158, 1202-9	3.8	166
33	A transforming growth factorbeta1 signal peptide variant increases secretion in vitro and is associated with increased incidence of invasive breast cancer. <i>Cancer Research</i> , 2003 , 63, 2610-5	10.1	238
32	Cross-sectional association between total level and type of alcohol consumption and glycosylated haemoglobin level: the EPIC-Norfolk Study. <i>European Journal of Clinical Nutrition</i> , 2002 , 56, 882-90	5.2	29
31	Physical inactivity is associated with lower forced expiratory volume in 1 second : European Prospective Investigation into Cancer-Norfolk Prospective Population Study. <i>American Journal of Epidemiology</i> , 2002 , 156, 139-47	3.8	83
30	Patterns of alcohol consumption in 10 European countries participating in the European Prospective Investigation into Cancer and Nutrition (EPIC) project. <i>Public Health Nutrition</i> , 2002 , 5, 1287-9	3.3	102
29	Variability of fish consumption within the 10 European countries participating in the European Investigation into Cancer and Nutrition (EPIC) study. <i>Public Health Nutrition</i> , 2002 , 5, 1273-85	3.3	203
28	Frequency of eating and concentrations of serum cholesterol in the Norfolk population of the European prospective investigation into cancer (EPIC-Norfolk): cross sectional study. <i>BMJ: British Medical Journal</i> , 2001 , 323, 1286-8		57
27	Patterns of physical activity and ultrasound attenuation by heel bone among Norfolk cohort of European Prospective Investigation of Cancer (EPIC Norfolk): population based study. <i>BMJ: British Medical Journal</i> , 2001 , 322, 140		34
26	Fruit and vegetable intake and population glycosylated haemoglobin levels: the EPIC-Norfolk Study. <i>European Journal of Clinical Nutrition</i> , 2001 , 55, 342-8	5.2	100
25	Polymorphisms in CYP1A1 and smoking: no association with breast cancer risk. <i>Carcinogenesis</i> , 2001 , 22, 1797-800	4.6	44
24	Prevalence and family history of colorectal cancer: implications for screening. <i>Journal of Medical Screening</i> , 2001 , 8, 69-72	1.4	27
23	Glycated haemoglobin, diabetes, and mortality in men in Norfolk cohort of european prospective investigation of cancer and nutrition (EPIC-Norfolk). <i>BMJ: British Medical Journal</i> , 2001 , 322, 15-8		641
22	Fat consumption and HbA(1c) levels: the EPIC-Norfolk study. <i>Diabetes Care</i> , 2001 , 24, 1911-6	14.6	60
21	Cigarette smoking and glycaemia: the EPIC-Norfolk Study. European Prospective Investigation into Cancer. <i>International Journal of Epidemiology</i> , 2001 , 30, 547-54	7.8	98

20	Self reported non-insulin dependent diabetes, family history, and risk of prevalent colorectal cancer: population based, cross sectional study. <i>Journal of Epidemiology and Community Health</i> , 2001 , 55, 804-5	5.1	15
19	Relation between plasma ascorbic acid and mortality in men and women in EPIC-Norfolk prospective study: a prospective population study. European Prospective Investigation into Cancer and Nutrition. <i>Lancet, The</i> , 2001 , 357, 657-63	4.0	435
18	Use of biological markers to validate self-reported dietary intake in a random sample of the European Prospective Investigation into Cancer United Kingdom Norfolk cohort. <i>American Journal of Clinical Nutrition</i> , 2001 , 74, 188-96	7	180
17	Nutritional methods in the European Prospective Investigation of Cancer in Norfolk. <i>Public Health Nutrition</i> , 2001 , 4, 847-58	3.3	279
16	DINER (Data Into Nutrients for Epidemiological Research) - a new data-entry program for nutritional analysis in the EPIC-Norfolk cohort and the 7-day diary method. <i>Public Health Nutrition</i> , 2001 , 4, 1253-65	3.3	114
15	Mammographic parenchymal patterns and breast cancer natural history--a case-control study. <i>Acta Oncologica</i> , 2001 , 40, 461-5	3.2	12
14	Estimation of disease progression parameters from case-control data: application to mammographic patterns and breast cancer natural history. <i>Journal of Epidemiology and Biostatistics</i> , 2001 , 6, 235-42		4
13	A common variant in BRCA2 is associated with both breast cancer risk and prenatal viability. <i>Nature Genetics</i> , 2000 , 26, 362-4	36.3	134
12	Size, node status and grade of breast tumours: association with mammographic parenchymal patterns. <i>European Radiology</i> , 2000 , 10, 157-61	8	53
11	Intake of Selected Vitamins and Bioflavonoids from Vitamin and Mineral Supplements by Subjects in the EPIC-NORFOLK Study, UK 2000 , 25-26		
10	Polymorphisms in the human aromatase cytochrome P450 gene (CYP19) and breast cancer risk. <i>Carcinogenesis</i> , 2000 , 21, 189-93	4.6	60
9	Vitamin C and hyperglycemia in the European Prospective Investigation into Cancer--Norfolk (EPIC-Norfolk) study: a population-based study. <i>Diabetes Care</i> , 2000 , 23, 726-32	14.6	110
8	High-risk mammographic parenchymal patterns, hormone replacement therapy and other risk factors: a case-control study. <i>International Journal of Epidemiology</i> , 2000 , 29, 629-36	7.8	64
7	The effect of age, sex, and education on food consumption of a middle-aged English cohort-EPIC in East Anglia. <i>Preventive Medicine</i> , 2000 , 30, 26-34	4.3	100
6	The extent of linkage disequilibrium in four populations with distinct demographic histories. <i>American Journal of Human Genetics</i> , 2000 , 67, 1544-54	11	170
5	Smoking and high-risk mammographic parenchymal patterns: a case-control study. <i>Breast Cancer Research</i> , 2000 , 2, 59-63	8.3	12
4	No association between androgen or vitamin D receptor gene polymorphisms and risk of breast cancer. <i>Carcinogenesis</i> , 1999 , 20, 2131-5	4.6	96
3	High-risk mammographic parenchymal patterns and anthropometric measures: a case-control study. <i>British Journal of Cancer</i> , 1999 , 81, 1257-61	8.7	44

2	Mammographic parenchymal patterns and mode of detection: implications for the breast screening programme. <i>Journal of Medical Screening</i> , 1998 , 5, 207-12	1.4	78
1	A large cross-ancestry meta-analysis of genome-wide association studies identifies 69 novel risk loci for primary open-angle glaucoma and includes a genetic link with Alzheimer’s disease		4