

# Kenichiro Yabuki

## List of Publications by Year in descending order

Source: <https://exaly.com/author-pdf/4836932/publications.pdf>

Version: 2024-02-01

16  
papers

180  
citations

1307594

7  
h-index

1125743

13  
g-index

19  
all docs

19  
docs citations

19  
times ranked

316  
citing authors

#	ARTICLE	IF	CITATIONS
1	Predictive value of the Hyodo score in endoscopic evaluation of aspiration during swallowing. <i>Auris Nasus Larynx</i> , 2018, 45, 1214-1220.	1.2	37
2	Role of 18F-FDG PET in detecting primary site in the patient with primary unknown carcinoma. <i>European Archives of Oto-Rhino-Laryngology</i> , 2010, 267, 1785-1792.	1.6	31
3	Predictive and Prognostic Value of Metabolic Tumor Volume (MTV) in Patients with Laryngeal Carcinoma Treated by Radiotherapy (RT) / Concurrent Chemoradiotherapy (CCRT). <i>PLoS ONE</i> , 2015, 10, e0117924.	2.5	23
4	Lymph node ratio as a prognostic factor for survival in patients with head and neck squamous cell carcinoma. <i>Auris Nasus Larynx</i> , 2018, 45, 846-853.	1.2	19
5	The applicability of new TNM classification for humanpapilloma virus-related oropharyngeal cancer in the 8th edition of the AJCC/UICC TNM staging system in Japan: A single-centre study. <i>Auris Nasus Larynx</i> , 2018, 45, 558-565.	1.2	18
6	Limitations of PET and PET/CT in detecting upper gastrointestinal synchronous cancer in patients with head and neck carcinoma. <i>European Archives of Oto-Rhino-Laryngology</i> , 2013, 270, 727-733.	1.6	11
7	Efficacy and safety of postoperative bio-chemoradiotherapy using cetuximab and docetaxel for high-risk head and neck cancer patients in Japan. <i>Cancer Chemotherapy and Pharmacology</i> , 2017, 80, 203-207.	2.3	9
8	Surgery-based versus radiation-based treatment strategy for a high metabolic volume laryngeal cancer. <i>Laryngoscope</i> , 2017, 127, 862-867.	2.0	9
9	Prognostic significance of metabolic tumor volume in patients with piriform sinus carcinoma treated by radiotherapy with or without concurrent chemotherapy. <i>Head and Neck</i> , 2016, 38, 1666-1671.	2.0	6
10	Predictive markers, including total lesion glycolysis, for the response of lymph node(s) metastasis from head and neck squamous cell carcinoma treated by chemoradiotherapy. <i>International Journal of Clinical Oncology</i> , 2016, 21, 224-230.	2.2	6
11	The Second-Look Procedure for Transoral Videolaryngoscopic Surgery for T1 and T2 Laryngeal, Oropharyngeal, and Hypopharyngeal Cancer Patients: Protocol for a Nonrandomized Clinical Trial. <i>JMIR Research Protocols</i> , 2017, 6, e235.	1.0	5
12	Postoperative Bio-Chemoradiotherapy Using Cetuximab and Docetaxel in Patients With Cis-Platinum Intolerant Core High-Risk Head and Neck Cancer: Protocol of a Phase 2 Nonrandomized Clinical Trial. <i>JMIR Research Protocols</i> , 2018, 7, e11003.	1.0	3
13	The Association between the Metabolic Tumor Volume and the Prognosis in Patients with Laryngeal Carcinoma. <i>Koutou (the LARYNX JAPAN)</i> , 2016, 28, 57-60.	0.1	3
14	A Case of Intraorbital Abscess Developing as a Complication of Acute Dacryocystitis. <i>Journal of Otolaryngology of Japan</i> , 2017, 120, 722-726.	0.1	0
15	Lymph node ratio as a prognostic factor for survival in patients with head and neck squamous cell carcinoma. <i>Journal of Otolaryngology of Japan</i> , 2019, 122, 1009-1010.	0.1	0
16	A case of adult deep neck abscess caused by <i>Streptococcus pyogenes</i> . <i>Journal of Japan Society for Head and Neck Surgery</i> , 2018, 28, 69-73.	0.0	0