

Peter M Zygmunt

List of Publications by Year in Descending Order

Source: <https://exaly.com/author-pdf/4838648/peter-m-zygmunt-publications-by-year.pdf>

Version: 2024-04-27

This document has been generated based on the publications and citations recorded by exaly.com. For the latest version of this publication list, visit the link given above.

The third column is the impact factor (IF) of the journal, and the fourth column is the number of citations of the article.

40
papers

6,879
citations

27
h-index

42
g-index

42
ext. papers

7,430
ext. citations

9.3
avg, IF

5.21
L-index

#	Paper	IF	Citations
40	Paracetamol analogues conjugated by FAAH induce TRPV1-mediated antinociception without causing acute liver toxicity. <i>European Journal of Medicinal Chemistry</i> , 2021 , 213, 113042	6.8	2
39	Calcium activates purified human TRPA1 with and without its N-terminal ankyrin repeat domain in the absence of calmodulin. <i>Cell Calcium</i> , 2020 , 90, 102228	4	7
38	Human TRPA1 is an inherently mechanosensitive bilayer-gated ion channel. <i>Cell Calcium</i> , 2020 , 91, 102255	16	
37	Electrophile-Induced Conformational Switch of the Human TRPA1 Ion Channel Detected by Mass Spectrometry. <i>International Journal of Molecular Sciences</i> , 2020 , 21,	6.3	4
36	Oxidation of methionine residues activates the high-threshold heat-sensitive ion channel TRPV2. <i>Proceedings of the National Academy of Sciences of the United States of America</i> , 2019 , 116, 24359-24365	11.5	21
35	The antipyretic effect of paracetamol occurs independent of transient receptor potential ankyrin 1-mediated hypothermia and is associated with prostaglandin inhibition in the brain. <i>FASEB Journal</i> , 2018 , 32, 5751-5759	0.9	20
34	TRPA1-FGFR2 binding event is a regulatory oncogenic driver modulated by miRNA-142-3p. <i>Nature Communications</i> , 2017 , 8, 947	17.4	26
33	The N-terminal Ankyrin Repeat Domain Is Not Required for Electrophile and Heat Activation of the Purified Mosquito TRPA1 Receptor. <i>Journal of Biological Chemistry</i> , 2016 , 291, 26899-26912	5.4	13
32	Human TRPA1 is a heat sensor displaying intrinsic U-shaped thermosensitivity. <i>Scientific Reports</i> , 2016 , 6, 28763	4.9	76
31	Photosensitization in Porphyrrias and Photodynamic Therapy Involves TRPA1 and TRPV1. <i>Journal of Neuroscience</i> , 2016 , 36, 5264-78	6.6	52
30	H2S and NO cooperatively regulate vascular tone by activating a neuroendocrine HNO-TRPA1-CGRP signalling pathway. <i>Nature Communications</i> , 2014 , 5, 4381	17.4	267
29	TRPA1. <i>Handbook of Experimental Pharmacology</i> , 2014 , 222, 583-630	3.2	153
28	Human TRPA1 is intrinsically cold- and chemosensitive with and without its N-terminal ankyrin repeat domain. <i>Proceedings of the National Academy of Sciences of the United States of America</i> , 2014 , 111, 16901-6	11.5	108
27	Monoacylglycerols activate TRPV1--a link between phospholipase C and TRPV1. <i>PLoS ONE</i> , 2013 , 8, e81638	3.8	104
26	Fatty acid amide hydrolase-dependent generation of antinociceptive drug metabolites acting on TRPV1 in the brain. <i>PLoS ONE</i> , 2013 , 8, e70690	3.7	44
25	TRPA1 contributes to the acute inflammatory response and mediates carrageenan-induced paw edema in the mouse. <i>Scientific Reports</i> , 2012 , 2, 380	4.9	63
24	TRPA1 mediates spinal antinociception induced by acetaminophen and the cannabinoid (9)-tetrahydrocannabinol. <i>Nature Communications</i> , 2011 , 2, 551	17.4	206

23	TRPV1 and TRPA1 stimulation induces MUC5B secretion in the human nasal airway in vivo. <i>Clinical Physiology and Functional Imaging</i> , 2011 , 31, 435-44	2.4	20
22	TRPV1 in brain is involved in acetaminophen-induced antinociception. <i>PLoS ONE</i> , 2010 , 5, e12748	3.7	105
21	Distribution and function of the hydrogen sulfide-sensitive TRPA1 ion channel in rat urinary bladder. <i>European Urology</i> , 2008 , 53, 391-9	10.2	230
20	Pungent products from garlic activate the sensory ion channel TRPA1. <i>Proceedings of the National Academy of Sciences of the United States of America</i> , 2005 , 102, 12248-52	11.5	645
19	Conversion of acetaminophen to the bioactive N-acylphenolamine AM404 via fatty acid amide hydrolase-dependent arachidonic acid conjugation in the nervous system. <i>Journal of Biological Chemistry</i> , 2005 , 280, 31405-12	5.4	284
18	Vascular effects of anandamide and N-acylvanillylamines in the human forearm and skin microcirculation. <i>British Journal of Pharmacology</i> , 2005 , 146, 171-9	8.6	34
17	Endogenous unsaturated C18 N-acylethanolamines are vanilloid receptor (TRPV1) agonists. <i>Journal of Biological Chemistry</i> , 2005 , 280, 38496-504	5.4	143
16	Mustard oils and cannabinoids excite sensory nerve fibres through the TRP channel ANKTM1. <i>Nature</i> , 2004 , 427, 260-5	50.4	1514
15	Delta 9-tetrahydrocannabinol and cannabinol activate capsaicin-sensitive sensory nerves via a CB1 and CB2 cannabinoid receptor-independent mechanism. <i>Journal of Neuroscience</i> , 2002 , 22, 4720-7	6.6	99
14	Mechanisms underlying tissue selectivity of anandamide and other vanilloid receptor agonists. <i>Molecular Pharmacology</i> , 2002 , 62, 705-13	4.3	45
13	Effects of inhibitors of small- and intermediate-conductance calcium-activated potassium channels, inwardly-rectifying potassium channels and Na(+)/K(+) ATPase on EDHF relaxations in the rat hepatic artery. <i>British Journal of Pharmacology</i> , 2000 , 129, 1490-6	8.6	46
12	Involvement of sensory nerves in vasodilator responses to acetylcholine and potassium ions in rat hepatic artery. <i>British Journal of Pharmacology</i> , 2000 , 130, 27-32	8.6	10
11	The anandamide transport inhibitor AM404 activates vanilloid receptors. <i>European Journal of Pharmacology</i> , 2000 , 396, 39-42	5.3	207
10	Vanilloid receptors on sensory nerves mediate the vasodilator action of anandamide. <i>Nature</i> , 1999 , 400, 452-7	50.4	1815
9	Interactions between endothelium-derived relaxing factors in the rat hepatic artery: focus on regulation of EDHF. <i>British Journal of Pharmacology</i> , 1998 , 124, 992-1000	8.6	39
8	Characterization of endothelium-dependent relaxation in guinea pig basilar artery - effect of hypoxia and role of cytochrome P450 mono-oxygenase. <i>Journal of Vascular Research</i> , 1998 , 35, 285-94	1.9	39
7	Characterization of the potassium channels involved in EDHF-mediated relaxation in cerebral arteries. <i>British Journal of Pharmacology</i> , 1997 , 120, 1344-50	8.6	104
6	Role of potassium channels in endothelium-dependent relaxation resistant to nitroarginine in the rat hepatic artery. <i>British Journal of Pharmacology</i> , 1996 , 117, 1600-6	8.6	176

5	Effects of cytochrome P450 inhibitors on EDHF-mediated relaxation in the rat hepatic artery. <i>British Journal of Pharmacology</i> , 1996 , 118, 1147-52	8.6	73
4	Substance P-induced relaxation and hyperpolarization in human cerebral arteries. <i>British Journal of Pharmacology</i> , 1995 , 115, 889-94	8.6	46
3	Calcium antagonistic properties of the sesquiterpene T-cadinol: a comparison with nimodipine in the isolated rat aorta. <i>Basic and Clinical Pharmacology and Toxicology</i> , 1991 , 69, 173-7		15
2	Human TRPA1 is an inherently mechanosensitive bilayer-gated ion channel		3
1	Calcium activates purified human TRPA1 with and without its N-terminal ankyrin repeat domain in the absence of calmodulin		1