

Giuseppe M Campo

List of Publications by Year in Descending Order

Source: <https://exaly.com/author-pdf/4840409/giuseppe-m-campo-publications-by-year.pdf>

Version: 2024-04-27

This document has been generated based on the publications and citations recorded by exaly.com. For the latest version of this publication list, visit the link given above.

The third column is the impact factor (IF) of the journal, and the fourth column is the number of citations of the article.

161
papers

4,839
citations

38
h-index

60
g-index

170
ext. papers

5,182
ext. citations

5.2
avg, IF

4.67
L-index

#	Paper	IF	Citations
161	miR9 inhibits 6-mer HA-induced cytokine production and apoptosis in human chondrocytes by reducing NF-kB activation.. <i>Archives of Biochemistry and Biophysics</i> , 2022 , 718, 109139	4.1	0
160	miR146a up-regulation is involved in small HA oligosaccharides-induced pro-inflammatory response in human chondrocytes. <i>Biochimica Et Biophysica Acta - General Subjects</i> , 2021 , 1865, 129731	4	3
159	Endocan, a novel inflammatory marker, is upregulated in human chondrocytes stimulated with IL-1 beta. <i>Molecular and Cellular Biochemistry</i> , 2021 , 476, 1589-1597	4.2	2
158	Hyaluronan Fragmentation During Inflammatory Pathologies: A Signal that Empowers Tissue Damage. <i>Mini-Reviews in Medicinal Chemistry</i> , 2020 , 20, 54-65	3.2	9
157	Biglycan and atherosclerosis: Lessons from high cardiovascular risk conditions. <i>Biochimica Et Biophysica Acta - Molecular and Cell Biology of Lipids</i> , 2020 , 1865, 158545	5	13
156	Hyaluronan oligosaccharides modulate inflammatory response, NIS and thyroglobulin expression in human thyrocytes. <i>Archives of Biochemistry and Biophysics</i> , 2020 , 694, 108598	4.1	5
155	Hyaluronan fragments produced during tissue injury: A signal amplifying the inflammatory response. <i>Archives of Biochemistry and Biophysics</i> , 2019 , 663, 228-238	4.1	15
154	Serglycin as part of IL-1 β induced inflammation in human chondrocytes. <i>Archives of Biochemistry and Biophysics</i> , 2019 , 669, 80-86	4.1	8
153	ECaryophyllene Mitigates Collagen Antibody Induced Arthritis (CAIA) in Mice Through a Cross-Talk between CB2 and PPAR- γ Receptors. <i>Biomolecules</i> , 2019 , 9,	5.9	29
152	Exploiting Curcumin Synergy With Natural Products Using Quantitative Analysis of Dose-Effect Relationships in an Experimental Model of Osteoarthritis. <i>Frontiers in Pharmacology</i> , 2019 , 10, 1347	5.6	13
151	The proteoglycan biglycan mediates inflammatory response by activating TLR-4 in human chondrocytes: Inhibition by specific siRNA and high polymerized Hyaluronan. <i>Archives of Biochemistry and Biophysics</i> , 2018 , 640, 75-82	4.1	12
150	Serglycin is involved in inflammatory response in articular mouse chondrocytes. <i>Biochemical and Biophysical Research Communications</i> , 2018 , 499, 506-512	3.4	11
149	Endothelial progenitor cells and rheumatic disease modifying therapy. <i>Vascular Pharmacology</i> , 2018 , 108, 8-14	5.9	5
148	Hyaluronan in experimental injured/inflamed cartilage: In vivo studies. <i>Life Sciences</i> , 2018 , 193, 132-140	6.8	19
147	Hyaluronan in the experimental injury of the cartilage: biochemical action and protective effects. <i>Inflammation Research</i> , 2018 , 67, 5-20	7.2	21
146	6-Mer Hyaluronan Oligosaccharides Modulate Neuroinflammation and β Synuclein Expression in Neuron-Like SH-SY5Y Cells. <i>Journal of Cellular Biochemistry</i> , 2016 , 117, 2835-2843	4.7	15
145	Inhibition of small HA fragment activity and stimulation of A2A adenosine receptor pathway limit apoptosis and reduce cartilage damage in experimental arthritis. <i>Histochemistry and Cell Biology</i> , 2015 , 143, 531-43	2.4	20

144	Beta-arrestin 1 is involved in the catabolic response stimulated by hyaluronan degradation in mouse chondrocytes. <i>Cell and Tissue Research</i> , 2015 , 361, 567-79	4.2	6
143	Beta-arrestin-2 negatively modulates inflammation response in mouse chondrocytes induced by 4-mer hyaluronan oligosaccharide. <i>Molecular and Cellular Biochemistry</i> , 2015 , 399, 201-8	4.2	17
142	Evaluation of putative cytotoxic activity of crude extracts from <i>Onopordum acanthium</i> leaves and <i>Spartium junceum</i> flowers against the U-373 glioblastoma cell line. <i>Pakistan Journal of Pharmaceutical Sciences</i> , 2015 , 28, 1225-32	0.4	6
141	Inhibition of the hyaluronan oligosaccharides inflammatory response: reduction of adenosine 2A receptor activation by EPAC and PKA. <i>Cell Biochemistry and Function</i> , 2014 , 32, 692-701	4.2	4
140	MiRNome expression is deregulated in the peripheral lymphoid compartment of multiple myeloma. <i>British Journal of Haematology</i> , 2014 , 165, 801-13	4.5	16
139	The SOD mimic MnTM-2-PyP(5+) reduces hyaluronan degradation-induced inflammation in mouse articular chondrocytes stimulated with Fe (II) plus ascorbate. <i>International Journal of Biochemistry and Cell Biology</i> , 2013 , 45, 1610-9	5.6	17
138	Combined treatment with hyaluronan inhibitor Pep-1 and a selective adenosine A2 receptor agonist reduces inflammation in experimental arthritis. <i>Innate Immunity</i> , 2013 , 19, 462-78	2.7	11
137	4-mer hyaluronan oligosaccharides stimulate inflammation response in synovial fibroblasts in part via TAK-1 and in part via p38-MAPK. <i>Current Medicinal Chemistry</i> , 2013 , 20, 1162-72	4.3	27
136	Adenosine A2A receptor activation and hyaluronan fragment inhibition reduce inflammation in mouse articular chondrocytes stimulated with interleukin-1. <i>FEBS Journal</i> , 2012 , 279, 2120-33	5.7	36
135	Hyaluronan differently modulates TLR-4 and the inflammatory response in mouse chondrocytes. <i>BioFactors</i> , 2012 , 38, 69-76	6.1	59
134	Protein kinase a mediated anti-inflammatory effects exerted by adenosine treatment in mouse chondrocytes stimulated with IL-1. <i>BioFactors</i> , 2012 , 38, 429-39	6.1	9
133	Inhibition of hyaluronan synthesis reduced inflammatory response in mouse synovial fibroblasts subjected to collagen-induced arthritis. <i>Archives of Biochemistry and Biophysics</i> , 2012 , 518, 42-52	4.1	25
132	Hyaluronan in part mediates IL-1beta-induced inflammation in mouse chondrocytes by up-regulating CD44 receptors. <i>Gene</i> , 2012 , 494, 24-35	3.8	36
131	The stimulation of adenosine 2A receptor reduces inflammatory response in mouse articular chondrocytes treated with hyaluronan oligosaccharides. <i>Matrix Biology</i> , 2012 , 31, 338-51	11.4	21
130	The inhibition of hyaluronan degradation reduced pro-inflammatory cytokines in mouse synovial fibroblasts subjected to collagen-induced arthritis. <i>Journal of Cellular Biochemistry</i> , 2012 , 113, 1852-67	4.7	53
129	6-Mer hyaluronan oligosaccharides increase IL-18 and IL-33 production in mouse synovial fibroblasts subjected to collagen-induced arthritis. <i>Innate Immunity</i> , 2012 , 18, 675-84	2.7	20
128	Systemic administration of high-molecular weight hyaluronan stimulates wound healing in genetically diabetic mice. <i>Biochimica Et Biophysica Acta - Molecular Basis of Disease</i> , 2011 , 1812, 752-9	6.9	45
127	Hyaluronan reduces inflammation in experimental arthritis by modulating TLR-2 and TLR-4 cartilage expression. <i>Biochimica Et Biophysica Acta - Molecular Basis of Disease</i> , 2011 , 1812, 1170-81	6.9	83

126	Polydeoxyribonucleotide reduces cytokine production and the severity of collagen-induced arthritis by stimulation of adenosine A _{2A} receptor. <i>Arthritis and Rheumatism</i> , 2011 , 63, 3364-71		65
125	Hyaluronan reduces inflammation in experimental arthritis by modulating TLR-2 and TLR-4 cartilage expression. <i>Biochimica Et Biophysica Acta - Molecular Basis of Disease</i> , 2011 , 1812, 1170-1181	6.9	1
124	Differential effect of molecular mass hyaluronan on lipopolysaccharide-induced damage in chondrocytes. <i>Innate Immunity</i> , 2010 , 16, 48-63	2.7	24
123	Molecular cloning and characterization of adult <i>Sparus aurata</i> hemoglobin genes. <i>OMICS A Journal of Integrative Biology</i> , 2010 , 14, 187-200	3.8	2
122	Molecular size hyaluronan differently modulates toll-like receptor-4 in LPS-induced inflammation in mouse chondrocytes. <i>Biochimie</i> , 2010 , 92, 204-15	4.6	121
121	High-molecular weight hyaluronan reduced renal PKC activation in genetically diabetic mice. <i>Biochimica Et Biophysica Acta - Molecular Basis of Disease</i> , 2010 , 1802, 1118-30	6.9	19
120	Small hyaluronan oligosaccharides induce inflammation by engaging both toll-like-4 and CD44 receptors in human chondrocytes. <i>Biochemical Pharmacology</i> , 2010 , 80, 480-90	6	115
119	Glycosaminoglycans modulate inflammation and apoptosis in LPS-treated chondrocytes. <i>Journal of Cellular Biochemistry</i> , 2009 , 106, 83-92	4.7	70
118	Differential effect of molecular size HA in mouse chondrocytes stimulated with PMA. <i>Biochimica Et Biophysica Acta - General Subjects</i> , 2009 , 1790, 1353-67	4	43
117	Glycosaminoglycans reduced inflammatory response by modulating toll-like receptor-4 in LPS-stimulated chondrocytes. <i>Archives of Biochemistry and Biophysics</i> , 2009 , 491, 7-15	4.1	47
116	Effect of cytokines on hyaluronan synthase activity and response to oxidative stress by fibroblasts. <i>British Journal of Biomedical Science</i> , 2009 , 66, 28-36	1.6	11
115	The antioxidant activity of chondroitin-4-sulphate, in carbon tetrachloride-induced acute hepatitis in mice, involves NF-kappaB and caspase activation. <i>British Journal of Pharmacology</i> , 2008 , 155, 945-56	8.6	45
114	Chondroitin-4-sulphate inhibits NF-kB translocation and caspase activation in collagen-induced arthritis in mice. <i>Osteoarthritis and Cartilage</i> , 2008 , 16, 1474-83	6.2	38
113	Hemoglobin system of <i>Sparus aurata</i> : changes in fishes farmed under extreme conditions. <i>Science of the Total Environment</i> , 2008 , 403, 148-53	10.2	19
112	Chondroitin-4-sulphate reduced oxidative injury in caerulein-induced pancreatitis in mice: the involvement of NF-kappaB translocation and apoptosis activation. <i>Experimental Biology and Medicine</i> , 2008 , 233, 741-52	3.7	14
111	Transient increase with strenuous exercise of plasma levels of glycosaminoglycans in humans and horses. <i>Connective Tissue Research</i> , 2008 , 49, 416-25	3.3	3
110	Purified human plasma glycosaminoglycans reduced NF-kappaB activation, pro-inflammatory cytokine production and apoptosis in LPS-treated chondrocytes. <i>Innate Immunity</i> , 2008 , 14, 233-46	2.7	19
109	The antioxidant effect exerted by TGF-1beta-stimulated hyaluronan production reduced NF-kB activation and apoptosis in human fibroblasts exposed to FeSo ₄ plus ascorbate. <i>Molecular and Cellular Biochemistry</i> , 2008 , 311, 167-77	4.2	16

108	NF-kB and caspases are involved in the hyaluronan and chondroitin-4-sulphate-exerted antioxidant effect in fibroblast cultures exposed to oxidative stress. <i>Journal of Applied Toxicology</i> , 2008 , 28, 509-17	4.1	30
107	Identification and gene expression of versican during early development of <i>Xenopus</i> . <i>International Journal of Developmental Biology</i> , 2008 , 52, 993-8	1.9	11
106	Differential effect of growth factors on hyaluronan synthase gene expression in fibroblasts exposed to oxidative stress. <i>Biochemistry (Moscow)</i> , 2007 , 72, 974-82, 4 p.	2.9	2
105	Chondroitin sulphate: antioxidant properties and beneficial effects. <i>Mini-Reviews in Medicinal Chemistry</i> , 2006 , 6, 1311-20	3.2	26
104	Lymphocytes from patients with early stage of B-cell chronic lymphocytic leukaemia and long survival synthesize decorin. <i>Biochimie</i> , 2006 , 88, 1933-9	4.6	9
103	Antioxidant activity of chondroitin sulfate. <i>Advances in Pharmacology</i> , 2006 , 53, 417-31	5.7	26
102	TNF-alpha, IFN-gamma, and IL-1beta modulate hyaluronan synthase expression in human skin fibroblasts: synergistic effect by concomital treatment with FeSO4 plus ascorbate. <i>Molecular and Cellular Biochemistry</i> , 2006 , 292, 169-78	4.2	35
101	Purified human chondroitin-4-sulfate reduced MMP/TIMP imbalance induced by iron plus ascorbate in human fibroblast cultures. <i>Cell Biology International</i> , 2006 , 30, 21-30	4.5	14
100	Purified human plasma glycosaminoglycans limit oxidative injury induced by iron plus ascorbate in skin fibroblast cultures. <i>Toxicology in Vitro</i> , 2005 , 19, 561-72	3.6	24
99	Extracellular superoxide dismutase (EC-SOD) gene mutations screening in a sample of Mediterranean population. <i>Mutation Research - Fundamental and Molecular Mechanisms of Mutagenesis</i> , 2005 , 578, 143-8	3.3	14
98	Oxidative stress in myotonic dystrophy type 1. <i>Free Radical Research</i> , 2005 , 39, 771-6	4	37
97	Effects of AT1 receptor antagonist losartan on sICAM-1 and TNF-alpha levels in uncomplicated hypertensive patients. <i>Angiology</i> , 2004 , 55, 195-203	2.1	15
96	Identification of paraoxonase 3 gene (PON3) missense mutations in a population of southern Italy. <i>Mutation Research - Fundamental and Molecular Mechanisms of Mutagenesis</i> , 2004 , 546, 75-80	3.3	25
95	The antioxidant and antifibrogenic effects of the glycosaminoglycans hyaluronic acid and chondroitin-4-sulphate in a subchronic rat model of carbon tetrachloride-induced liver fibrogenesis. <i>Chemico-Biological Interactions</i> , 2004 , 148, 125-38	5	46
94	Reduction of DNA fragmentation and hydroxyl radical production by hyaluronic acid and chondroitin-4-sulphate in iron plus ascorbate-induced oxidative stress in fibroblast cultures. <i>Free Radical Research</i> , 2004 , 38, 601-11	4	30
93	Glycosaminoglycans reduce oxidative damage induced by copper (Cu ²⁺), iron (Fe ²⁺) and hydrogen peroxide (H ₂ O ₂) in human fibroblast cultures. <i>Glycoconjugate Journal</i> , 2004 , 20, 133-41	3	38
92	Levetiracetam protects against kainic acid-induced toxicity. <i>Life Sciences</i> , 2004 , 74, 1253-64	6.8	49
91	Hyaluronic acid and chondroitin-4-sulphate treatment reduces damage in carbon tetrachloride-induced acute rat liver injury. <i>Life Sciences</i> , 2004 , 74, 1289-305	6.8	45

90	A protective effect of the synthetic coumarine derivative Cloricromene against DNB-colitis in the rat. <i>Life Sciences</i> , 2004 , 74, 2749-56	6.8	3
89	Administration of hyaluronic acid and chondroitin-4-sulfate limits endogenous antioxidant depletion and reduces cell damage in experimental acute pancreatitis. <i>Pancreas</i> , 2004 , 28, E45-53	2.6	13
88	Characterization of serum glycosaminoglycan (GAG) chains and native proteoglycan fractions in the ostrich. <i>Veterinary Research Communications</i> , 2003 , 27 Suppl 1, 599-601	2.9	2
87	Efficacy of treatment with glycosaminoglycans on experimental collagen-induced arthritis in rats. <i>Arthritis Research</i> , 2003 , 5, R122-31		139
86	Lipid peroxidation inhibition reduces NF-kappaB activation and attenuates cerulein-induced pancreatitis. <i>Free Radical Research</i> , 2003 , 37, 425-35	4	25
85	Aromatic trap analysis of free radicals production in experimental collagen-induced arthritis in the rat: protective effect of glycosaminoglycans treatment. <i>Free Radical Research</i> , 2003 , 37, 257-68	4	39
84	The effect of the phytoestrogen genistein on plasma nitric oxide concentrations, endothelin-1 levels and endothelium dependent vasodilation in postmenopausal women. <i>Atherosclerosis</i> , 2002 , 163, 339-47	3.1	191
83	Improved high-performance liquid chromatographic method to estimate aminosugars and its application to glycosaminoglycan determination in plasma and serum. <i>Biomedical Applications</i> , 2001 , 765, 151-60		33
82	Reduction of carbon tetrachloride-induced rat liver injury by IRFI 042, a novel dual vitamin E-like antioxidant. <i>Free Radical Research</i> , 2001 , 34, 379-93	4	61
81	Oxidative stress causes nuclear factor-kappaB activation in acute hypovolemic hemorrhagic shock. <i>Free Radical Biology and Medicine</i> , 2001 , 30, 1055-66	7.8	62
80	Effects of simvastatin treatment on sICAM-1 and sE-selectin levels in hypercholesterolemic subjects. <i>Atherosclerosis</i> , 2001 , 155, 143-7	3.1	28
79	Raxofelast, a hydrophilic vitamin E-like antioxidant, stimulates wound healing in genetically diabetic mice. <i>Surgery</i> , 2001 , 129, 467-77	3.6	24
78	Inhibition of lipid peroxidation restores impaired vascular endothelial growth factor expression and stimulates wound healing and angiogenesis in the genetically diabetic mouse. <i>Diabetes</i> , 2001 , 50, 667-74	0.9	197
77	Protective effects of cyclosporin-A in splanchnic artery occlusion shock. <i>British Journal of Pharmacology</i> , 2000 , 130, 339-44	8.6	8
76	Genistein supplementation and estrogen replacement therapy improve endothelial dysfunction induced by ovariectomy in rats. <i>Cardiovascular Research</i> , 2000 , 45, 454-62	9.9	117
75	IRFI 042, a novel dual vitamin E-like antioxidant, inhibits activation of nuclear factor-kappaB and reduces the inflammatory response in myocardial ischemia-reperfusion injury. <i>Cardiovascular Research</i> , 2000 , 47, 515-28	9.9	56
74	The reduction of myocardial damage and leukocyte polymorphonuclear accumulation following coronary artery occlusion by the tyrosine kinase inhibitor tyrphostin AG 556. <i>Life Sciences</i> , 2000 , 67, 2615-29	6.8	6
73	Effects of <i>Hypericum perforatum</i> on levels of 5-hydroxytryptamine, noradrenaline and dopamine in the cortex, diencephalon and brainstem of the rat. <i>Journal of Pharmacy and Pharmacology</i> , 1999 , 51, 723-8	4.8	63

72	Recombinant human erythropoietin inhibits iNOS activity and reverts vascular dysfunction in splanchnic artery occlusion shock. <i>British Journal of Pharmacology</i> , 1999 , 127, 482-8	8.6	56
71	Tacrolimus suppresses tumour necrosis factor-alpha and protects against splanchnic artery occlusion shock. <i>British Journal of Pharmacology</i> , 1999 , 127, 498-504	8.6	13
70	Adrenocorticotropin reverses vascular dysfunction and protects against splanchnic artery occlusion shock. <i>British Journal of Pharmacology</i> , 1999 , 128, 816-22	8.6	26
69	Cardioprotection by the phytoestrogen genistein in experimental myocardial ischaemia-reperfusion injury. <i>British Journal of Pharmacology</i> , 1999 , 128, 1683-90	8.6	71
68	Cyclosporin-A reduces leukocyte accumulation and protects against myocardial ischaemia reperfusion injury in rats. <i>European Journal of Pharmacology</i> , 1999 , 364, 159-68	5.3	43
67	The lazaroïd, U-74389G, inhibits inducible nitric oxide synthase activity, reverses vascular failure and protects against endotoxin shock. <i>European Journal of Pharmacology</i> , 1999 , 369, 49-55	5.3	19
66	Effect of sulfatide on acute lung injury during endotoxemia in rats. <i>Life Sciences</i> , 1999 , 65, 2541-52	6.8	8
65	Leptin increases serotonin turnover by inhibition of brain nitric oxide synthesis. <i>Journal of Clinical Investigation</i> , 1999 , 104, 975-82	15.9	128
64	Inhibition of tumour necrosis factor and reversal of endotoxin-induced shock by U-83836E, a second generation lazaroïd in rats. <i>British Journal of Pharmacology</i> , 1998 , 124, 1293-9	8.6	12
63	Sulfatide reduces leucocyte accumulation and reverts vascular failure in splanchnic artery occlusion shock. <i>European Journal of Pharmacology</i> , 1998 , 361, 101-8	5.3	2
62	Determination of clozapine, desmethylclozapine and clozapine N-oxide in human plasma by reversed-phase high-performance liquid chromatography with ultraviolet detection. <i>Biomedical Applications</i> , 1998 , 714, 299-308		35
61	Beneficial effect of raxofelast, an hydrophilic vitamin E analogue, in the rat heart after ischemia and reperfusion injury. <i>Journal of Molecular and Cellular Cardiology</i> , 1998 , 30, 1493-503	5.8	27
60	Effects of picotamide on release of endothelin-1, thromboxane and prostacycline after treadmill stress in patients with peripheral artery disease. <i>Angiology</i> , 1998 , 49, 879-84	2.1	5
59	Protective effects of the new lazaroïd "U-83836E" in splanchnic artery occlusion (SAO) shock. <i>Free Radical Research</i> , 1998 , 28, 477-84	4	5
58	Multiple organ failure following zymosan-induced peritonitis is mediated by nitric oxide. <i>Shock</i> , 1997 , 8, 268-75	3.4	53
57	Antioxidant activity of U-83836E, a second generation lazaroïd, during myocardial ischemia/reperfusion injury. <i>Free Radical Research</i> , 1997 , 27, 577-90	4	21
56	17Beta-oestradiol reduces cardiac leukocyte accumulation in myocardial ischaemia reperfusion injury in rat. <i>European Journal of Pharmacology</i> , 1997 , 335, 185-92	5.3	81
55	Raxofelast (IRFI 016): A New Hydrophilic Vitamin E-Like Antioxidant Agent. <i>Cardiovascular Drug Reviews</i> , 1997 , 15, 157-173		13

54	The effects of recombinant human granulocyte-colony stimulating factor on vascular dysfunction and splanchnic ischaemia-reperfusion injury. <i>British Journal of Pharmacology</i> , 1997 , 120, 333-9	8.6	23
53	The involvement of tumour necrosis factor-alpha in the protective effects of 17 beta oestradiol in splanchnic ischaemia-reperfusion injury. <i>British Journal of Pharmacology</i> , 1997 , 121, 1782-8	8.6	19
52	Relationship between plasma desipramine levels, CYP2D6 phenotype and clinical response to desipramine: a prospective study. <i>European Journal of Clinical Pharmacology</i> , 1997 , 51, 395-8	2.8	93
51	Effect of ketoconazole on the pharmacokinetics of imipramine and desipramine in healthy subjects. <i>British Journal of Clinical Pharmacology</i> , 1997 , 43, 315-8	3.8	34
50	Effects of S-ethylisothiourea, a potent inhibitor of nitric oxide synthase, alone or in combination with a nitric oxide donor in splanchnic artery occlusion shock. <i>British Journal of Pharmacology</i> , 1996 , 119, 23-8	8.6	8
49	Endotoxin tolerance impairs a pertussis-toxin-sensitive G-protein regulating tumour necrosis factor release by macrophages from tumour-bearing rats. <i>Pharmacological Research</i> , 1996 , 33, 203-9	10.2	5
48	Leukocyte integrin very late antigen-4/vascular cell adhesion molecule-1 adhesion pathway in splanchnic artery occlusion shock. <i>European Journal of Pharmacology</i> , 1996 , 318, 153-60	5.3	3
47	Effects of cloricromene on the levels of endothelin and on the microcirculatory function in peripheral atherosclerotic arteriopathies. <i>Pharmacology</i> , 1996 , 52, 8-15	2.3	
46	Monocytes and lymphocytes as active participants in the pathogenesis of experimental shock. <i>Inflammation Research</i> , 1996 , 45, 398-404	7.2	7
45	Thrombolytic therapy with urokinase reduces increased circulating endothelial adhesion molecules in acute myocardial infarction. <i>Inflammation Research</i> , 1996 , 45, 14-9	7.2	37
44	Phenobarbital induces the 2-hydroxylation of desipramine. <i>Therapeutic Drug Monitoring</i> , 1996 , 18, 60-4	3.2	30
43	The effect of carbamazepine on the 2-hydroxylation of desipramine. <i>Psychopharmacology</i> , 1995 , 117, 413-6	4.7	24
42	Multiple actions of the coumarine derivative cloricromene and its protective effects on ischemic brain injury. <i>Naunyn-Schmiedeberg's Archives of Pharmacology</i> , 1995 , 351, 209-15	3.4	14
41	Soluble E-selectin levels in acute human myocardial infarction. <i>International Journal of Microcirculation, Clinical and Experimental</i> , 1995 , 15, 80-4		2
40	E-selectin involvement in the pathogenesis of splanchnic artery occlusion shock. <i>European Journal of Pharmacology</i> , 1995 , 272, 223-9	5.3	7
39	G 619, a dual thromboxane synthase inhibitor and thromboxane A2 receptor antagonist, inhibits tumor necrosis factor-alpha biosynthesis. <i>European Journal of Pharmacology</i> , 1995 , 286, 31-9	5.3	13
38	Effects of gallopamil on epinephrine and norepinephrine plasmatic levels and on TxB2 and beta-tg release in patients with coronary artery disease during adrenergic stimulus with cold pressor test. <i>Pharmacological Research</i> , 1995 , 32, 49-55	10.2	
37	Tumour necrosis factor mediates E-selectin production and leukocyte accumulation in myocardial ischaemia-reperfusion injury. <i>Pharmacological Research</i> , 1995 , 31, 281-8	10.2	18

36	Improved survival and reversal of endothelial dysfunction by the 21-aminosteroid, U-74389G in splanchnic ischaemia-reperfusion injury in the rat. <i>British Journal of Pharmacology</i> , 1995 , 115, 395-400	8.6	25
35	Protective effects of IRFI-016, a new antioxidant agent, in myocardial damage, following coronary artery occlusion and reperfusion in the rat. <i>Pharmacology</i> , 1994 , 48, 157-66	2.3	12
34	Protective effects of L-659,989, a platelet-activating factor receptor antagonist, in myocardial ischemia and reperfusion in rats. <i>Journal of Cardiovascular Pharmacology</i> , 1994 , 23, 7-12	3.1	12
33	TCV-309, a novel platelet activating factor antagonist, inhibits leukocyte accumulation and protects against splanchnic artery occlusion shock. <i>Agents and Actions</i> , 1994 , 42, 128-34		8
32	Reduction of myocardial infarct size in rat by IRFI-048, a selective analogue of vitamin E. <i>Free Radical Biology and Medicine</i> , 1994 , 16, 427-35	7.8	10
31	CYP2D6-related oxidation polymorphism in Italy. <i>Pharmacological Research</i> , 1994 , 29, 281-9	10.2	16
30	Antibodies against intercellular adhesion molecule 1 protect against myocardial ischaemia-reperfusion injury in rat. <i>European Journal of Pharmacology</i> , 1994 , 264, 143-9	5.3	31
29	Central serotonergic system involvement in the anorexia induced by NG-nitro-L-arginine, an inhibitor of nitric oxide synthase. <i>European Journal of Pharmacology</i> , 1994 , 255, 51-5	5.3	52
28	Food deprivation increases brain nitric oxide synthase and depresses brain serotonin levels in rats. <i>Neuropharmacology</i> , 1994 , 33, 83-6	5.5	84
27	Contribution of intercellular adhesion molecule 1 (ICAM-1) to the pathogenesis of splanchnic artery occlusion shock in the rat. <i>British Journal of Pharmacology</i> , 1994 , 113, 912-6	8.6	17
26	Participation of tumour necrosis factor and nitric oxide in the mediation of vascular dysfunction in splanchnic artery occlusion shock. <i>British Journal of Pharmacology</i> , 1994 , 113, 1153-8	8.6	29
25	E-selectin in the pathogenesis of experimental myocardial ischemia-reperfusion injury. <i>European Journal of Pharmacology - Environmental Toxicology and Pharmacology Section</i> , 1994 , 270, 45-51		12
24	Beneficial effects of BAY u3405, a novel thromboxane A2 receptor antagonist, in splanchnic artery occlusion shock. <i>Pharmacology</i> , 1994 , 49, 376-85	2.3	3
23	Decreased plasma concentrations of imipramine and desipramine following cholestyramine intake in depressed patients. <i>Therapeutic Drug Monitoring</i> , 1994 , 16, 432-4	3.2	2
22	Tumor necrosis factor involvement in myocardial ischaemia-reperfusion injury. <i>European Journal of Pharmacology</i> , 1993 , 237, 223-30	5.3	51
21	Anorectic activity of NG-nitro-L-arginine, an inhibitor of brain nitric oxide synthase, in obese Zucker rats. <i>European Journal of Pharmacology</i> , 1993 , 230, 125-8	5.3	82
20	The effect of cloricromene, a coumarine derivative, on leukocyte accumulation, myocardial necrosis and TNF-alpha production in myocardial ischaemia-reperfusion injury. <i>Life Sciences</i> , 1993 , 53, 341-55	6.8	19
19	Splanchnic artery occlusion shock: vinblastine-induced leukopenia reduces tumour necrosis factor and thromboxane A2 formation, and increases survival rate. <i>Pharmacological Research</i> , 1993 , 27, 61-71	10.2	10

18	Effect of fluvoxamine on the pharmacokinetics of imipramine and desipramine in healthy subjects. <i>Therapeutic Drug Monitoring</i> , 1993 , 15, 243-6	3.2	126
17	G619, a dual thromboxane synthase inhibitor and thromboxane A2 receptor antagonist, reduces myocardial damage and polymorphonuclear leukocyte accumulation following coronary artery occlusion and reperfusion in rats. <i>Pharmacology</i> , 1993 , 47, 167-75	2.3	7
16	G 619: A Novel Dual Thromboxane Synthase Inhibitor and Thromboxane A2 Receptor Antagonist. <i>Cardiovascular Drug Reviews</i> , 1993 , 11, 116-125		
15	Reduction of myocardial leukocyte accumulation and myocardial infarct size following administration of BAY u3405, a thromboxane A2 receptor antagonist, in myocardial ischaemia-reperfusion injury. <i>Agents and Actions</i> , 1993 , 39, 143-9		6
14	Protective effects of G 619, a dual thromboxane synthase inhibitor and thromboxane A2 receptor antagonist, in splanchnic artery occlusion shock. <i>Journal of Cardiovascular Pharmacology</i> , 1992 , 19, 115-9 ^{3.1}		13
13	Evidence for a role of nitric oxide in hypovolemic hemorrhagic shock. <i>Journal of Cardiovascular Pharmacology</i> , 1992 , 19, 982-6	3.1	63
12	Interaction between fluvoxamine and imipramine/desipramine in four patients. <i>Therapeutic Drug Monitoring</i> , 1992 , 14, 194-6	3.2	70
11	Debrisoquine oxidation in an Italian population: a study in healthy subjects and in schizophrenic patients. <i>Pharmacological Research</i> , 1992 , 25, 43-50	10.2	12
10	Protective effects of endotoxin tolerance in tumor bearing rats. <i>Pharmacological Research</i> , 1992 , 26 Suppl 2, 162-3	10.2	
9	Evidence that nitric oxide modulates drinking behaviour. <i>Neuropharmacology</i> , 1992 , 31, 761-4	5.5	79
8	Platelet activating factor interaction with tumor necrosis factor and myocardial depressant factor in splanchnic artery occlusion shock. <i>European Journal of Pharmacology</i> , 1992 , 222, 13-9	5.3	33
7	Cloricromene, a coumarine derivative, protects against lethal endotoxin shock in rats. <i>European Journal of Pharmacology</i> , 1992 , 210, 107-13	5.3	25
6	Protective effect of cloricromene, a coumarine derivative, in hypovolemic hemorrhagic shock in the rat. <i>Journal of Cardiovascular Pharmacology</i> , 1991 , 17, 261-6	3.1	19
5	Changes in urine volume and urinary electrolyte excretion after intracerebroventricular injection of arecoline and hemicholinium-3. <i>Life Sciences</i> , 1991 , 48, 2097-107	6.8	3
4	Platelet activating factor involvement in splanchnic artery occlusion shock in rats. <i>European Journal of Pharmacology</i> , 1991 , 192, 47-53	5.3	23
3	Antihypertensive activity of indolepyruvic acid: a keto analogue of tryptophan. <i>Journal of Cardiovascular Pharmacology</i> , 1990 , 15, 102-8	3.1	6
2	Effects of fructose 1,6-diphosphate on splanchnic artery occlusion shock in the rat. <i>Resuscitation</i> , 1989 , 18, 299-307	4	16
1	Effects of fructose-1,6-diphosphate (FDP) in splanchnic artery occlusion (SAO) shock in the rat. <i>Pharmacological Research Communications</i> , 1988 , 20, 371		1

