

Yvonne Forsell

List of Publications by Year in descending order

Source: <https://exaly.com/author-pdf/4846061/publications.pdf>

Version: 2024-02-01

86
papers

2,208
citations

236612

25
h-index

264894

42
g-index

87
all docs

87
docs citations

87
times ranked

3497
citing authors

#	ARTICLE	IF	CITATIONS
1	Internet-Based Cognitive Behavioral Therapy for Depression. <i>JAMA Psychiatry</i> , 2021, 78, 361.	6.0	398
2	Physical exercise and internet-based cognitive-behavioural therapy in the treatment of depression: Randomised controlled trial. <i>British Journal of Psychiatry</i> , 2015, 207, 227-234.	1.7	118
3	Physical Activity Patterns of People Affected by Depressive and Anxiety Disorders as Measured by Accelerometers: A Cross-Sectional Study. <i>PLoS ONE</i> , 2015, 10, e0115894.	1.1	91
4	Cannabis use, depression and anxiety: A 3-year prospective population-based study. <i>Journal of Affective Disorders</i> , 2016, 193, 103-108.	2.0	78
5	Psychotropic drug use in elderly people with and without dementia. <i>International Journal of Geriatric Psychiatry</i> , 2001, 16, 900-906.	1.3	72
6	Exercise, Physical Activity, and Sedentary Behavior in the Treatment of Depression: Broadening the Scientific Perspectives and Clinical Opportunities. <i>Frontiers in Psychiatry</i> , 2016, 7, 36.	1.3	71
7	Morbidity and Comorbidity in Relation to Functional Status: A Community-Based Study of the Oldest Old (90+ Years). <i>Journal of the American Geriatrics Society</i> , 2000, 48, 1462-1469.	1.3	63
8	Training fast or slow? Exercise for depression: A randomized controlled trial. <i>Preventive Medicine</i> , 2016, 91, 123-131.	1.6	60
9	Early exposure to antibiotic drugs and risk for psychiatric disorders: a population-based study. <i>Translational Psychiatry</i> , 2019, 9, 317.	2.4	60
10	A three-year follow-up of major depression, dysthymia, minor depression and subsyndromal depression: results from a population-based study. <i>Depression and Anxiety</i> , 2007, 24, 62-65.	2.0	58
11	Exercise and internet-based cognitive-behavioural therapy for depression: multicentre randomised controlled trial with 12-month follow-up. <i>British Journal of Psychiatry</i> , 2016, 209, 414-420.	1.7	57
12	Interleukin-6 and depressive symptom severity in response to physical exercise. <i>Psychiatry Research</i> , 2017, 252, 270-276.	1.7	49
13	Somebody to lean on: Social relationships predict post-treatment depression severity in adults. <i>Psychiatry Research</i> , 2017, 249, 261-267.	1.7	46
14	Long-term effects of exercise at different intensity levels on depression: A randomized controlled trial. <i>Preventive Medicine</i> , 2017, 105, 37-46.	1.6	46
15	NR3C1 hypermethylation in depressed and bullied adolescents. <i>Translational Psychiatry</i> , 2018, 8, 121.	2.4	46
16	Common Perceived Barriers and Facilitators for Reducing Sedentary Behaviour among Office Workers. <i>International Journal of Environmental Research and Public Health</i> , 2018, 15, 792.	1.2	38
17	Hypomethylation of MAOA ^{x3} 's first exon region in depression: A replication study. <i>Psychiatry Research</i> , 2015, 226, 389-391.	1.7	37
18	School environment and mental health in early adolescence - a longitudinal study in Sweden (KUPOL). <i>BMC Psychiatry</i> , 2016, 16, 243.	1.1	36

#	ARTICLE	IF	CITATIONS
19	Severity of Depression, Anxious Distress and the Risk of Cardiovascular Disease in a Swedish Population-Based Cohort. PLoS ONE, 2015, 10, e0140742.	1.1	34
20	The validity of the symptom checklist depression and anxiety subscales: A general population study in Sweden. Journal of Affective Disorders, 2015, 183, 247-252.	2.0	31
21	Comparison of DSM-5 Classifications of Alcohol Use Disorders With Those of DSM-IV, DSM-III-R, and ICD-10 in a General Population Sample in Sweden. Journal of Studies on Alcohol and Drugs, 2015, 76, 773-780.	0.6	30
22	Twelve-week physical exercise does not have a long-lasting effect on kynurenines in plasma of depressed patients. Neuropsychiatric Disease and Treatment, 2017, Volume 13, 967-972.	1.0	30
23	The Pathway to Meeting Need for Mental Health Services in Sweden. Psychiatric Services, 2006, 57, 114-119.	1.1	29
24	Adults with mild to moderate depression exhibit more alcohol related problems compared to the general adult population: a cross sectional study. BMC Public Health, 2015, 15, 542.	1.2	29
25	Effects of a synbiotic on symptoms, and daily functioning in attention deficit hyperactivity disorder "A double-blind randomized controlled trial. Brain, Behavior, and Immunity, 2020, 89, 9-19.	2.0	29
26	Well-Being in an Adult Swedish Population. Social Indicators Research, 2005, 74, 313-325.	1.4	27
27	Impact of Childhood Adversity and Vasopressin receptor 1a Variation on Social Interaction in Adulthood: A Cross-Sectional Study. PLoS ONE, 2015, 10, e0136436.	1.1	27
28	hTERT genetic variation in depression. Journal of Affective Disorders, 2016, 189, 62-69.	2.0	25
29	Effect of neuroticism on risk of cardiovascular disease in depressed persons - a Swedish population-based cohort study. BMC Cardiovascular Disorders, 2017, 17, 185.	0.7	25
30	Are changes in occupational physical activity level compensated by changes in exercise behavior?. European Journal of Public Health, 2018, 28, 940-943.	0.1	24
31	Effects of internet-based cognitive behavioural therapy and physical exercise on sick leave and employment in primary care patients with depression: two subgroup analyses. Occupational and Environmental Medicine, 2018, 75, 52-58.	1.3	22
32	Internet-based cognitive behaviour therapy and physical exercise "Effects studied by automated telephone assessments in mental ill-health patients; a randomized controlled trial. British Journal of Clinical Psychology, 2016, 55, 414-428.	1.7	21
33	Longitudinal association between tobacco use and the onset of depressive symptoms among Swedish adolescents: the Kupol cohort study. European Child and Adolescent Psychiatry, 2019, 28, 695-704.	2.8	21
34	Cost-effectiveness of internet-based cognitive-behavioural therapy and physical exercise for depression. BJPsych Open, 2018, 4, 265-273.	0.3	19
35	Is Sense of Coherence Lowered by Negative Life Events?. Journal of Happiness Studies, 2011, 12, 475-492.	1.9	17
36	Habitual physical activity levels predict treatment outcomes in depressed adults: A prospective cohort study. Preventive Medicine, 2016, 88, 53-58.	1.6	17

#	ARTICLE	IF	CITATIONS
37	Stress, depressive status and telomere length: Does social interaction and coping strategy play a mediating role?. <i>Journal of Affective Disorders</i> , 2017, 222, 138-145.	2.0	16
38	The serotonin transporter promoter variant (5-HTTLPR) and childhood adversity are associated with the personality trait openness to experience. <i>Psychiatry Research</i> , 2017, 257, 322-326.	1.7	16
39	Severity of depression, anxious distress and the risk of type 2 diabetes – a population-based cohort study in Sweden. <i>BMC Public Health</i> , 2019, 19, 1174.	1.2	15
40	Childhood adversity increases methylation in the GRIN2B gene. <i>Journal of Psychiatric Research</i> , 2021, 132, 38-43.	1.5	14
41	Association of Preeclampsia and Perinatal Complications With Offspring Neurodevelopmental and Psychiatric Disorders. <i>JAMA Network Open</i> , 2022, 5, e2145719.	2.8	14
42	BDNF Val66Met and childhood adversity on response to physical exercise and internet-based cognitive behavioural therapy in depressed Swedish adults. <i>Journal of Psychiatric Research</i> , 2017, 93, 50-58.	1.5	12
43	Utilization of psychiatric care and antidepressants among people with different severity of depression: a population-based cohort study in Stockholm, Sweden. <i>Social Psychiatry and Psychiatric Epidemiology</i> , 2018, 53, 607-615.	1.6	12
44	Exercise effects on cognitive functioning in young adults with first-episode psychosis: FitForLife. <i>Psychological Medicine</i> , 2019, 49, 431-439.	2.7	12
45	Do unfavourable alcohol, smoking, nutrition and physical activity predict sustained leisure time sedentary behaviour? A population-based cohort study. <i>Preventive Medicine</i> , 2017, 101, 23-27.	1.6	11
46	Neuropeptide Y, stressful life events and personality trait conscientiousness: Preliminary associations from a Swedish longitudinal study. <i>Psychiatry Research</i> , 2018, 263, 48-53.	1.7	10
47	The Effect of Smartphone Apps Versus Supervised Exercise on Physical Activity, Cardiorespiratory Fitness, and Body Composition Among Individuals With Mild-to-Moderate Mobility Disability: Randomized Controlled Trial. <i>JMIR MHealth and UHealth</i> , 2020, 8, e14615.	1.8	10
48	Psychiatric Symptoms, Social Disability, Low Wellbeing and Need for Treatment: Data from a Population-Based Study. <i>International Journal of Social Psychiatry</i> , 2004, 50, 195-203.	1.6	9
49	Psychological Symptoms in Late Adolescence and Long-Term Risk of Suicide and Suicide Attempt. <i>Suicide and Life-Threatening Behavior</i> , 2018, 48, 315-327.	0.9	9
50	Structured physical exercise and recovery from first episode psychosis in young adults, the FitForLife study. <i>Psychiatry Research</i> , 2018, 267, 346-353.	1.7	9
51	Unfavourable sedentary and physical activity behaviour before and after retirement: a population-based cohort study. <i>BMJ Open</i> , 2020, 10, e037659.	0.8	9
52	The cannabinoid receptor-1 gene interacts with stressful life events to increase the risk for problematic alcohol use. <i>Scientific Reports</i> , 2022, 12, 4963.	1.6	9
53	KIBRA genetic polymorphism and cognitive dysfunction in depression. <i>Psychiatry Research</i> , 2015, 226, 405-406.	1.7	8
54	Association of Catechol-O-methyltransferase (COMT Val158Met) with future risk of cardiovascular disease in depressed individuals - a Swedish population-based cohort study. <i>BMC Medical Genetics</i> , 2018, 19, 126.	2.1	8

#	ARTICLE	IF	CITATIONS
55	Changes in alcohol consumption after treatment for depression: a secondary analysis of the Swedish randomised controlled study REGASSA. <i>BMJ Open</i> , 2019, 9, e028236.	0.8	7
56	Latent class growth modelling for the evaluation of intervention outcomes: example from a physical activity intervention. <i>Journal of Behavioral Medicine</i> , 2021, 44, 622-629.	1.1	7
57	Path analysis of the chronicity of depression using the comprehensive developmental model framework. <i>Nordic Journal of Psychiatry</i> , 2016, 70, 380-391.	0.7	6
58	An eHealth program versus a standard care supervised health program and associated health outcomes in individuals with mobility disability: study protocol for a randomized controlled trial. <i>Trials</i> , 2018, 19, 258.	0.7	6
59	A pragmatic randomised trial of two counselling models at the Swedish national alcohol helpline. <i>BMC Psychiatry</i> , 2019, 19, 213.	1.1	6
60	Single-nucleotide polymorphism in the human TIA1 gene interacts with stressful life events to predict the development of pathological anxiety symptoms in a Swedish population. <i>Journal of Affective Disorders</i> , 2020, 260, 597-603.	2.0	6
61	Cannabis use, subsequent other illicit drug use and drug use disorders: A 16-year follow-up study among Swedish adults. <i>Addictive Behaviors</i> , 2020, 106, 106390.	1.7	6
62	Cortisol Concentration as Predictor of Tobacco Initiation in Adolescents: Results From a Population-Based Swedish Cohort. <i>Journal of Adolescent Health</i> , 2021, 68, 758-764.	1.2	6
63	DNA methylation of the glucocorticoid receptor gene predicts substance use in adolescence: longitudinal data from over 1000 young individuals. <i>Translational Psychiatry</i> , 2021, 11, 477.	2.4	6
64	Ethnic heterogeneity, social capital and psychological distress in Sweden. <i>Health and Place</i> , 2018, 52, 70-84.	1.5	5
65	Death of a Parent and the Risk of Ischemic Heart Disease and Stroke in Denmark and Sweden. <i>JAMA Network Open</i> , 2022, 5, e2218178.	2.8	5
66	Conceptual Design of an Extended Reality Exercise Game for the Elderly. <i>Applied Sciences (Switzerland)</i> , 2022, 12, 6436.	1.3	5
67	Psychosocial correlates with depressive symptoms six years after a first episode of psychosis as compared with findings from a general population sample. <i>BMC Psychiatry</i> , 2004, 4, 29.	1.1	4
68	Working conditions, serotonin transporter gene polymorphism (5-HTTLPR) and anxiety disorders: A prospective cohort study. <i>Journal of Affective Disorders</i> , 2013, 151, 652-659.	2.0	4
69	Increased screening of alcohol habits among patients with depression is needed. <i>Scandinavian Journal of Public Health</i> , 2014, 42, 658-659.	1.2	4
70	Comparing counselling models for the hazardous use of alcohol at the Swedish National Alcohol Helpline: study protocol for a randomised controlled trial. <i>Trials</i> , 2017, 18, 257.	0.7	4
71	Does depressed persons with non-cardiovascular morbidity have a higher risk of CVD? A population-based cohort study in Sweden. <i>BMC Cardiovascular Disorders</i> , 2019, 19, 260.	0.7	4
72	Childhood adversities, negative life events and outcomes of non-pharmacological treatments for depression in primary care: A secondary analysis of a randomized controlled trial. <i>Journal of Psychiatric Research</i> , 2019, 110, 152-158.	1.5	4

#	ARTICLE	IF	CITATIONS
73	Association of changes in cardiorespiratory fitness with health-related quality of life in young adults with mobility disability: secondary analysis of a randomized controlled trial of mobile app versus supervised training. BMC Public Health, 2020, 20, 1721.	1.2	4
74	Death of a parent during childhood and blood pressure in youth: a population-based cohort study of Swedish men. BMJ Open, 2021, 11, e043657.	0.8	4
75	FitForLife: study protocol for a randomized controlled trial. Trials, 2015, 16, 553.	0.7	3
76	Exercise Reduces Salivary Morning Cortisol Levels in Patients with Depression. Molecular Neuropsychiatry, 2018, 4, 196-203.	3.0	3
77	Physical exercise is associated with a reduction in plasma levels of fractalkine, TGF- β 1, eotaxin-1 and IL-6 in younger adults with mobility disability. PLoS ONE, 2022, 17, e0263173.	1.1	3
78	A Review of Extended Reality Exercise Games for Elderly. , 2022, , .		3
79	Impact of Expectations on the Effects of Exercise on Psychological Distress. American Journal of Health Behavior, 2014, 38, 650-656.	0.6	2
80	Physical exercise is associated with a reduction in inflammatory biomarkers in first-episode psychosis: A pilot study of CRP, SAA, sICAM-1 and sVCAM-1. Schizophrenia Research, 2021, 228, 316-318.	1.1	2
81	Laparoscopic Nissen fundoplication versus 90° and 180° fundoplication for gastroesophageal reflux disease. European Surgery - Acta Chirurgica Austriaca, 2022, 54, 66.	0.3	2
82	Health behaviours as a predictor of quitting hazardous alcohol use in the Stockholm Public Health Cohort. European Journal of Public Health, 2018, 28, 527-532.	0.1	1
83	No association of cigarette smoking and depressive symptoms with cortisol concentration in adolescents. Results from a population-based Swedish cohort. Psychiatry Research, 2021, 301, 113968.	1.7	1
84	Utilization of healthcare and prescription medicines after non-pharmacological interventions for depression - A 3-year register follow-up of an RCT in primary care. Preventive Medicine Reports, 2022, 25, 101658.	0.8	1
85	Tobacco use does not influence the response to non-pharmacologic depression treatment: A secondary analysis of the Regassa randomized controlled trial. Psychiatry Research, 2018, 261, 442-448.	1.7	0
86	Is There a Dose-Response Relationship between Acute Physical Activity and Sleep Length? A Longitudinal Study with Children and Adolescents Living in Sweden. Children, 2021, 8, 808.	0.6	0