

Yvonne Forsell

List of Publications by Year in descending order

Source: <https://exaly.com/author-pdf/4846061/publications.pdf>

Version: 2024-02-01

86
papers

2,208
citations

236612

25
h-index

264894

42
g-index

87
all docs

87
docs citations

87
times ranked

3497
citing authors

| # | ARTICLE | IF | CITATIONS |
|----|---|-----|-----------|
| 1 | Utilization of healthcare and prescription medicines after non-pharmacological interventions for depression - A 3-year register follow-up of an RCT in primary care. Preventive Medicine Reports, 2022, 25, 101658. | 0.8 | 1 |
| 2 | Laparoscopic Nissen fundoplication versus 90° and 180° fundoplication for gastroesophageal reflux disease. European Surgery - Acta Chirurgica Austriaca, 2022, 54, 66. | 0.3 | 2 |
| 3 | Association of Preeclampsia and Perinatal Complications With Offspring Neurodevelopmental and Psychiatric Disorders. JAMA Network Open, 2022, 5, e2145719. | 2.8 | 14 |
| 4 | Physical exercise is associated with a reduction in plasma levels of fractalkine, TGF- β 1, eotaxin-1 and IL-6 in younger adults with mobility disability. PLoS ONE, 2022, 17, e0263173. | 1.1 | 3 |
| 5 | A Review of Extended Reality Exercise Games for Elderly. , 2022, , . | | 3 |
| 6 | The cannabinoid receptor-1 gene interacts with stressful life events to increase the risk for problematic alcohol use. Scientific Reports, 2022, 12, 4963. | 1.6 | 9 |
| 7 | Death of a Parent and the Risk of Ischemic Heart Disease and Stroke in Denmark and Sweden. JAMA Network Open, 2022, 5, e2218178. | 2.8 | 5 |
| 8 | Conceptual Design of an Extended Reality Exercise Game for the Elderly. Applied Sciences (Switzerland), 2022, 12, 6436. | 1.3 | 5 |
| 9 | Childhood adversity increases methylation in the GRIN2B gene. Journal of Psychiatric Research, 2021, 132, 38-43. | 1.5 | 14 |
| 10 | Cortisol Concentration as Predictor of Tobacco Initiation in Adolescents: Results From a Population-Based Swedish Cohort. Journal of Adolescent Health, 2021, 68, 758-764. | 1.2 | 6 |
| 11 | Physical exercise is associated with a reduction in inflammatory biomarkers in first-episode psychosis: A pilot study of CRP, SAA, sICAM-1 and sVCAM-1. Schizophrenia Research, 2021, 228, 316-318. | 1.1 | 2 |
| 12 | Latent class growth modelling for the evaluation of intervention outcomes: example from a physical activity intervention. Journal of Behavioral Medicine, 2021, 44, 622-629. | 1.1 | 7 |
| 13 | Death of a parent during childhood and blood pressure in youth: a population-based cohort study of Swedish men. BMJ Open, 2021, 11, e043657. | 0.8 | 4 |
| 14 | Internet-Based Cognitive Behavioral Therapy for Depression. JAMA Psychiatry, 2021, 78, 361. | 6.0 | 398 |
| 15 | No association of cigarette smoking and depressive symptoms with cortisol concentration in adolescents. Results from a population-based Swedish cohort. Psychiatry Research, 2021, 301, 113968. | 1.7 | 1 |
| 16 | Is There a Dose-Response Relationship between Acute Physical Activity and Sleep Length? A Longitudinal Study with Children and Adolescents Living in Sweden. Children, 2021, 8, 808. | 0.6 | 0 |
| 17 | DNA methylation of the glucocorticoid receptor gene predicts substance use in adolescence: longitudinal data from over 1000 young individuals. Translational Psychiatry, 2021, 11, 477. | 2.4 | 6 |
| 18 | Single-nucleotide polymorphism in the human TIA1 gene interacts with stressful life events to predict the development of pathological anxiety symptoms in a Swedish population. Journal of Affective Disorders, 2020, 260, 597-603. | 2.0 | 6 |

| # | ARTICLE | IF | CITATIONS |
|----|---|-----|-----------|
| 19 | Association of changes in cardiorespiratory fitness with health-related quality of life in young adults with mobility disability: secondary analysis of a randomized controlled trial of mobile app versus supervised training. BMC Public Health, 2020, 20, 1721. | 1.2 | 4 |
| 20 | Unfavourable sedentary and physical activity behaviour before and after retirement: a population-based cohort study. BMJ Open, 2020, 10, e037659. | 0.8 | 9 |
| 21 | Effects of a synbiotic on symptoms, and daily functioning in attention deficit hyperactivity disorder "A double-blind randomized controlled trial. Brain, Behavior, and Immunity, 2020, 89, 9-19. | 2.0 | 29 |
| 22 | Cannabis use, subsequent other illicit drug use and drug use disorders: A 16-year follow-up study among Swedish adults. Addictive Behaviors, 2020, 106, 106390. | 1.7 | 6 |
| 23 | The Effect of Smartphone Apps Versus Supervised Exercise on Physical Activity, Cardiorespiratory Fitness, and Body Composition Among Individuals With Mild-to-Moderate Mobility Disability: Randomized Controlled Trial. JMIR MHealth and UHealth, 2020, 8, e14615. | 1.8 | 10 |
| 24 | Exercise effects on cognitive functioning in young adults with first-episode psychosis: FitForLife. Psychological Medicine, 2019, 49, 431-439. | 2.7 | 12 |
| 25 | A pragmatic randomised trial of two counselling models at the Swedish national alcohol helpline. BMC Psychiatry, 2019, 19, 213. | 1.1 | 6 |
| 26 | Severity of depression, anxious distress and the risk of type 2 diabetes "a population-based cohort study in Sweden. BMC Public Health, 2019, 19, 1174. | 1.2 | 15 |
| 27 | Changes in alcohol consumption after treatment for depression: a secondary analysis of the Swedish randomised controlled study REGASSA. BMJ Open, 2019, 9, e028236. | 0.8 | 7 |
| 28 | Early exposure to antibiotic drugs and risk for psychiatric disorders: a population-based study. Translational Psychiatry, 2019, 9, 317. | 2.4 | 60 |
| 29 | Does depressed persons with non-cardiovascular morbidity have a higher risk of CVD? A population-based cohort study in Sweden. BMC Cardiovascular Disorders, 2019, 19, 260. | 0.7 | 4 |
| 30 | Childhood adversities, negative life events and outcomes of non-pharmacological treatments for depression in primary care: A secondary analysis of a randomized controlled trial. Journal of Psychiatric Research, 2019, 110, 152-158. | 1.5 | 4 |
| 31 | Longitudinal association between tobacco use and the onset of depressive symptoms among Swedish adolescents: the Kupol cohort study. European Child and Adolescent Psychiatry, 2019, 28, 695-704. | 2.8 | 21 |
| 32 | Neuropeptide Y, stressful life events and personality trait conscientiousness: Preliminary associations from a Swedish longitudinal study. Psychiatry Research, 2018, 263, 48-53. | 1.7 | 10 |
| 33 | Utilization of psychiatric care and antidepressants among people with different severity of depression: a population-based cohort study in Stockholm, Sweden. Social Psychiatry and Psychiatric Epidemiology, 2018, 53, 607-615. | 1.6 | 12 |
| 34 | Tobacco use does not influence the response to non-pharmacologic depression treatment: A secondary analysis of the Regassa randomized controlled trial. Psychiatry Research, 2018, 261, 442-448. | 1.7 | 0 |
| 35 | Are changes in occupational physical activity level compensated by changes in exercise behavior?. European Journal of Public Health, 2018, 28, 940-943. | 0.1 | 24 |
| 36 | Psychological Symptoms in Late Adolescence and Long-Term Risk of Suicide and Suicide Attempt. Suicide and Life-Threatening Behavior, 2018, 48, 315-327. | 0.9 | 9 |

| # | ARTICLE | IF | CITATIONS |
|----|---|-----|-----------|
| 37 | Effects of internet-based cognitive behavioural therapy and physical exercise on sick leave and employment in primary care patients with depression: two subgroup analyses. <i>Occupational and Environmental Medicine</i> , 2018, 75, 52-58. | 1.3 | 22 |
| 38 | Exercise Reduces Salivary Morning Cortisol Levels in Patients with Depression. <i>Molecular Neuropsychiatry</i> , 2018, 4, 196-203. | 3.0 | 3 |
| 39 | Ethnic heterogeneity, social capital and psychological distress in Sweden. <i>Health and Place</i> , 2018, 52, 70-84. | 1.5 | 5 |
| 40 | Structured physical exercise and recovery from first episode psychosis in young adults, the FitForLife study. <i>Psychiatry Research</i> , 2018, 267, 346-353. | 1.7 | 9 |
| 41 | Association of Catechol-O-methyltransferase (COMT Val158Met) with future risk of cardiovascular disease in depressed individuals - a Swedish population-based cohort study. <i>BMC Medical Genetics</i> , 2018, 19, 126. | 2.1 | 8 |
| 42 | Common Perceived Barriers and Facilitators for Reducing Sedentary Behaviour among Office Workers. <i>International Journal of Environmental Research and Public Health</i> , 2018, 15, 792. | 1.2 | 38 |
| 43 | An eHealth program versus a standard care supervised health program and associated health outcomes in individuals with mobility disability: study protocol for a randomized controlled trial. <i>Trials</i> , 2018, 19, 258. | 0.7 | 6 |
| 44 | Cost-effectiveness of internet-based cognitive behavioural therapy and physical exercise for depression. <i>BJPsych Open</i> , 2018, 4, 265-273. | 0.3 | 19 |
| 45 | Health behaviours as a predictor of quitting hazardous alcohol use in the Stockholm Public Health Cohort. <i>European Journal of Public Health</i> , 2018, 28, 527-532. | 0.1 | 1 |
| 46 | NR3C1 hypermethylation in depressed and bullied adolescents. <i>Translational Psychiatry</i> , 2018, 8, 121. | 2.4 | 46 |
| 47 | Somebody to lean on: Social relationships predict post-treatment depression severity in adults. <i>Psychiatry Research</i> , 2017, 249, 261-267. | 1.7 | 46 |
| 48 | Interleukin-6 and depressive symptom severity in response to physical exercise. <i>Psychiatry Research</i> , 2017, 252, 270-276. | 1.7 | 49 |
| 49 | Do unfavourable alcohol, smoking, nutrition and physical activity predict sustained leisure time sedentary behaviour? A population-based cohort study. <i>Preventive Medicine</i> , 2017, 101, 23-27. | 1.6 | 11 |
| 50 | BDNF Val66Met and childhood adversity on response to physical exercise and internet-based cognitive behavioural therapy in depressed Swedish adults. <i>Journal of Psychiatric Research</i> , 2017, 93, 50-58. | 1.5 | 12 |
| 51 | Stress, depressive status and telomere length: Does social interaction and coping strategy play a mediating role?. <i>Journal of Affective Disorders</i> , 2017, 222, 138-145. | 2.0 | 16 |
| 52 | The serotonin transporter promoter variant (5-HTTLPR) and childhood adversity are associated with the personality trait openness to experience. <i>Psychiatry Research</i> , 2017, 257, 322-326. | 1.7 | 16 |
| 53 | Long-term effects of exercise at different intensity levels on depression: A randomized controlled trial. <i>Preventive Medicine</i> , 2017, 105, 37-46. | 1.6 | 46 |
| 54 | Comparing counselling models for the hazardous use of alcohol at the Swedish National Alcohol Helpline: study protocol for a randomised controlled trial. <i>Trials</i> , 2017, 18, 257. | 0.7 | 4 |

| # | ARTICLE | IF | CITATIONS |
|----|--|-----|-----------|
| 55 | Effect of neuroticism on risk of cardiovascular disease in depressed persons - a Swedish population-based cohort study. <i>BMC Cardiovascular Disorders</i> , 2017, 17, 185. | 0.7 | 25 |
| 56 | Twelve-week physical exercise does not have a long-lasting effect on kynurenines in plasma of depressed patients. <i>Neuropsychiatric Disease and Treatment</i> , 2017, Volume 13, 967-972. | 1.0 | 30 |
| 57 | Exercise, Physical Activity, and Sedentary Behavior in the Treatment of Depression: Broadening the Scientific Perspectives and Clinical Opportunities. <i>Frontiers in Psychiatry</i> , 2016, 7, 36. | 1.3 | 71 |
| 58 | Habitual physical activity levels predict treatment outcomes in depressed adults: A prospective cohort study. <i>Preventive Medicine</i> , 2016, 88, 53-58. | 1.6 | 17 |
| 59 | Training fast or slow? Exercise for depression: A randomized controlled trial. <i>Preventive Medicine</i> , 2016, 91, 123-131. | 1.6 | 60 |
| 60 | School environment and mental health in early adolescence - a longitudinal study in Sweden (KUPOL). <i>BMC Psychiatry</i> , 2016, 16, 243. | 1.1 | 36 |
| 61 | Exercise and internet-based cognitive-behavioural therapy for depression: multicentre randomised controlled trial with 12-month follow-up. <i>British Journal of Psychiatry</i> , 2016, 209, 414-420. | 1.7 | 57 |
| 62 | Internet-based cognitive behaviour therapy and physical exercise - Effects studied by automated telephone assessments in mental ill-health patients; a randomized controlled trial. <i>British Journal of Clinical Psychology</i> , 2016, 55, 414-428. | 1.7 | 21 |
| 63 | Cannabis use, depression and anxiety: A 3-year prospective population-based study. <i>Journal of Affective Disorders</i> , 2016, 193, 103-108. | 2.0 | 78 |
| 64 | Path analysis of the chronicity of depression using the comprehensive developmental model framework. <i>Nordic Journal of Psychiatry</i> , 2016, 70, 380-391. | 0.7 | 6 |
| 65 | hTERT genetic variation in depression. <i>Journal of Affective Disorders</i> , 2016, 189, 62-69. | 2.0 | 25 |
| 66 | Comparison of DSM-5 Classifications of Alcohol Use Disorders With Those of DSM-IV, DSM-III-R, and ICD-10 in a General Population Sample in Sweden. <i>Journal of Studies on Alcohol and Drugs</i> , 2015, 76, 773-780. | 0.6 | 30 |
| 67 | FitForLife: study protocol for a randomized controlled trial. <i>Trials</i> , 2015, 16, 553. | 0.7 | 3 |
| 68 | Physical Activity Patterns of People Affected by Depressive and Anxiety Disorders as Measured by Accelerometers: A Cross-Sectional Study. <i>PLoS ONE</i> , 2015, 10, e0115894. | 1.1 | 91 |
| 69 | Impact of Childhood Adversity and Vasopressin receptor 1a Variation on Social Interaction in Adulthood: A Cross-Sectional Study. <i>PLoS ONE</i> , 2015, 10, e0136436. | 1.1 | 27 |
| 70 | The validity of the symptom checklist depression and anxiety subscales: A general population study in Sweden. <i>Journal of Affective Disorders</i> , 2015, 183, 247-252. | 2.0 | 31 |
| 71 | KIBRA genetic polymorphism and cognitive dysfunction in depression. <i>Psychiatry Research</i> , 2015, 226, 405-406. | 1.7 | 8 |
| 72 | Hypomethylation of MAOA's first exon region in depression: A replication study. <i>Psychiatry Research</i> , 2015, 226, 389-391. | 1.7 | 37 |

| # | ARTICLE | IF | CITATIONS |
|----|--|-----|-----------|
| 73 | Physical exercise and internet-based cognitive-behavioural therapy in the treatment of depression: Randomised controlled trial. <i>British Journal of Psychiatry</i> , 2015, 207, 227-234. | 1.7 | 118 |
| 74 | Adults with mild to moderate depression exhibit more alcohol related problems compared to the general adult population: a cross sectional study. <i>BMC Public Health</i> , 2015, 15, 542. | 1.2 | 29 |
| 75 | Severity of Depression, Anxious Distress and the Risk of Cardiovascular Disease in a Swedish Population-Based Cohort. <i>PLoS ONE</i> , 2015, 10, e0140742. | 1.1 | 34 |
| 76 | Impact of Expectations on the Effects of Exercise on Psychological Distress. <i>American Journal of Health Behavior</i> , 2014, 38, 650-656. | 0.6 | 2 |
| 77 | Increased screening of alcohol habits among patients with depression is needed. <i>Scandinavian Journal of Public Health</i> , 2014, 42, 658-659. | 1.2 | 4 |
| 78 | Working conditions, serotonin transporter gene polymorphism (5-HTTLPR) and anxiety disorders: A prospective cohort study. <i>Journal of Affective Disorders</i> , 2013, 151, 652-659. | 2.0 | 4 |
| 79 | Is Sense of Coherence Lowered by Negative Life Events?. <i>Journal of Happiness Studies</i> , 2011, 12, 475-492. | 1.9 | 17 |
| 80 | A three-year follow-up of major depression, dysthymia, minor depression and subsyndromal depression: results from a population-based study. <i>Depression and Anxiety</i> , 2007, 24, 62-65. | 2.0 | 58 |
| 81 | The Pathway to Meeting Need for Mental Health Services in Sweden. <i>Psychiatric Services</i> , 2006, 57, 114-119. | 1.1 | 29 |
| 82 | Well-Being in an Adult Swedish Population. <i>Social Indicators Research</i> , 2005, 74, 313-325. | 1.4 | 27 |
| 83 | Psychiatric Symptoms, Social Disability, Low Wellbeing and Need for Treatment: Data from a Population-Based Study. <i>International Journal of Social Psychiatry</i> , 2004, 50, 195-203. | 1.6 | 9 |
| 84 | Psychosocial correlates with depressive symptoms six years after a first episode of psychosis as compared with findings from a general population sample. <i>BMC Psychiatry</i> , 2004, 4, 29. | 1.1 | 4 |
| 85 | Psychotropic drug use in elderly people with and without dementia. <i>International Journal of Geriatric Psychiatry</i> , 2001, 16, 900-906. | 1.3 | 72 |
| 86 | Morbidity and Comorbidity in Relation to Functional Status: A Community-Based Study of the Oldest Old (90+ Years). <i>Journal of the American Geriatrics Society</i> , 2000, 48, 1462-1469. | 1.3 | 63 |