

Yvonne Forsell

List of Publications by Year in descending order

Source: <https://exaly.com/author-pdf/4846061/publications.pdf>

Version: 2024-02-01

86
papers

2,208
citations

236925
25
h-index

265206
42
g-index

87
all docs

87
docs citations

87
times ranked

3497
citing authors

#	ARTICLE	IF	CITATIONS
1	Utilization of healthcare and prescription medicines after non-pharmacological interventions for depression - A 3-year register follow-up of an RCT in primary care. Preventive Medicine Reports, 2022, 25, 101658.	1.8	1
2	Laparoscopic Nissen fundoplication versus 90° and 180° fundoplication for gastroesophageal reflux disease. European Surgery - Acta Chirurgica Austriaca, 2022, 54, 66.	0.7	2
3	Association of Preeclampsia and Perinatal Complications With Offspring Neurodevelopmental and Psychiatric Disorders. JAMA Network Open, 2022, 5, e2145719.	5.9	14
4	Physical exercise is associated with a reduction in plasma levels of fractalkine, TGF- β 1, eotaxin-1 and IL-6 in younger adults with mobility disability. PLoS ONE, 2022, 17, e0263173.	2.5	3
5	A Review of Extended Reality Exercise Games for Elderly. , 2022, , .		3
6	The cannabinoid receptor-1 gene interacts with stressful life events to increase the risk for problematic alcohol use. Scientific Reports, 2022, 12, 4963.	3.3	9
7	Death of a Parent and the Risk of Ischemic Heart Disease and Stroke in Denmark and Sweden. JAMA Network Open, 2022, 5, e2218178.	5.9	5
8	Conceptual Design of an Extended Reality Exercise Game for the Elderly. Applied Sciences (Switzerland), 2022, 12, 6436.	2.5	5
9	Childhood adversity increases methylation in the GRIN2B gene. Journal of Psychiatric Research, 2021, 132, 38-43.	3.1	14
10	Cortisol Concentration as Predictor of Tobacco Initiation in Adolescents: Results From a Population-Based Swedish Cohort. Journal of Adolescent Health, 2021, 68, 758-764.	2.5	6
11	Physical exercise is associated with a reduction in inflammatory biomarkers in first-episode psychosis: A pilot study of CRP, SAA, sICAM-1 and sVCAM-1. Schizophrenia Research, 2021, 228, 316-318.	2.0	2
12	Latent class growth modelling for the evaluation of intervention outcomes: example from a physical activity intervention. Journal of Behavioral Medicine, 2021, 44, 622-629.	2.1	7
13	Death of a parent during childhood and blood pressure in youth: a population-based cohort study of Swedish men. BMJ Open, 2021, 11, e043657.	1.9	4
14	Internet-Based Cognitive Behavioral Therapy for Depression. JAMA Psychiatry, 2021, 78, 361.	11.0	398
15	No association of cigarette smoking and depressive symptoms with cortisol concentration in adolescents. Results from a population-based Swedish cohort. Psychiatry Research, 2021, 301, 113968.	3.3	1
16	Is There a Dose-Response Relationship between Acute Physical Activity and Sleep Length? A Longitudinal Study with Children and Adolescents Living in Sweden. Children, 2021, 8, 808.	1.5	0
17	DNA methylation of the glucocorticoid receptor gene predicts substance use in adolescence: longitudinal data from over 1000 young individuals. Translational Psychiatry, 2021, 11, 477.	4.8	6
18	Single-nucleotide polymorphism in the human TIA1 gene interacts with stressful life events to predict the development of pathological anxiety symptoms in a Swedish population. Journal of Affective Disorders, 2020, 260, 597-603.	4.1	6

#	ARTICLE	IF	CITATIONS
19	Association of changes in cardiorespiratory fitness with health-related quality of life in young adults with mobility disability: secondary analysis of a randomized controlled trial of mobile app versus supervised training. BMC Public Health, 2020, 20, 1721.	2.9	4
20	Unfavourable sedentary and physical activity behaviour before and after retirement: a population-based cohort study. BMJ Open, 2020, 10, e037659.	1.9	9
21	Effects of a synbiotic on symptoms, and daily functioning in attention deficit hyperactivity disorder “A double-blind randomized controlled trial. Brain, Behavior, and Immunity, 2020, 89, 9-19.	4.1	29
22	Cannabis use, subsequent other illicit drug use and drug use disorders: A 16-year follow-up study among Swedish adults. Addictive Behaviors, 2020, 106, 106390.	3.0	6
23	The Effect of Smartphone Apps Versus Supervised Exercise on Physical Activity, Cardiorespiratory Fitness, and Body Composition Among Individuals With Mild-to-Moderate Mobility Disability: Randomized Controlled Trial. JMIR MHealth and UHealth, 2020, 8, e14615.	3.7	10
24	Exercise effects on cognitive functioning in young adults with first-episode psychosis: FitForLife. Psychological Medicine, 2019, 49, 431-439.	4.5	12
25	A pragmatic randomised trial of two counselling models at the Swedish national alcohol helpline. BMC Psychiatry, 2019, 19, 213.	2.6	6
26	Severity of depression, anxious distress and the risk of type 2 diabetes “a population-based cohort study in Sweden. BMC Public Health, 2019, 19, 1174.	2.9	15
27	Changes in alcohol consumption after treatment for depression: a secondary analysis of the Swedish randomised controlled study REGASSA. BMJ Open, 2019, 9, e028236.	1.9	7
28	Early exposure to antibiotic drugs and risk for psychiatric disorders: a population-based study. Translational Psychiatry, 2019, 9, 317.	4.8	60
29	Does depressed persons with non-cardiovascular morbidity have a higher risk of CVD? A population-based cohort study in Sweden. BMC Cardiovascular Disorders, 2019, 19, 260.	1.7	4
30	Childhood adversities, negative life events and outcomes of non-pharmacological treatments for depression in primary care: A secondary analysis of a randomized controlled trial. Journal of Psychiatric Research, 2019, 110, 152-158.	3.1	4
31	Longitudinal association between tobacco use and the onset of depressive symptoms among Swedish adolescents: the Kupol cohort study. European Child and Adolescent Psychiatry, 2019, 28, 695-704.	4.7	21
32	Neuropeptide Y, stressful life events and personality trait conscientiousness: Preliminary associations from a Swedish longitudinal study. Psychiatry Research, 2018, 263, 48-53.	3.3	10
33	Utilization of psychiatric care and antidepressants among people with different severity of depression: a population-based cohort study in Stockholm, Sweden. Social Psychiatry and Psychiatric Epidemiology, 2018, 53, 607-615.	3.1	12
34	Tobacco use does not influence the response to non-pharmacologic depression treatment: A secondary analysis of the Regassa randomized controlled trial. Psychiatry Research, 2018, 261, 442-448.	3.3	0
35	Are changes in occupational physical activity level compensated by changes in exercise behavior?. European Journal of Public Health, 2018, 28, 940-943.	0.3	24
36	Psychological Symptoms in Late Adolescence and Long-term Risk of Suicide and Suicide Attempt. Suicide and Life-Threatening Behavior, 2018, 48, 315-327.	1.9	9

#	ARTICLE	IF	CITATIONS
37	Effects of internet-based cognitive behavioural therapy and physical exercise on sick leave and employment in primary care patients with depression: two subgroup analyses. Occupational and Environmental Medicine, 2018, 75, 52-58.	2.8	22
38	Exercise Reduces Salivary Morning Cortisol Levels in Patients with Depression. Molecular Neuropsychiatry, 2018, 4, 196-203.	2.9	3
39	Ethnic heterogeneity, social capital and psychological distress in Sweden. Health and Place, 2018, 52, 70-84.	3.3	5
40	Structured physical exercise and recovery from first episode psychosis in young adults, the FitForLife study. Psychiatry Research, 2018, 267, 346-353.	3.3	9
41	Association of Catechol-O-methyltransferase (COMT Val158Met) with future risk of cardiovascular disease in depressed individuals - a Swedish population-based cohort study. BMC Medical Genetics, 2018, 19, 126.	2.1	8
42	Common Perceived Barriers and Facilitators for Reducing Sedentary Behaviour among Office Workers. International Journal of Environmental Research and Public Health, 2018, 15, 792.	2.6	38
43	An eHealth program versus a standard care supervised health program and associated health outcomes in individuals with mobility disability: study protocol for a randomized controlled trial. Trials, 2018, 19, 258.	1.6	6
44	Cost-effectiveness of internet-based cognitive behavioural therapy and physical exercise for depression. BJPsych Open, 2018, 4, 265-273.	0.7	19
45	Health behaviours as a predictor of quitting hazardous alcohol use in the Stockholm Public Health Cohort. European Journal of Public Health, 2018, 28, 527-532.	0.3	1
46	NR3C1 hypermethylation in depressed and bullied adolescents. Translational Psychiatry, 2018, 8, 121.	4.8	46
47	Somebody to lean on: Social relationships predict post-treatment depression severity in adults. Psychiatry Research, 2017, 249, 261-267.	3.3	46
48	Interleukin-6 and depressive symptom severity in response to physical exercise. Psychiatry Research, 2017, 252, 270-276.	3.3	49
49	Do unfavourable alcohol, smoking, nutrition and physical activity predict sustained leisure time sedentary behaviour? A population-based cohort study. Preventive Medicine, 2017, 101, 23-27.	3.4	11
50	BDNF Val66Met and childhood adversity on response to physical exercise and internet-based cognitive behavioural therapy in depressed Swedish adults. Journal of Psychiatric Research, 2017, 93, 50-58.	3.1	12
51	Stress, depressive status and telomere length: Does social interaction and coping strategy play a mediating role?. Journal of Affective Disorders, 2017, 222, 138-145.	4.1	16
52	The serotonin transporter promoter variant (5-HTTLPR) and childhood adversity are associated with the personality trait openness to experience. Psychiatry Research, 2017, 257, 322-326.	3.3	16
53	Long-term effects of exercise at different intensity levels on depression: A randomized controlled trial. Preventive Medicine, 2017, 105, 37-46.	3.4	46
54	Comparing counselling models for the hazardous use of alcohol at the Swedish National Alcohol Helpline: study protocol for a randomised controlled trial. Trials, 2017, 18, 257.	1.6	4

#	ARTICLE	IF	CITATIONS
55	Effect of neuroticism on risk of cardiovascular disease in depressed persons - a Swedish population-based cohort study. BMC Cardiovascular Disorders, 2017, 17, 185.	1.7	25
56	Twelve-week physical exercise does not have a long-lasting effect on kynurenines in plasma of depressed patients. Neuropsychiatric Disease and Treatment, 2017, Volume 13, 967-972.	2.2	30
57	Exercise, Physical Activity, and Sedentary Behavior in the Treatment of Depression: Broadening the Scientific Perspectives and Clinical Opportunities. Frontiers in Psychiatry, 2016, 7, 36.	2.6	71
58	Habitual physical activity levels predict treatment outcomes in depressed adults: A prospective cohort study. Preventive Medicine, 2016, 88, 53-58.	3.4	17
59	Training fast or slow? Exercise for depression: A randomized controlled trial. Preventive Medicine, 2016, 91, 123-131.	3.4	60
60	School environment and mental health in early adolescence - a longitudinal study in Sweden (KUPOL). BMC Psychiatry, 2016, 16, 243.	2.6	36
61	Exercise and internet-based cognitive-behavioural therapy for depression: multicentre randomised controlled trial with 12-month follow-up. British Journal of Psychiatry, 2016, 209, 414-420.	2.8	57
62	Internet-based cognitive behaviour therapy and physical exercise - Effects studied by automated telephone assessments in mental ill-health patients; a randomized controlled trial. British Journal of Clinical Psychology, 2016, 55, 414-428.	3.5	21
63	Cannabis use, depression and anxiety: A 3-year prospective population-based study. Journal of Affective Disorders, 2016, 193, 103-108.	4.1	78
64	Path analysis of the chronicity of depression using the comprehensive developmental model framework. Nordic Journal of Psychiatry, 2016, 70, 380-391.	1.3	6
65	hTERT genetic variation in depression. Journal of Affective Disorders, 2016, 189, 62-69.	4.1	25
66	Comparison of DSM-5 Classifications of Alcohol Use Disorders With Those of DSM-IV, DSM-III-R, and ICD-10 in a General Population Sample in Sweden. Journal of Studies on Alcohol and Drugs, 2015, 76, 773-780.	1.0	30
67	FitForLife: study protocol for a randomized controlled trial. Trials, 2015, 16, 553.	1.6	3
68	Physical Activity Patterns of People Affected by Depressive and Anxiety Disorders as Measured by Accelerometers: A Cross-Sectional Study. PLoS ONE, 2015, 10, e0115894.	2.5	91
69	Impact of Childhood Adversity and Vasopressin receptor 1a Variation on Social Interaction in Adulthood: A Cross-Sectional Study. PLoS ONE, 2015, 10, e0136436.	2.5	27
70	The validity of the symptom checklist depression and anxiety subscales: A general population study in Sweden. Journal of Affective Disorders, 2015, 183, 247-252.	4.1	31
71	KIBRA genetic polymorphism and cognitive dysfunction in depression. Psychiatry Research, 2015, 226, 405-406.	3.3	8
72	Hypomethylation of MAOA's first exon region in depression: A replication study. Psychiatry Research, 2015, 226, 389-391.	3.3	37

#	ARTICLE	IF	CITATIONS
73	Physical exercise and internet-based cognitive-behavioural therapy in the treatment of depression: Randomised controlled trial. <i>British Journal of Psychiatry</i> , 2015, 207, 227-234.	2.8	118
74	Adults with mild to moderate depression exhibit more alcohol related problems compared to the general adult population: a cross sectional study. <i>BMC Public Health</i> , 2015, 15, 542.	2.9	29
75	Severity of Depression, Anxious Distress and the Risk of Cardiovascular Disease in a Swedish Population-Based Cohort. <i>PLoS ONE</i> , 2015, 10, e0140742.	2.5	34
76	Impact of Expectations on the Effects of Exercise on Psychological Distress. <i>American Journal of Health Behavior</i> , 2014, 38, 650-656.	1.4	2
77	Increased screening of alcohol habits among patients with depression is needed. <i>Scandinavian Journal of Public Health</i> , 2014, 42, 658-659.	2.3	4
78	Working conditions, serotonin transporter gene polymorphism (5-HTTLPR) and anxiety disorders: A prospective cohort study. <i>Journal of Affective Disorders</i> , 2013, 151, 652-659.	4.1	4
79	Is Sense of Coherence Lowered by Negative Life Events?. <i>Journal of Happiness Studies</i> , 2011, 12, 475-492.	3.2	17
80	A three-year follow-up of major depression, dysthymia, minor depression and subsyndromal depression: results from a population-based study. <i>Depression and Anxiety</i> , 2007, 24, 62-65.	4.1	58
81	The Pathway to Meeting Need for Mental Health Services in Sweden. <i>Psychiatric Services</i> , 2006, 57, 114-119.	2.0	29
82	Well-Being in an Adult Swedish Population. <i>Social Indicators Research</i> , 2005, 74, 313-325.	2.7	27
83	Psychiatric Symptoms, Social Disability, Low Wellbeing and Need for Treatment: Data from a Population-Based Study. <i>International Journal of Social Psychiatry</i> , 2004, 50, 195-203.	3.1	9
84	Psychosocial correlates with depressive symptoms six years after a first episode of psychosis as compared with findings from a general population sample. <i>BMC Psychiatry</i> , 2004, 4, 29.	2.6	4
85	Psychotropic drug use in elderly people with and without dementia. <i>International Journal of Geriatric Psychiatry</i> , 2001, 16, 900-906.	2.7	72
86	Morbidity and Comorbidity in Relation to Functional Status: A Community-Based Study of the Oldest Old (90+ Years). <i>Journal of the American Geriatrics Society</i> , 2000, 48, 1462-1469.	2.6	63