

Jean Charron

List of Publications by Year in descending order

Source: <https://exaly.com/author-pdf/4853107/publications.pdf>

Version: 2024-02-01

58
papers

6,107
citations

147801

31
h-index

155660

55
g-index

61
all docs

61
docs citations

61
times ranked

8125
citing authors

#	ARTICLE	IF	CITATIONS
1	RAG-2-deficient mice lack mature lymphocytes owing to inability to initiate V(D)J rearrangement. <i>Cell</i> , 1992, 68, 855-867.	28.9	2,426
2	Embryonic death of Mek1-deficient mice reveals a role for this kinase in angiogenesis in the labyrinthine region of the placenta. <i>Current Biology</i> , 1999, 9, 369-376.	3.9	313
3	Embryonic lethality in mice homozygous for a targeted disruption of the N-myc gene.. <i>Genes and Development</i> , 1992, 6, 2248-2257.	5.9	280
4	c-Raf, but Not B-Raf, Is Essential for Development of K-Ras Oncogene-Driven Non-Small Cell Lung Carcinoma. <i>Cancer Cell</i> , 2011, 19, 652-663.	16.8	260
5	Prolonged Mek1/2 suppression impairs the developmental potential of embryonic stem cells. <i>Nature</i> , 2017, 548, 219-223.	27.8	211
6	Structure of the rat pro-opiomelanocortin (POMC) gene. <i>FEBS Letters</i> , 1985, 193, 54-58.	2.8	174
7	Specification of axial identity in the mouse: role of the Hoxa-5 (Hox1.3) gene.. <i>Genes and Development</i> , 1993, 7, 2085-2096.	5.9	169
8	<i>Mek2</i> Is Dispensable for Mouse Growth and Development. <i>Molecular and Cellular Biology</i> , 2003, 23, 4778-4787.	2.3	164
9	Mouse and human phenotypes indicate a critical conserved role for ERK2 signaling in neural crest development. <i>Proceedings of the National Academy of Sciences of the United States of America</i> , 2008, 105, 17115-17120.	7.1	159
10	Role of <i>Plk2</i> (<i>Snk</i>) in Mouse Development and Cell Proliferation. <i>Molecular and Cellular Biology</i> , 2003, 23, 6936-6943.	2.3	146
11	MEK Is a Key Regulator of Gliogenesis in the Developing Brain. <i>Neuron</i> , 2012, 75, 1035-1050.	8.1	145
12	Mek1/2 MAPK Kinases Are Essential for Mammalian Development, Homeostasis, and Raf-Induced Hyperplasia. <i>Developmental Cell</i> , 2007, 12, 615-629.	7.0	132
13	Pro-opiomelanocortin gene: A model for negative regulation of transcription by glucocorticoids. <i>Journal of Cellular Biochemistry</i> , 1987, 35, 293-304.	2.6	109
14	Phosphorylation Is Involved in the Activation of Metal-regulatory Transcription Factor 1 in Response to Metal Ions. <i>Journal of Biological Chemistry</i> , 2001, 276, 41879-41888.	3.4	107
15	Activated MEK Suppresses Activation of PKR and Enables Efficient Replication and In Vivo Oncolysis by γ^3 1 34.5 Mutants of Herpes Simplex Virus 1. <i>Journal of Virology</i> , 2006, 80, 1110-1120.	3.4	103
16	Glucocorticoid inhibition of transcription from episomal proopioidmelanocortin gene promoter.. <i>Proceedings of the National Academy of Sciences of the United States of America</i> , 1986, 83, 8903-8907.	7.1	98
17	B-RAF kinase drives developmental axon growth and promotes axon regeneration in the injured mature CNS. <i>Journal of Experimental Medicine</i> , 2014, 211, 801-814.	8.5	86
18	Requirement for <i>Map2k1</i> (<i>Mek1</i>) in extra-embryonic ectoderm during placentogenesis. <i>Development (Cambridge)</i> , 2006, 133, 3429-3440.	2.5	79

#	ARTICLE	IF	CITATIONS
19	The Leydig Cell MEK/ERK Pathway Is Critical for Maintaining a Functional Population of Adult Leydig Cells and for Fertility. <i>Molecular Endocrinology</i> , 2011, 25, 1211-1222.	3.7	64
20	Crucial requirement of ERK/MAPK signaling in respiratory tract development. <i>Development (Cambridge)</i> , 2014, 141, 3197-3211.	2.5	62
21	Mitogen-Activated Protein Kinase (MAPK) Pathway Regulates Branching by Remodeling Epithelial Cell Adhesion. <i>PLoS Genetics</i> , 2014, 10, e1004193.	3.5	59
22	Essential role of the ERK/MAPK pathway in blood-placental barrier formation. <i>Development (Cambridge)</i> , 2014, 141, 2825-2837.	2.5	56
23	Selective Role for Mek1 but not Mek2 in the Induction of Epidermal Neoplasia. <i>Cancer Research</i> , 2009, 69, 3772-3778.	0.9	54
24	Schnurri-3 regulates ERK downstream of WNT signaling in osteoblasts. <i>Journal of Clinical Investigation</i> , 2013, 123, 4010-4022.	8.2	53
25	Tissue-specific activity of the pro-opiomelanocortin (POMC) gene and repression by glucocorticoids. <i>Genome</i> , 1989, 31, 510-519.	2.0	47
26	<i>Map2k1</i> and <i>Map2k2</i> genes contribute to the normal development of syncytiotrophoblasts during placentation. <i>Development (Cambridge)</i> , 2009, 136, 1363-1374.	2.5	47
27	Rapamycin Induces Mitogen-activated Protein (MAP) Kinase Phosphatase-1 (MKP-1) Expression through Activation of Protein Kinase B and Mitogen-activated Protein Kinase Kinase Pathways. <i>Journal of Biological Chemistry</i> , 2013, 288, 33966-33977.	3.4	47
28	Reduced Fertility in Male Mice Deficient in the Zinc Metallopeptidase NL1. <i>Molecular and Cellular Biology</i> , 2004, 24, 4428-4437.	2.3	37
29	Defective Development of the Embryonic Liver in N-myc-Deficient Mice. <i>Developmental Biology</i> , 1998, 195, 16-28.	2.0	36
30	Epithelial inactivation of <i>Yy1</i> abrogates lung branching morphogenesis. <i>Development (Cambridge)</i> , 2015, 142, 2981-2995.	2.5	35
31	Functional redundancy of the kinases MEK1 and MEK2: Rescue of the <i>Mek1</i> mutant phenotype by <i>Mek2</i> knock-in reveals a protein threshold effect. <i>Science Signaling</i> , 2016, 9, ra9.	3.6	32
32	Anesthesia-induced hypothermia mediates decreased ARC gene and protein expression through ERK/MAPK inactivation. <i>Scientific Reports</i> , 2013, 3, 1388.	3.3	28
33	MEK1/2 Inhibition Promotes Macrophage Reparative Properties. <i>Journal of Immunology</i> , 2017, 198, 862-872.	0.8	25
34	<i>Mek1/2</i> gene dosage determines tissue response to oncogenic Ras signaling in the skin. <i>Oncogene</i> , 2009, 28, 1485-1495.	5.9	24
35	Bromodeoxyuridine resistance in CHO cells occurs in three discrete steps. <i>Somatic Cell Genetics</i> , 1982, 8, 207-222.	2.7	19
36	ERK (MAPK) does not phosphorylate tau under physiological conditions in vivo or in vitro. <i>Neurobiology of Aging</i> , 2015, 36, 901-902.	3.1	19

#	ARTICLE	IF	CITATIONS
37	MEK1 dependent and independent ERK activation regulates IL-10 and IL-12 production in bone marrow derived macrophages. Cellular Signalling, 2015, 27, 2068-2076.	3.6	19
38	<i>Mek1</i> <i>Y130C</i> mice recapitulate aspects of the human Cardio-Facio-Cutaneous syndrome. DMM Disease Models and Mechanisms, 2018, 11, .	2.4	19
39	Generation of normal lymphocytes derived from N-myc-deficient embryonic stem cells. International Immunology, 1995, 7, 1637-1647.	4.0	18
40	Lung development requires an active ERK/MAPK pathway in the lung mesenchyme. Developmental Dynamics, 2017, 246, 72-82.	1.8	18
41	N-Myc Shares Cellular Functions with c-Myc. DNA and Cell Biology, 2000, 19, 353-364.	1.9	16
42	MEK1 regulates pulmonary macrophage inflammatory responses and resolution of acute lung injury. JCI Insight, 2019, 4, .	5.0	16
43	mTOR signaling regulates gastric epithelial progenitor homeostasis and gastric tumorigenesis via MEK1-ERKs and BMP-Smad1 pathways. Cell Reports, 2021, 35, 109069.	6.4	13
44	mTOR Activation Initiates Renal Cell Carcinoma Development by Coordinating ERK and p38MAPK. Cancer Research, 2021, 81, 3174-3186.	0.9	12
45	Identification of N-myc Regulatory Regions Involved in Embryonic Expression. Pediatric Research, 2002, 51, 48-56.	2.3	10
46	Implication of MEK1 and MEK2 in the establishment of the blood-placenta barrier during placentogenesis in mouse. Reproductive BioMedicine Online, 2012, 25, 58-67.	2.4	10
47	MEK2 Negatively Regulates Lipopolysaccharide-Mediated IL-1 ² Production through HIF-1 [±] Expression. Journal of Immunology, 2019, 202, 1815-1825.	0.8	10
48	MEK/ERK Signaling in β -Cells Bifunctionally Regulates β -Cell Mass and Glucose-Stimulated Insulin Secretion Response to Maintain Glucose Homeostasis. Diabetes, 2021, 70, 1519-1535.	0.6	9
49	Analysis of Deoxycytidine (dC) Deaminase Activity in Herpes Simplex Virus-infected or HSV TK-transformed Cells: Association with Mycoplasma Contamination but Not with Virus Infection. Journal of General Virology, 1981, 57, 245-250.	2.9	8
50	Cooperative Action of Multiple <i>cis</i> -Acting Elements Is Required for N- <i>myc</i> Expression in Branchial Arches: Specific Contribution of GATA3. Molecular and Cellular Biology, 2010, 30, 5348-5363.	2.3	8
51	<i>Mek1</i> and <i>Mek2</i> Functional Redundancy in Erythropoiesis. Frontiers in Cell and Developmental Biology, 2021, 9, 639022.	3.7	5
52	MEK2 Is Essential for the Ovarian Response to the LHR Signal.. Biology of Reproduction, 2009, 81, 155-155.	2.7	5
53	Fine-tuning of MEK signaling is pivotal for limiting B and T cell activation. Cell Reports, 2022, 38, 110223.	6.4	3
54	B-RAF kinase drives developmental axon growth and promotes axon regeneration in the injured mature CNS. Journal of Cell Biology, 2014, 205, 20520IA78.	5.2	1

#	ARTICLE	IF	CITATIONS
55	MAP2K2 Delays Recovery in Murine Models of Acute Lung Injury and Associates with ARDS Outcome. American Journal of Respiratory Cell and Molecular Biology, 2022, , .	2.9	1
56	22-P002 A cooperative action of multiple cis-acting elements is required for N-myc expression in branchial arches: Specific contribution of GATA3. Mechanisms of Development, 2009, 126, S329.	1.7	0
57	Essential role of the ERK/MAPK pathway in blood-placental barrier formation. Journal of Cell Science, 2014, 127, e1-e1.	2.0	0
58	Implication des kinases MEK1 et MEK2 dans la maturation du syst�me immunitaire chez la souris. Medecine/Sciences, 2022, 38, 529-532.	0.2	0