

Linda K Medlin

List of Publications by Year in descending order

Source: <https://exaly.com/author-pdf/485525/publications.pdf>

Version: 2024-02-01

78
papers

5,897
citations

109321

35
h-index

74163

75
g-index

81
all docs

81
docs citations

81
times ranked

4789
citing authors

#	ARTICLE	IF	CITATIONS
1	Application of the 1/4Aqua microarray for pathogenic organisms across a marine/freshwater interface. <i>Harmful Algae</i> , 2020, 92, 101703.	4.8	5
2	Phylogenetic reconstruction of diatoms using a seven-gene dataset, multiple outgroups, and morphological data for a total evidence approach. <i>Phycologia</i> , 2020, 59, 422-436.	1.4	6
3	Mini review: Diatom species as seen through a molecular window. <i>Revista Brasileira De Botanica</i> , 2018, 41, 457-469.	1.3	9
4	Seasonal dynamics of freshwater pathogens as measured by microarray at Lake Sapanca, a drinking water source in the north-eastern part of Turkey. <i>Environmental Monitoring and Assessment</i> , 2018, 190, 42.	2.7	4
5	Molecular detection of harmful cyanobacteria and expression of their toxin genes in Dutch lakes using multi-probe RNA chips. <i>Harmful Algae</i> , 2018, 72, 25-35.	4.8	5
6	Multivariate analyses document host specificity, differences in the diatom metaphyton vs. epiphyton, and seasonality that structure the epiphytic diatom community. <i>Estuarine, Coastal and Shelf Science</i> , 2018, 213, 314-330.	2.1	4
7	Evolution of the diatoms: IX. Two datasets resolving monophyletic Classes of diatoms are used to explore the validity of adding short clone library sequences to the analysis. <i>European Journal of Phycology</i> , 2017, 52, 90-103.	2.0	5
8	Microarray (phylochip) analysis of freshwater pathogens at several sites along the Northern German coast transecting both estuarine and freshwaters. <i>Applied Microbiology and Biotechnology</i> , 2017, 101, 871-886.	3.6	13
9	Molecular Techniques for the Detection of Organisms in Aquatic Environments, with Emphasis on Harmful Algal Bloom Species. <i>Sensors</i> , 2017, 17, 1184.	3.8	70
10	Opinion: Can coalescent models explain deep divergences in the diatoms and argue for the acceptance of paraphyletic taxa at all taxonomic hierarchies?. <i>Nova Hedwigia</i> , 2016, 102, 107-128.	0.4	11
11	Electrochemical RNA genosensors for toxic algal species: enhancing selectivity and sensitivity. <i>Talanta</i> , 2016, 161, 560-566.	5.5	23
12	Evolution of the diatoms: major steps in their evolution and a review of the supporting molecular and morphological evidence. <i>Phycologia</i> , 2016, 55, 79-103.	1.4	70
13	A validated UPLC-MS/MS method for the surveillance of ten aquatic biotoxins in European brackish and freshwater systems. <i>Harmful Algae</i> , 2016, 55, 31-40.	4.8	53
14	Application of microarrays (phylochips) for analysis of community diversity by species identification. <i>Perspectives in Phycology</i> , 2016, 3, 93-106.	1.9	5
15	Detection of Emerging and Re-Emerging Pathogens in Surface Waters Close to an Urban Area. <i>International Journal of Environmental Research and Public Health</i> , 2015, 12, 5505-5527.	2.6	37
16	New Insights into Plagiogrammaceae (Bacillariophyta) Based on Multigene Phylogenies and Morphological Characteristics with the Description of a New Genus and Three New Species. <i>PLoS ONE</i> , 2015, 10, e0139300.	2.5	29
17	An assessment of RNA content in <i>Prymnesium parvum</i> , <i>Prymnesium polylepis</i> cf. <i>Chattonella</i> sp. and <i>Karlodinium veneficum</i> under varying environmental conditions for calibrating an RNA microarray for species detection. <i>FEMS Microbiology Ecology</i> , 2014, 88, 140-159.	2.7	8
18	Validation of the detection of <i>Pseudo-nitzschia</i> spp. using specific RNA probes tested in a microarray format: Calibration of signal based on variability of RNA content with environmental conditions. <i>Harmful Algae</i> , 2014, 37, 183-193.	4.8	10

#	ARTICLE	IF	CITATIONS
19	Validation of the detection of <i>Alexandrium</i> species using specific RNA probes tested in a microarray format: Calibration of signal using variability of RNA content with environmental conditions. <i>Harmful Algae</i> , 2014, 37, 17-27.	4.8	10
20	Evolution of the Diatoms: VIII. Re-Examination of the SSU-Rrna Gene Using Multiple Outgroups and a Cladistic Analysis of Valve Features. <i>Journal of Biodiversity Bioprospecting and Development</i> , 2014, 01, .	0.4	8
21	Molecular tools for monitoring harmful algal blooms. <i>Environmental Science and Pollution Research</i> , 2013, 20, 6683-6685.	5.3	33
22	Microarray testing for the presence of toxic algae monitoring programme in Galicia (NW Spain). <i>Environmental Science and Pollution Research</i> , 2013, 20, 6778-6793.	5.3	18
23	Note: steps taken to optimise probe specificity and signal intensity prior to field validation of the MIDTAL (Microarray for the Detection of Toxic Algae). <i>Environmental Science and Pollution Research</i> , 2013, 20, 6686-6689.	5.3	6
24	Molecular probes and microarrays for the detection of toxic algae in the genera <i>Dinophysis</i> and <i>Phalacroma</i> (Dinophyta). <i>Environmental Science and Pollution Research</i> , 2013, 20, 6733-6750.	5.3	21
25	Introduction to project MIDTAL: its methods and samples from Arcachon Bay, France. <i>Environmental Science and Pollution Research</i> , 2013, 20, 6690-6704.	5.3	37
26	Review: advances in electrochemical genosensors-based methods for monitoring blooms of toxic algae. <i>Environmental Science and Pollution Research</i> , 2013, 20, 6838-6850.	5.3	17
27	Testing a Microarray to Detect and Monitor Toxic Microalgae in Arcachon Bay in France. <i>Microarrays (Basel, Switzerland)</i> , 2013, 2, 1-23.	1.4	23
28	Genomics and Genetics of Diatoms. <i>Advances in Botanical Research</i> , 2012, 64, 245-284.	1.1	15
29	Molecular Detection, Quantification, and Diversity Evaluation of Microalgae. <i>Marine Biotechnology</i> , 2012, 14, 129-142.	2.4	96
30	Evaluation of probe orientation and effect of the digoxigenin-enzymatic label in a sandwich hybridization format to develop toxic algae biosensors. <i>Harmful Algae</i> , 2011, 10, 489-494.	4.8	17
31	Molecular phylogeny of the family Bacillariaceae based on 18S rDNA sequences: focus on freshwater <i>Nitzschia</i> of the section <i>Lanceolatae</i> . <i>Diatom Research</i> , 2011, 26, 273-291.	1.2	63
32	A Review of the Evolution of the Diatoms from the Origin of the Lineage to Their Populations. <i>Cellular Origin and Life in Extreme Habitats</i> , 2011, , 93-118.	0.3	18
33	Electrochemical performance of a DNA-based sensor device for detecting toxic algae. <i>Sensors and Actuators B: Chemical</i> , 2011, 153, 71-77.	7.8	33
34	The Permian–Triassic mass extinction forces the radiation of the modern marine phytoplankton. <i>Phycologia</i> , 2011, 50, 684-693.	1.4	14
35	Refining cryptophyte identification: matching cell fixation methods to FISH hybridisation of cryptomonads. <i>Journal of Applied Phycology</i> , 2010, 22, 725-731.	2.8	5
36	Pursuit of a natural classification of diatoms: An incorrect comparison of published data. <i>European Journal of Phycology</i> , 2010, 45, 155-166.	2.0	12

#	ARTICLE	IF	CITATIONS
37	Correctly assigning original discoveries to original authors. <i>Molecular Phylogenetics and Evolution</i> , 2009, 50, 407-408.	2.7	3
38	The <i>Phaeodactylum</i> genome reveals the evolutionary history of diatom genomes. <i>Nature</i> , 2008, 456, 239-244.	27.8	1,458
39	MOLECULAR EVIDENCE CONFIRMS SISTER RELATIONSHIP OF <i>ARDISSONEA</i> , <i>CLIMACOSPHEA</i> , AND <i>TOXARIUM</i> WITHIN THE BIPOLAR CENTRIC DIATOMS (BACILLARIOPHYTA, MEDIOPHYCEAE), AND CLADISTIC ANALYSES CONFIRM THAT EXTREMELY ELONGATED SHAPE HAS ARISEN TWICE IN THE DIATOMS. <i>Journal of Phycology</i> , 2008, 44, 1340-1348.	2.3	44
40	phylochip – a program for analysing hierarchical probe sets. <i>Molecular Ecology Resources</i> , 2008, 8, 99-102.	4.8	16
41	Colorimetric detection of the toxic dinoflagellate <i>Alexandrium minutum</i> using sandwich hybridization in a microtiter plate assay. <i>Harmful Algae</i> , 2008, 7, 137-145.	4.8	48
42	The ALEX CHIP – Development of a DNA chip for identification and monitoring of <i>Alexandrium</i> . <i>Harmful Algae</i> , 2008, 7, 485-494.	4.8	65
43	MORPHOLOGICAL AND MOLECULAR INVESTIGATIONS OF NAVICULOID DIATOMS. II. SELECTED GENERA AND FAMILIES. <i>Diatom Research</i> , 2008, 23, 283-329.	1.2	65
44	MORPHOLOGICAL AND MOLECULAR INVESTIGATIONS OF NAVICULOID DIATOMS. III. <i>HIPPODONTA</i> AND <i>NAVICULA</i> S. S. <i>Diatom Research</i> , 2008, 23, 331-347.	1.2	28
45	Feasibility of Assessing the Community Composition of Prasinophytes at the Helgoland Roads Sampling Site with a DNA Microarray. <i>Applied and Environmental Microbiology</i> , 2008, 74, 5305-5316.	3.1	37
46	Molecular assessment of phylogenetic relationships in selected species/genera in the naviculoid diatoms (Bacillariophyta). I. The genus <i>Placoneis</i> . <i>Nova Hedwigia</i> , 2007, 85, 331-352.	0.4	66
47	An assessment of cryptic genetic diversity within the <i>Cyclotella meneghiniana</i> species complex (Bacillariophyta) based on nuclear and plastid genes, and amplified fragment length polymorphisms. <i>European Journal of Phycology</i> , 2007, 42, 47-60.	2.0	58
48	Picobiliphytes: A Marine Picoplanktonic Algal Group with Unknown Affinities to Other Eukaryotes. <i>Science</i> , 2007, 315, 253-255.	12.6	202
49	CONTINUED IDEAS ON THE EVOLUTION OF SILICA METABOLISM. <i>Diatom Research</i> , 2007, 22, 217-226.	1.2	6
50	MOLECULAR PHYLOGENY OF SELECTED MEMBERS OF THE ORDER THALASSIOSIRALES (BACILLARIOPHYTA) AND EVOLUTION OF THE FULTOPORTULA1. <i>Journal of Phycology</i> , 2006, 42, 121-138.	2.3	93
51	Evolution of the diatoms: insights from fossil, biological and molecular data. <i>Phycologia</i> , 2006, 45, 361-402.	1.4	374
52	Picoeukaryotic Plankton Diversity at the Helgoland Time Series Site as Assessed by Three Molecular Methods. <i>Microbial Ecology</i> , 2006, 52, 53-71.	2.8	107
53	Automated detection and enumeration for toxic algae by solid-phase cytometry and the introduction of a new probe for <i>Prymnesium parvum</i> (Haptophyta: Prymnesiophyceae). <i>Journal of Plankton Research</i> , 2006, 28, 643-657.	1.8	39
54	DIVERSITY IN THE GENUS <i>SKELETONEMA</i> (BACILLARIOPHYCEAE). II. AN ASSESSMENT OF THE TAXONOMY OF COSTATUM-LIKE SPECIES WITH THE DESCRIPTION OF FOUR NEW SPECIES. <i>Journal of Phycology</i> , 2005, 41, 151-176.	2.3	336

#	ARTICLE	IF	CITATIONS
55	Electrochemical detection of the toxic dinoflagellate <i>Alexandrium ostenfeldii</i> with a DNA-biosensor. <i>Biosensors and Bioelectronics</i> , 2005, 20, 1349-1357.	10.1	96
56	DNA microchips for phytoplankton: The fluorescent wave of the future. <i>Nova Hedwigia</i> , 2004, 79, 321-327.	0.4	38
57	Evolution of the diatoms: V. Morphological and cytological support for the major clades and a taxonomic revision. <i>Phycologia</i> , 2004, 43, 245-270.	1.4	321
58	Using fluorescently-labelled rRNA probes for hierarchical estimation of phytoplankton diversity a mini-review. <i>Nova Hedwigia</i> , 2004, 79, 313-320.	0.4	39
59	PHYLOGENETIC POSITION OF TOXARIUM, A PENNATE-LIKE LINEAGE WITHIN CENTRIC DIATOMS (BACILLARIOPHYCEAE)1. <i>Journal of Phycology</i> , 2003, 39, 185-197.	2.3	108
60	The Application of a Molecular Clock Based on Molecular Sequences and the Fossil Record to Explain Biogeographic Distributions Within the <i>Alexandrium tamarense</i> "Species Complex" (Dinophyceae). <i>Molecular Biology and Evolution</i> , 2003, 20, 1015-1027.	8.9	179
61	WHY SILICA OR BETTER YET WHY NOT SILICA? SPECULATIONS AS TO WHY THE DIATOMS UTILISE SILICA AS THEIR CELL WALL MATERIAL. <i>Diatom Research</i> , 2002, 17, 453-459.	1.2	25
62	Oligonucleotide Probes for the Identification of Three Algal Groups by Dot Blot and Fluorescent Whole-Cell Hybridization. <i>Journal of Eukaryotic Microbiology</i> , 2000, 47, 76-84.	1.7	142
63	BOLIDOMONAS: A NEW GENUS WITH TWO SPECIES BELONGING TO A NEW ALGAL CLASS, THE BOLIDOPHYCEAE (HETEROKONTA). <i>Journal of Phycology</i> , 1999, 35, 368-381.	2.3	225
64	Phylogenetic relationships of the "golden algae"™ (haptophytes, heterokont chromophytes) and their plastids. <i>Plant Systematics and Evolution Supplementum = Entwicklungsgeschichte Und Systematik Der Pflanzen Supplementum</i> , 1997, , 187-219.	1.5	114
65	Evolution of the diatoms (Bacillariophyta). II. Nuclear-encoded small- subunit rRNA sequence comparisons confirm a paraphyletic origin for the centric diatoms. <i>Molecular Biology and Evolution</i> , 1996, 13, 67-75.	8.9	114
66	Evolution of the Diatoms (Bacillariophyta). <i>Molecular Phylogenetics and Evolution</i> , 1996, 6, 391-407.	2.7	146
67	<i>Synedropsis</i> gen. nov., a genus of araphid diatoms associated with sea ice. <i>Phycologia</i> , 1994, 33, 248-270.	1.4	63
68	Genetic differentiation among three colony-forming species of <i>Phaeocystis</i> : further evidence for the phylogeny of the Prymnesiophyta. <i>Phycologia</i> , 1994, 33, 199-212.	1.4	107
69	EVIDENCE FOR PARALLEL EVOLUTION OF FRUSTULE SHAPE IN TWO LINES OF PENNATE DIATOMS FROM THE EPIPHYTON. <i>Diatom Research</i> , 1991, 6, 109-124.	1.2	11
70	MORPHOLOGICAL AND GENETIC VARIATION WITHIN THE DIATOM SKELETONEMA COSTATUM (BACILLARIOPHYTA): EVIDENCE FOR A NEW SPECIES, SKELETONEMA PSEUDOCOSTATUM1. <i>Journal of Phycology</i> , 1991, 27, 514-524.	2.3	146
71	COMPARISON OF RESTRICTION ENDONUCLEASE SITES IN THE SMALL SUBUNIT 16S-like rRNA GENE FROM THE MAJOR GENERA OF THE FAMILY BACILLARIACEAE. <i>Diatom Research</i> , 1990, 5, 63-71.	1.2	6
72	TAXONOMIC STUDIES OF MARINE GOMPHONEMOID DIATOMS. <i>Diatom Research</i> , 1986, 1, 205-225.	1.2	50

#	ARTICLE	IF	CITATIONS
73	A LIGHT AND ELECTRON MICROSCOPIC STUDY OF THE EPIPSAMMIC DIATOM <i>CATENULA ADHAERENS</i> MERESCHKOWSKY. <i>Diatom Research</i> , 1986, 1, 283-290.	1.2	20
74	AN INVESTIGATION OF THE CELL WALL COMPONENTS OF <i>ACTINOCYCLUS SUBTILIS</i> (BACILLARIOPHYCEAE). <i>Journal of Phycology</i> , 1986, 22, 466-479.	2.3	24
75	A reappraisal of the diatom genus <i>Rhoiconeis</i> and the description of <i>Campylopyxis</i> , gen. nov.. <i>British Phycological Journal</i> , 1985, 20, 313-328.	1.2	20
76	Successional Sequences of Microbial Colonization on Three Species of Rhodophycean Macroalgae. <i>Annals of Botany</i> , 1985, 56, 399-413.	2.9	11
77	STRUCTURE, LIFE HISTORY AND SYSTEMATICS OF <i>RHOICOSPHENIA</i> (BACILLARIOPHYTA). IV. CORRELATION OF SIZE REDUCTION WITH CHANGES IN VALVE MORPHOLOGY OF <i>RH. GENUFLEXA</i> 1. <i>Journal of Phycology</i> , 1984, 20, 101-108.	2.3	26
78	Determination of the efficiency of filtration of cultures from microalgae and bacteria using hollow fiber filters. <i>Environmental Science: Water Research and Technology</i> , 0, , .	2.4	0