
## Kiyotaka Uchiyama

List of Publications by Year in descending order

Source: https://exaly.com/author-pdf/4872266/publications.pdf Version: 2024-02-01



#	Article	IF	CITATIONS
1	Efficacy of dexmedetomidine on peritoneal dialysis catheter insertion. International Urology and Nephrology, 2022, 54, 209-215.	1.4	2
2	Eosinophilic Reaction at the Time of Catheter Insertion Predicts Survival in Patients Initiating Peritoneal Dialysis. Blood Purification, 2022, 51, 355-364.	1.8	3
3	Home-Based Exercise Program Ameliorates Renal Function Decline in Patients With CKD Stage 4. Kidney International Reports, 2022, 7, 899-903.	0.8	2
4	Selfâ€assessment sheet submission rate predicts technique survival in patients initiating peritoneal dialysis. Nephrology, 2022, 27, 501-509.	1.6	2
5	Serum thymus and activation-regulated chemokine level is associated with the severity of chronic kidney disease-associated pruritus in patients undergoing peritoneal dialysis. Peritoneal Dialysis International, 2022, , 089686082210854.	2.3	1
6	High body mass index is a risk factor for transition to hemodialysis or hybrid therapy and peritoneal dialysis-related infection in Japanese patients undergoing peritoneal dialysis. International Urology and Nephrology, 2022, 54, 3193-3202.	1.4	6
7	Late Dialysis Modality Education Could Negatively Predict Peritoneal Dialysis Selection. Journal of Clinical Medicine, 2022, 11, 4042.	2.4	0
8	Sarcomatoid renal cell carcinoma with autosomal dominant polycystic kidney disease: a case report and literature review. CEN Case Reports, 2021, 10, 199-207.	0.9	2
9	The effect of trichlormethiazide in autosomal dominant polycystic kidney disease patients receiving tolvaptan: a randomized crossover controlled trial. Scientific Reports, 2021, 11, 17666.	3.3	14
10	Peritoneal dialysisâ€related peritonitis caused by <scp><i>Paracoccus yeei</i></scp> . Therapeutic Apheresis and Dialysis, 2021, 25, 715-717.	0.9	3
11	Exercise Parameters Predict Technique Survival in Patients on Peritoneal Dialysis. Blood Purification, 2021, 50, 380-389.	1.8	9
12	Effects of exercise on residual renal function in patients undergoing peritoneal dialysis: A postâ€hoc analysis of a randomized controlled trial. Therapeutic Apheresis and Dialysis, 2020, 24, 668-676.	0.9	8
13	Home-based Aerobic Exercise and Resistance Training in Peritoneal Dialysis Patients: A Randomized Controlled Trial. Scientific Reports, 2019, 9, 2632.	3.3	37
14	Long-Term Peritoneal Dialysis in 2 Patients with Takayasu's Arteritis. Peritoneal Dialysis International, 2017, 37, 122-123.	2.3	1
15	Baseline and Timeâ€Averaged Values Predicting Residual Renal Function Decline Rate in Japanese Peritoneal Dialysis Patients. Therapeutic Apheresis and Dialysis, 2017, 21, 599-605.	0.9	10
16	Importance of Neurogenic Bladder as a Cause of Drainage Failure. Peritoneal Dialysis International, 2016, 36, 232-233.	2.3	3
17	Effect of tolvaptan in a patient with autosomal dominant polycystic kidney disease after living donor liver transplantation. CEN Case Reports, 2016, 5, 227-231.	0.9	3