

Richard Bende

List of Publications by Year in descending order

Source: <https://exaly.com/author-pdf/4872357/publications.pdf>

Version: 2024-02-01

54
papers

3,696
citations

218381

26
h-index

174990

52
g-index

56
all docs

56
docs citations

56
times ranked

4040
citing authors

#	ARTICLE	IF	CITATIONS
1	Kinetics of Gag-specific cytotoxic T lymphocyte responses during the clinical course of HIV-1 infection: a longitudinal analysis of rapid progressors and long-term asymptomatics.. Journal of Experimental Medicine, 1995, 181, 1365-1372.	4.2	647
2	Expression of connective tissue growth factor in human renal fibrosis. Kidney International, 1998, 53, 853-861.	2.6	512
3	CD20 deficiency in humans results in impaired T cell-independent antibody responses. Journal of Clinical Investigation, 2010, 120, 214-222.	3.9	324
4	Among B cell non-Hodgkin's lymphomas, MALT lymphomas express a unique antibody repertoire with frequent rheumatoid factor reactivity. Journal of Experimental Medicine, 2005, 201, 1229-1241.	4.2	193
5	The hepatocyte growth factor/Met pathway controls proliferation and apoptosis in multiple myeloma. Leukemia, 2003, 17, 764-774.	3.3	145
6	Proteins encoded by open reading frames 3 and 4 of the genome of Lelystad virus (Arteriviridae) are structural proteins of the virion. Journal of Virology, 1996, 70, 4767-4772.	1.5	142
7	A mutated B cell chronic lymphocytic leukemia subset that recognizes and responds to fungi. Journal of Experimental Medicine, 2013, 210, 59-70.	4.2	132
8	IL-21 is expressed in Hodgkin lymphoma and activates STAT5: evidence that activated STAT5 is required for Hodgkin lymphomagenesis. Blood, 2008, 111, 4706-4715.	0.6	117
9	Molecular pathways in follicular lymphoma. Leukemia, 2007, 21, 18-29.	3.3	116
10	Associations between HLA Frequencies and Pathogenic Features of Human Immunodeficiency Virus Type 1 Infection in Seroconverters from the Amsterdam Cohort of Homosexual Men. Journal of Infectious Diseases, 1994, 169, 1244-1249.	1.9	114
11	Primary Follicular Lymphoma of the Small Intestine. American Journal of Pathology, 2003, 162, 105-113.	1.9	96
12	The majority of cutaneous marginal zone B-cell lymphomas expresses class-switched immunoglobulins and develops in a T-helper type 2 inflammatory environment. Blood, 2008, 112, 3355-3361.	0.6	92
13	Kinetics of Connective Tissue Growth Factor Expression during Experimental Proliferative Glomerulonephritis. Journal of the American Society of Nephrology: JASN, 2001, 12, 472-484.	3.0	82
14	Expression of activation-induced cytidine deaminase is confined to B-cell non-Hodgkin's lymphomas of germinal-center phenotype. Cancer Research, 2003, 63, 3894-8.	0.4	77
15	Chronic inflammatory disease, lymphoid tissue neogenesis and extranodal marginal zone B-cell lymphomas. Haematologica, 2009, 94, 1109-1123.	1.7	72
16	Variable heavy chain gene analysis of follicular lymphomas: correlation between heavy chain isotype expression and somatic mutation load. Blood, 2000, 95, 2922-2929.	0.6	71
17	Germinal centers in human lymph nodes contain reactivated memory B cells. Journal of Experimental Medicine, 2007, 204, 2655-2665.	4.2	60
18	Variable heavy-chain gene analysis of follicular lymphomas: subclone selection rather than clonal evolution over time. Blood, 2001, 98, 238-240.	0.6	54

#	ARTICLE	IF	CITATIONS
19	Comparison of human B cell antigen receptor complexes: membrane-expressed forms of immunoglobulin (Ig)M, IgD, and IgG are associated with structurally related heterodimers.. <i>Journal of Experimental Medicine</i> , 1992, 175, 1511-1519.	4.2	52
20	Effects of IL-4, IL-5, and IL-6 on growth and immunoglobulin production of Epstein-Barr virus-infected human B cells. <i>Cellular Immunology</i> , 1992, 143, 310-323.	1.4	40
21	A novel chronic lymphocytic leukemia subset expressing mutated IGHV3-7-encoded rheumatoid factor B-cell receptors that are functionally proficient. <i>Leukemia</i> , 2013, 27, 738-740.	3.3	36
22	Stereotypic Rheumatoid Factors That Are Frequently Expressed in Mucosa-Associated Lymphoid Tissue-Type Lymphomas Are Rare in the Labial Salivary Glands of Patients With Sjögren's Syndrome. <i>Arthritis and Rheumatology</i> , 2015, 67, 1074-1083.	2.9	36
23	Graft-versus-host-like disease complicating thymoma: Lack of AIRE expression as a cause of non-hereditary autoimmunity?. <i>Immunology Letters</i> , 2007, 114, 31-37.	1.1	34
24	Nucleocapsid protein N of Lelystad virus: expression by recombinant baculovirus, immunological properties, and suitability for detection of serum antibodies. <i>Vaccine Journal</i> , 1995, 2, 652-656.	2.6	34
25	Salivary Gland Mucosa-Associated Lymphoid Tissue-Type Lymphoma From Sjögren's Syndrome Patients in the Majority Express Rheumatoid Factors Affinity-Selected for IgG. <i>Arthritis and Rheumatology</i> , 2020, 72, 1330-1340.	2.9	30
26	The DNA Damage Response Regulates RAG1/2 Expression in Pre-B Cells through ATM-FOXO1 Signaling. <i>Journal of Immunology</i> , 2016, 197, 2918-2929.	0.4	27
27	Chlamydia psittaci-negative ocular adnexal marginal zone B-cell lymphomas have biased VH4-34 immunoglobulin gene expression and proliferate in a distinct inflammatory environment. <i>Leukemia</i> , 2012, 26, 1647-1653.	3.3	26
28	Variable heavy chain gene analysis of follicular lymphomas: correlation between heavy chain isotype expression and somatic mutation load. <i>Blood</i> , 2000, 95, 2922-9.	0.6	26
29	Follicular lymphoma with a novel t(14;18) breakpoint involving the immunoglobulin heavy chain switch mu region indicates an origin from germinal center B cells. <i>Blood</i> , 2002, 99, 716-718.	0.6	24
30	Follicular Dendritic Cells Catalyze Hepatocyte Growth Factor (HGF) Activation in the Germinal Center Microenvironment by Secreting the Serine Protease HGF Activator. <i>Journal of Immunology</i> , 2005, 175, 2807-2813.	0.4	24
31	Identification of a novel stereotypic IGHV4-59/IGHJ5-encoded B-cell receptor subset expressed by various B-cell lymphomas with high affinity rheumatoid factor activity. <i>Haematologica</i> , 2016, 101, e200-e203.	1.7	24
32	NF- κ B and AKT signaling prevent DNA damage in transformed pre-B cells by suppressing RAG1/2 expression and activity. <i>Blood</i> , 2015, 126, 1324-1335.	0.6	23
33	B-Lymphoblastic Lymphomas Evolving from Follicular Lymphomas Co-Express Surrogate Light Chains and Mutated Gamma Heavy Chains. <i>American Journal of Pathology</i> , 2016, 186, 3273-3284.	1.9	23
34	AKT signaling restrains tumor suppressive functions of FOXO transcription factors and GSK3 kinase in multiple myeloma. <i>Blood Advances</i> , 2020, 4, 4151-4164.	2.5	20
35	Acquisition of N-Glycosylation Sites in Immunoglobulin Heavy Chain Genes During Local Expansion in Parotid Salivary Glands of Primary Sjögren Patients. <i>Frontiers in Immunology</i> , 2018, 9, 491.	2.2	19
36	Antigen receptors and somatic hypermutation in B-cell chronic lymphocytic leukemia with Richter's transformation. <i>Haematologica</i> , 2006, 91, 903-11.	1.7	18

#	ARTICLE	IF	CITATIONS
37	In Situ Analysis of the Variable Heavy Chain Gene of an IgM/IgG-Expressing Follicular Lymphoma. <i>American Journal of Pathology</i> , 2002, 160, 883-891.	1.9	15
38	AID splice variants lack deaminase activity. <i>Blood</i> , 2009, 113, 1862-1864.	0.6	15
39	Repertoire Analysis of B-Cells Located in Striated Ducts of Salivary Glands of Patients With Sjögren's Syndrome. <i>Frontiers in Immunology</i> , 2020, 11, 1486.	2.2	13
40	Biased Ig λ expression in hypermutated IgD multiple myelomas does not result from receptor revision. <i>Leukemia</i> , 2002, 16, 1358-1361.	3.3	12
41	Activation-Induced Cytidine Deaminase Splice Variants Are Defective Because of the Lack of Structural Support for the Catalytic Site. <i>Journal of Immunology</i> , 2010, 184, 2487-2491.	0.4	11
42	VH gene analysis of primary cutaneous B-cell lymphomas: evidence for ongoing somatic hypermutation and isotype switching. <i>Blood</i> , 1998, 92, 3857-64.	0.6	11
43	MALT lymphoma-derived rheumatoid factors are nonpolyreactive high-affinity antibodies. <i>Blood</i> , 2010, 116, 1818-1819.	0.6	10
44	Quantitative RT-PCR analysis of activation-induced cytidine deaminase expression in tissue samples from mantle cell lymphoma and B-cell chronic lymphocytic leukemia patients. <i>Blood</i> , 2005, 105, 2997-2999.	0.6	9
45	In vitro induction of antibody secretion of primary B-cell chronic lymphocytic leukaemia cells. <i>Leukemia</i> , 2015, 29, 244-247.	3.3	8
46	Immunoglobulin diversification in B cell malignancies: internal splicing of heavy chain variable region as a by-product of somatic hypermutation. <i>Leukemia</i> , 2002, 16, 636-644.	3.3	7
47	The NEDD8-activating enzyme inhibitor MLN4924 induces DNA damage in Ph ⁺ leukemia and sensitizes for ABL kinase inhibitors. <i>Cell Cycle</i> , 2019, 18, 2307-2322.	1.3	5
48	De novo gene mutations in normal human memory B cells. <i>Leukemia</i> , 2019, 33, 1219-1230.	3.3	4
49	Rheumatoid Factor Reactivity of Expanded CD ²¹ B Cells in Patients With Sjögren's Syndrome: Comment on the Article by Glauzy et al. <i>Arthritis and Rheumatology</i> , 2019, 71, 169-170.	2.9	4
50	Analysis of Variable Heavy and Light Chain Genes in Follicular Lymphomas of Different Heavy Chain Isotype. <i>Current Topics in Microbiology and Immunology</i> , 1999, 246, 217-224.	0.7	2
51	A Major Subset of Mutated CLL Expresses Affinity-selected and Functionally Proficient Rheumatoid Factors. <i>HemaSphere</i> , 2021, 5, e550.	1.2	1
52	Identification of a Novel B-CLL Subset Expressing Mutated Stereotyped B-Cell Receptors with Specificity for Yeast Mannan. <i>Blood</i> , 2011, 118, 623-623.	0.6	1
53	Expression of the IL-6 gene induces differentiation of a human monoclonal EBV-transformed B cell line. <i>Human Antibodies</i> , 1993, 4, 124-133.	0.6	0
54	Higher-order of chronic lymphocytic leukaemia (CLL) classification: shared antigenic specificities of stereotyped B-cell receptor subsets as defined by the European Research Initiative on CLL consortium. <i>British Journal of Haematology</i> , 2021, , .	1.2	0