

Josep Franch-Nadal

List of Publications by Citations

Source: <https://exaly.com/author-pdf/4873187/josep-franch-nadal-publications-by-citations.pdf>

Version: 2024-04-23

This document has been generated based on the publications and citations recorded by exaly.com. For the latest version of this publication list, visit the link given above.

The third column is the impact factor (IF) of the journal, and the fourth column is the number of citations of the article.

137
papers

3,221
citations

27
h-index

53
g-index

172
ext. papers

4,285
ext. citations

3.8
avg, IF

4.79
L-index

#	Paper	IF	Citations
137	Prevalence of diabetes mellitus and impaired glucose regulation in Spain: the Di@bet.es Study. <i>Diabetologia</i> , 2012 , 55, 88-93	10.3	586
136	Cardiovascular Events Associated With SGLT-2 Inhibitors Versus Other Glucose-Lowering Drugs: The CVD-REAL 2 Study. <i>Journal of the American College of Cardiology</i> , 2018 , 71, 2628-2639	15.1	263
135	Risk factors associated with alterations in carotid intima-media thickness in hypertension: baseline data from the European Lacidipine Study on Atherosclerosis. <i>Journal of Hypertension</i> , 1998 , 16, 949-61	1.9	214
134	Risk of Cause-Specific Death in Individuals With Diabetes: A Competing Risks Analysis. <i>Diabetes Care</i> , 2016 , 39, 1987-1995	14.6	165
133	Control of glycemia and cardiovascular risk factors in patients with type 2 diabetes in primary care in Catalonia (Spain). <i>Diabetes Care</i> , 2012 , 35, 774-9	14.6	151
132	Prevalence of chronic kidney disease in patients with type 2 diabetes in Spain: PERCEDIME2 study. <i>BMC Nephrology</i> , 2013 , 14, 46	2.7	60
131	Direct medical costs attributable to type 2 diabetes mellitus: a population-based study in Catalonia, Spain. <i>European Journal of Health Economics</i> , 2016 , 17, 1001-1010	3.6	58
130	Glycaemic control and antidiabetic treatment trends in primary care centres in patients with type 2 diabetes mellitus during 2007-2013 in Catalonia: a population-based study. <i>BMJ Open</i> , 2016 , 6, e012463 ³		49
129	Prevalencia, diagnóstico, tratamiento y control de la hipertensión arterial en España. Resultados del estudio Di@bet.es. <i>Revista Espanola De Cardiología</i> , 2016 , 69, 572-578	1.5	47
128	Fifteen years of continuous improvement of quality care of type 2 diabetes mellitus in primary care in Catalonia, Spain. <i>International Journal of Clinical Practice</i> , 2012 , 66, 289-98	2.9	47
127	Iodine intake in the adult population. Di@bet.es study. <i>Clinical Nutrition</i> , 2012 , 31, 882-8	5.9	45
126	Clinical inertia in the treatment of hyperglycemia in type 2 diabetes patients in primary care. <i>Current Medical Research and Opinion</i> , 2013 , 29, 1495-502	2.5	42
125	Chronic kidney disease in the type 2 diabetic patients: prevalence and associated variables in a random sample of 2642 patients of a Mediterranean area. <i>BMC Nephrology</i> , 2012 , 13, 87	2.7	39
124	Chronic Kidney Disease and Diabetic Retinopathy in Patients with Type 2 Diabetes. <i>PLoS ONE</i> , 2016 , 11, e0149448	3.7	39
123	Therapeutic inertia in patients treated with two or more antidiabetics in primary care: Factors predicting intensification of treatment. <i>Diabetes, Obesity and Metabolism</i> , 2018 , 20, 103-112	6.7	38
122	Older type 2 diabetic patients are more likely to achieve glycaemic and cardiovascular risk factors targets than younger patients: analysis of a primary care database. <i>International Journal of Clinical Practice</i> , 2015 , 69, 1486-95	2.9	38
121	Long-term cardiovascular risk in type 2 diabetic compared with nondiabetic first acute myocardial infarction patients: a population-based cohort study in southern Europe. <i>Diabetes Care</i> , 2010 , 33, 2004-9 ^{14.6}	14.6	38

120	Risk of cardiovascular events and death associated with initiation of SGLT2 inhibitors compared with DPP-4 inhibitors: an analysis from the CVD-REAL 2 multinational cohort study. <i>Lancet Diabetes and Endocrinology</i> , 2020 , 8, 606-615	18.1	37
119	Modifiable risk factors associated with prediabetes in men and women: a cross-sectional analysis of the cohort study in primary health care on the evolution of patients with prediabetes (PREDAPS-Study). <i>BMC Family Practice</i> , 2015 , 16, 5	2.6	33
118	Prevalence of the metabolic syndrome in Spain using regional cutoff points for waist circumference: the di@bet.es study. <i>Acta Diabetologica</i> , 2013 , 50, 615-23	3.9	30
117	Low Physical Activity and Its Association with Diabetes and Other Cardiovascular Risk Factors: A Nationwide, Population-Based Study. <i>PLoS ONE</i> , 2016 , 11, e0160959	3.7	30
116	Olive oil has a beneficial effect on impaired glucose regulation and other cardiometabolic risk factors. Di@bet.es study. <i>European Journal of Clinical Nutrition</i> , 2013 , 67, 911-6	5.2	27
115	Validity of self-reported diabetes in health interview surveys for measuring social inequalities in the prevalence of diabetes. <i>Journal of Epidemiology and Community Health</i> , 2012 , 66, e15	5.1	27
114	Diabetic retinopathy is associated with the presence and burden of subclinical carotid atherosclerosis in type 1 diabetes. <i>Cardiovascular Diabetology</i> , 2018 , 17, 66	8.7	25
113	Prevalence of diabetic retinopathy in individuals with type 2 diabetes who had recorded diabetic retinopathy from retinal photographs in Catalonia (Spain). <i>British Journal of Ophthalmology</i> , 2015 , 99, 1628-33	5.5	25
112	Variable patterns of obesity and cardiometabolic phenotypes and their association with lifestyle factors in the Di@bet.es study. <i>Nutrition, Metabolism and Cardiovascular Diseases</i> , 2014 , 24, 947-55	4.5	24
111	Incidence of diabetes mellitus in Spain as results of the nation-wide cohort di@bet.es study. <i>Scientific Reports</i> , 2020 , 10, 2765	4.9	24
110	Fatty liver index is a predictor of incident diabetes in patients with prediabetes: The PREDAPS study. <i>PLoS ONE</i> , 2018 , 13, e0198327	3.7	23
109	Ambient temperature and prevalence of obesity in the Spanish population: The Di@bet.es study. <i>Obesity</i> , 2014 , 22, 2328-32	8	23
108	Low-grade Inflammatory Marker Profile May Help to Differentiate Patients With LADA, Classic Adult-Onset Type 1 Diabetes, and Type 2 Diabetes. <i>Diabetes Care</i> , 2018 , 41, 862-868	14.6	22
107	Cardiometabolic Risk Profiles in Patients With Impaired Fasting Glucose and/or Hemoglobin A1c 5.7% to 6.4%: Evidence for a Gradient According to Diagnostic Criteria: The PREDAPS Study. <i>Medicine (United States)</i> , 2015 , 94, e1935	1.8	22
106	Differences in the Cardiometabolic Control in Type 2 Diabetes according to Gender and the Presence of Cardiovascular Disease: Results from the eControl Study. <i>International Journal of Endocrinology</i> , 2014 , 2014, 131709	2.7	22
105	Prevalence and coprevalence of chronic comorbid conditions in patients with type 2 diabetes in Catalonia: a population-based cross-sectional study. <i>BMJ Open</i> , 2019 , 9, e031281	3	22
104	Reference values for TSH may be inadequate to define hypothyroidism in persons with morbid obesity: Di@bet.es study. <i>Obesity</i> , 2017 , 25, 788-793	8	21
103	miR-10b and miR-223-3p in serum microvesicles signal progression from prediabetes to type 2 diabetes. <i>Journal of Endocrinological Investigation</i> , 2020 , 43, 451-459	5.2	21

102	Do Patients With High Vascular Risk Take Antihypertensive Medication Correctly? Cumple-MEMS Study. <i>Revista Espanola De Cardiologia</i> , 2012 , 65, 544-50	1.5	20
101	Trends in chronic complications of type 2 diabetic patients from Spanish primary health care centres (GEDAPS study): ten year-implementation of St. Vincent recommendations. <i>Primary Care Diabetes</i> , 2012 , 6, 11-8	2.4	20
100	The Association Between Poor Glycemic Control and Health Care Costs in People With Diabetes: A Population-Based Study. <i>Diabetes Care</i> , 2020 , 43, 751-758	14.6	18
99	Mediterranean diet adherence in individuals with prediabetes and unknown diabetes: the Di@bet.es Study. <i>Annals of Nutrition and Metabolism</i> , 2013 , 62, 339-46	4.5	17
98	Barriers associated with poor control in Spanish diabetic patients. A consensus study. <i>International Journal of Clinical Practice</i> , 2013 , 67, 888-94	2.9	17
97	[Cohort Study in Primary Health Care on the Evolution of Patients with Prediabetes (PREDAPS): basis and methodology]. <i>Revista Espanola De Salud Publica</i> , 2013 , 87, 121-35	0.4	17
96	Traditional risk factors alone could not explain the excess mortality in patients with diabetes: a national cohort study of older Spanish adults. <i>Diabetes Care</i> , 2012 , 35, 2503-9	14.6	17
95	Metabolic control and cardiovascular risk factors in type 2 diabetes mellitus patients according to diabetes duration. <i>Family Practice</i> , 2015 , 32, 27-34	1.9	15
94	Prevalence, clinical features and risk assessment of pre-diabetes in Spain: the prospective Mollerussa cohort study. <i>BMJ Open</i> , 2017 , 7, e015158	3	14
93	Clinical characteristics of type 2 diabetic patients on basal insulin therapy with adequate fasting glucose control who do not achieve HbA1c targets. <i>Journal of Diabetes</i> , 2017 , 9, 34-44	3.8	14
92	Evaluation of Health-Related Quality of Life according to Carbohydrate Metabolism Status: A Spanish Population-Based Study (Di@bet.es Study). <i>International Journal of Endocrinology</i> , 2012 , 2012, 872305	2.7	14
91	Noninsulin Antidiabetic Drugs for Patients with Type 2 Diabetes Mellitus: Are We Respecting Their Contraindications?. <i>Journal of Diabetes Research</i> , 2016 , 2016, 7502489	3.9	13
90	The impact of a programme to improve quality of care for people with type 2 diabetes on hard to reach groups: The GEDAPS study. <i>Primary Care Diabetes</i> , 2015 , 9, 211-8	2.4	12
89	Vitamin D deficiency is associated with poorer satisfaction with diabetes-related treatment and quality of life in patients with type 2 diabetes: a cross-sectional study. <i>Health and Quality of Life Outcomes</i> , 2018 , 16, 44	3	12
88	Impact of chronic kidney disease on the prevalence of cardiovascular disease in patients with type 2 diabetes in Spain: PERCEDIME2 study. <i>BMC Nephrology</i> , 2014 , 15, 150	2.7	12
87	Is diabetes mellitus correctly registered and classified in primary care? A population-based study in Catalonia, Spain. <i>Endocrinologia Y Nutricion: Organo De La Sociedad Espanola De Endocrinologia Y Nutricion</i> , 2016 , 63, 440-448		12
86	Estimaci3n del riesgo cardiovascular en Espa1a seg1n la 1gu1a europea sobre prevenci3n de la 1enfermedad cardiovascular en la 1pr1ctica cl1nica. <i>Revista Espanola De Cardiologia</i> , 2015 , 68, 417-425	1.5	11
85	Iron deficiency is associated with Hypothyroxinemia and Hypotriiodothyroninemia in the Spanish general adult population: Di@bet.es study. <i>Scientific Reports</i> , 2018 , 8, 6571	4.9	11

84	Factors determining high-sensitivity C-reactive protein values in the Spanish population. Di@bet.es study. <i>European Journal of Clinical Investigation</i> , 2013 , 43, 1-10	4.6	11
83	Subclinical Carotid Atherosclerosis in Asymptomatic Subjects With Type 2 Diabetes Mellitus. <i>Journal of Cardiovascular Nursing</i> , 2016 , 31, E1-7	2.1	10
82	Association of low oleic acid intake with diabetic retinopathy in type 2 diabetic patients: a case-control study. <i>Nutrition and Metabolism</i> , 2016 , 13, 40	4.6	10
81	Non-compliance and inertia in hypertensive Spaniards at high cardiovascular risk: CUMPLE study. <i>Current Medical Research and Opinion</i> , 2014 , 30, 11-7	2.5	9
80	The diabetic immigrant: cardiovascular risk factors and control. Contributions of the IDIME study. <i>Revista Espanola De Cardiologia</i> , 2013 , 66, 39-46	1.5	9
79	Understanding the physical, social, and emotional experiences of people with uncontrolled Type 2 diabetes: a qualitative study. <i>Patient Preference and Adherence</i> , 2016 , 10, 2323-2332	2.4	9
78	Prevalence of Diabetic Foot Disease in Patients with Diabetes Mellitus under Renal Replacement Therapy in Lleida, Spain. <i>BioMed Research International</i> , 2016 , 2016, 7217586	3	9
77	Glycaemic control after treatment intensification in patients with type 2 diabetes uncontrolled on two or more non-insulin antidiabetic drugs in a real-world setting. <i>Diabetes, Obesity and Metabolism</i> , 2019 , 21, 1373-1380	6.7	9
76	How Many Patients with Type 2 Diabetes Meet the Inclusion Criteria of the Cardiovascular Outcome Trials with SGLT2 Inhibitors? Estimations from a Population Database in a Mediterranean Area. <i>Journal of Diabetes Research</i> , 2019 , 2019, 2018374	3.9	9
75	Cross-sectional study about the use of telemedicine for type 2 diabetes mellitus management in Spain: patient's perspective. The EnREDA2 Study. <i>BMJ Open</i> , 2019 , 9, e028467	3	8
74	Factors affecting levels of urinary albumin excretion in the general population of Spain: the Di@bet.es study. <i>Clinical Science</i> , 2013 , 124, 269-77	6.5	8
73	Glucagon-like peptide-1 receptor agonists in patients with type 2 diabetes: real-world evidence from a Mediterranean area. <i>Current Medical Research and Opinion</i> , 2019 , 35, 1735-1744	2.5	7
72	Prevalence of plasma lipid abnormalities and its association with glucose metabolism in Spain: the di@bet.es study. <i>Clinica E Investigaci3n En Arteriosclerosis</i> , 2014 , 26, 107-14	1.4	7
71	Ambient temperature and prevalence of diabetes and insulin resistance in the Spanish population: Di@bet.es study. <i>European Journal of Endocrinology</i> , 2019 , 180, 273-280	6.5	7
70	Saxagliptin and risk of heart failure hospitalization: Concern or miscalculation?. <i>International Journal of Cardiology</i> , 2016 , 220, 573-4	3.2	7
69	Decreased iron stores are associated with cardiovascular disease in patients with type 2 diabetes both cross-sectionally and longitudinally. <i>Atherosclerosis</i> , 2018 , 272, 193-199	3.1	6
68	Consumo de f3rmacos relacionados con el tratamiento de la diabetes mellitus y otros factores de riesgo cardiovascular en la poblaci3n espa3ola. Estudio Di@bet.es. <i>Revista Espanola De Cardiologia</i> , 2013 , 66, 854-863	1.5	6
67	Inadequate Glycaemic Control and Therapeutic Management of Adults over 65 Years Old with Type 2 Diabetes Mellitus in Spain. <i>Journal of Nutrition, Health and Aging</i> , 2017 , 21, 1365-1370	5.2	6

66	Patient-reported outcomes in type 2 diabetes mellitus: patients and primary care physicians perspectives in the Spanish health care system. <i>Patient Preference and Adherence</i> , 2015 , 9, 1413-22	2.4	6
65	Prevalence of pre-diabetes and undiagnosed diabetes in the Mollerussa prospective observational cohort study in a semi-rural area of Catalonia. <i>BMJ Open</i> , 2020 , 10, e033332	3	5
64	Glycaemic control and treatment of type 2 diabetes in adults aged 75 years or older. <i>International Journal of Clinical Practice</i> , 2018 , 72, e13075	2.9	5
63	Investigar en diabetes desde una base de datos de atención primaria: la experiencia del Sistema de Información para el Desarrollo de la Investigación en Atención Primaria (SIDiAP). <i>Avances En Diabetología</i> , 2013 , 29, 169-174		5
62	La historia clínica electrónica y los registros sobre diabetes en España. <i>Avances En Diabetología</i> , 2011 , 27, 128-136		5
61	Half of patients with type 2 diabetes mellitus are at very high cardiovascular risk according to the ESC/EASD: data from a large Mediterranean population. <i>European Journal of Preventive Cardiology</i> , 2020 ,	3.9	5
60	Knowledge management through two virtual communities of practice (Endobloc and Pneumobloc). <i>Health Informatics Journal</i> , 2017 , 23, 170-180	3	4
59	Glucagon-Like Peptide-1 Receptor Agonists in Patients with Type 2 Diabetes: Prescription According to Reimbursement Constraints and Guideline Recommendations in Catalonia. <i>Journal of Clinical Medicine</i> , 2019 , 8,	5.1	4
58	Dairy Product Consumption and Metabolic Diseases in the Di@bet.es Study. <i>Nutrients</i> , 2019 , 11,	6.7	4
57	Lifestyle factors do not explain the difference on diabetes progression according to type of prediabetes: Results from a Spanish prospective cohort of prediabetic patients. <i>Diabetes Research and Clinical Practice</i> , 2019 , 153, 66-75	7.4	4
56	Comparison of different vascular risk engines in the identification of type 2 diabetes patients with high cardiovascular risk. <i>BMC Cardiovascular Disorders</i> , 2015 , 15, 121	2.3	4
55	Impact of the American Diabetes Association diagnosis criteria on high-risk Spanish population. IGT Research Group. Impaired glucose tolerance. <i>Diabetes Research and Clinical Practice</i> , 1999 , 46, 75-81	7.4	4
54	Analysis of the location of retinal lesions in central retinographies of patients with Type 2 diabetes. <i>Acta Ophthalmologica</i> , 2020 , 98, e13-e21	3.7	4
53	Prediabetes Is Independently Associated with Subclinical Carotid Atherosclerosis: An Observational Study in a Non-Urban Mediterranean Population. <i>Journal of Clinical Medicine</i> , 2020 , 9,	5.1	4
52	Evaluation of clinical and antidiabetic treatment characteristics of different sub-groups of patients with type 2 diabetes: Data from a Mediterranean population database. <i>Primary Care Diabetes</i> , 2021 , 15, 588-595	2.4	4
51	Diabetes and tuberculosis: a syndemic complicated by COVID-19. <i>Medicina Clínica</i> , 2021 , 157, 288-293	1	4
50	High levels of fasting glucose and glycosylated hemoglobin values are associated with hyperfiltration in a Spanish prediabetes cohort. The PREDAPS Study. <i>PLoS ONE</i> , 2019 , 14, e0222848	3.7	3
49	Ultrasound Tissue Characterization of Carotid Plaques Differs Between Patients with Type 1 Diabetes and Subjects without Diabetes. <i>Journal of Clinical Medicine</i> , 2019 , 8,	5.1	3

48	Spanish People with Type 2 Diabetes Show an Improved Adherence to the Mediterranean Diet. <i>Nutrients</i> , 2020 , 12,	6.7	3
47	Adiposity and Diabetes Risk in Adults with Prediabetes: Heterogeneity of Findings Depending on Age and Anthropometric Measure. <i>Obesity</i> , 2018 , 26, 1481-1490	8	3
46	Estudio observacional transversal en consultas de atención primaria y atención hospitalaria del perfil clínico del paciente con diabetes mellitus tipo 2 no controlado en fase inicial de tratamiento (estudio EUPHORIA). <i>Avances En Diabetología</i> , 2011 , 27, 69-77		3
45	Improving management of glycaemic control in people with T2DM in primary care: estimation of the impact on the clinical complications and associated costs. <i>BMC Health Services Research</i> , 2020 , 20, 803	2.9	3
44	Cardiovascular outcomes with sodium-glucose cotransporter-2 inhibitors vs other glucose-lowering drugs in 13 countries across three continents: analysis of CVD-REAL data. <i>Cardiovascular Diabetology</i> , 2021 , 20, 159	8.7	3
43	The Circulating Fatty Acid Transporter Soluble CD36 Is Not Associated with Carotid Atherosclerosis in Subjects with Type 1 and Type 2 Diabetes Mellitus. <i>Journal of Clinical Medicine</i> , 2020 , 9,	5.1	2
42	Prevalence and control of type 2 diabetes mellitus among primary care physicians in Spain. PRISMA Study. <i>Endocrinología, Diabetes Y Nutrición</i> , 2017 , 64, 265-271	1.3	2
41	El control metabólico en pacientes diabéticos inmigrantes. <i>Avances En Diabetología</i> , 2010 , 26, 107-111		2
40	Anthropologic study of immigrant patients with T2DM from Morocco to Spain. <i>European Diabetes Nursing</i> , 2010 , 7, 24-28		2
39	Therapeutic Inertia: Still a Long Way to Go That Cannot Be Postponed. <i>Diabetes Spectrum</i> , 2020 , 33, 50-57.	5.9	2
38	Atherogenic dyslipidemia, but not hyperglycemia, is an independent factor associated with liver fibrosis in subjects with type 2 diabetes and NAFLD: a population-based study. <i>European Journal of Endocrinology</i> , 2021 , 184, 587-596	6.5	2
37	Perfil lipoproteico avanzado en individuos con metabolismo glucémico normal y alterado. <i>Revista Espanola De Cardiología</i> , 2021 , 75, 22-22	1.5	2
36	Analysis of the Adherence and Safety of Second Oral Glucose-Lowering Therapy in Routine Practice From the Mediterranean Area: A Retrospective Cohort Study. <i>Frontiers in Endocrinology</i> , 2021 , 12, 708372	5.7	2
35	Short-term foot complications in Charcot neuroarthropathy: A retrospective study in tertiary care centres in Spain. <i>Endocrinología, Diabetes Y Nutrición</i> , 2018 , 65, 479-485	1.3	2
34	Cardiovascular and mortality benefits of sodium-glucose co-transporter-2 inhibitors in patients with type 2 diabetes mellitus: CVD-Real Catalonia. <i>Cardiovascular Diabetology</i> , 2021 , 20, 139	8.7	2
33	Epidemiological multicentre study on the education provided to patients with type 2 diabetes mellitus in the Spanish Health Care System. The Forma2 study. <i>Semergen</i> , 2018 , 44, 5-12	1.9	1
32	Treatment of Hyperglycaemia in Type 2 Diabetic Patients in a Primary Care Population Database in a Mediterranean Area (Catalonia, Spain). <i>Journal of Diabetes & Metabolism</i> , 2014 , 05,	0	1
31	Do Patients With High Vascular Risk Take Antihypertensive Medication Correctly? Cumple-MEMS Study. <i>Revista Espanola De Cardiología (English Ed)</i> , 2012 , 65, 544-550	0.7	1

30	Inercia e incumplimiento a partes iguales. <i>Hipertension Y Riesgo Vascular</i> , 2012 , 29, 1-3	0.5	1
29	Trends in the Degree of Control and Treatment of Cardiovascular Risk Factors in People With Type 2 Diabetes in a Primary Care Setting in Catalonia During 2007-2018.. <i>Frontiers in Endocrinology</i> , 2021 , 12, 810757	5.7	1
28	Therapeutic inertia in patients treated with two or more antidiabetics in primary care: Factors predicting intensification of treatment 2018 , 20, 103		1
27	Incidence and regression of metabolic syndrome in a representative sample of the Spanish population: results of the cohort di@bet.es study. <i>BMJ Open Diabetes Research and Care</i> , 2020 , 8,	4.5	1
26	Circulating CD5L is associated with cardiovascular events and all-cause mortality in individuals with chronic kidney disease. <i>Aging</i> , 2021 , 13, 22690-22709	5.6	1
25	Analysis of the effectiveness of second oral glucose-lowering therapy in routine clinical practice from the mediterranean area: A retrospective cohort study. <i>Diabetes Research and Clinical Practice</i> , 2021 , 171, 108616	7.4	1
24	Advanced lipoprotein profile in individuals with normal and impaired glucose metabolism. <i>Revista Espanola De Cardiologia (English Ed)</i> , 2021 , 75, 22-22	0.7	1
23	Incidence and regression of metabolic syndrome in a representative sample of the Spanish population: results of the cohort di@bet.es study 2020 , 8, e001715		1
22	Use of drugs related to the treatment of diabetes mellitus and other cardiovascular risk factors in the Spanish population. The Di@bet.es study. <i>Revista Espanola De Cardiologia (English Ed)</i> , 2013 , 66, 854-63	0.7	0
21	Adding questions about cardiovascular risk factors improve the ability of the ADA questionnaire to identify unknown diabetic patients in Spain. <i>Avances En Diabetologia</i> , 2010 , 26, 347-352		0
20	Associations Between Diabetic Retinopathy and Parkinson Disease: Results From the Catalanian Primary Care Cohort Study.. <i>Frontiers in Medicine</i> , 2021 , 8, 800973	4.9	0
19	The heterogeneity of reversion to normoglycemia according to prediabetes type is not explained by lifestyle factors. <i>Scientific Reports</i> , 2021 , 11, 9667	4.9	0
18	Risk factors for severe outcomes in people with diabetes hospitalised for COVID-19: a cross-sectional database study. <i>BMJ Open</i> , 2021 , 11, e051237	3	0
17	Physicians and Pharmacists Clinical Considerations for Elderly Patients with Type 2 Diabetes Mellitus: The IMPLICA2 Study. <i>Clinical Drug Investigation</i> , 2019 , 39, 73-84	3.2	0
16	Iodine Deficiency and Mortality in Spanish Adults: Di@bet.es Study. <i>Thyroid</i> , 2021 , 31, 106-114	6.2	0
15	Fatty liver index as a predictor for type 2 diabetes in subjects with normoglycemia in a nationwide cohort study. <i>Scientific Reports</i> , 2021 , 11, 16453	4.9	0
14	Incidence and characteristics of diabetic foot ulcers in subjects with type 2 diabetes in Catalanian primary care centres: An observational multicentre study. <i>Primary Care Diabetes</i> , 2021 , 15, 1033-1039	2.4	0
13	Diabetes and tuberculosis: A syndemic complicated by COVID-19. <i>Medicina Clínica (English Edition)</i> , 2021 , 157, 288-293	0.3	0

12	Mediterranean diet and diabetes risk in a cohort study of individuals with prediabetes: propensity score analyses.. <i>Diabetic Medicine</i> , 2021 , e14768	3.5	o
11	Heterogeneity in the association between prediabetes categories and reduction on glomerular filtration rate in a 5-year follow-up.. <i>Scientific Reports</i> , 2022 , 12, 7373	4.9	o
10	Prevalence and control of type 2 diabetes mellitus among primary care physicians in Spain. PRISMA Study. <i>Endocrinología Diabetes Y Nutrición (English Ed)</i> , 2017 , 64, 265-271	0.1	
9	Efectos hipoglucemiantes de las resinas de intercambio. Del secuestro de los ácidos biliares a la activación del metabolismo de la glucosa. <i>Clínica E Investigación En Arteriosclerosis</i> , 2011 , 23, 3-8	1.4	
8	Eficacia de los fármacos hipolipemiantes: un metaanálisis. <i>FMC Formación Médica Continuada En Atención Primaria</i> , 2006 , 13, 650	0	
7	Diabetes Treatment in Patients with Renal DiseaseHow Many Are Receiving Contraindicated Drugs?. <i>Diabetes</i> , 2018 , 67, 1647-P	0.9	
6	Multiple Chronic Comorbidities in a T2DM Mediterranean Population. <i>Diabetes</i> , 2018 , 67, 1646-P	0.9	
5	Compliance with the DPP-4 inhibitors dose adjustment recommendations based on renal function in a population database. <i>Endocrinología, Diabetes Y Nutrición</i> , 2021 , 69, 83-83	1.3	
4	Análisis de coste-efectividad de saxagliptina como tratamiento triple oral (con metformina y una sulfonilurea) en el manejo de la diabetes tipo 2 en España. <i>Pharmacoeconomics - Spanish Research Articles</i> , 2016 , 13, 25-35		
3	Classifying and communicating risks in prediabetes according to fasting glucose and/or glycated hemoglobin: PREDAPS cohort study. <i>Scandinavian Journal of Primary Health Care</i> , 2021 , 39, 355-363	2.7	
2	Compliance with the DPP-4 inhibitors dose adjustment recommendations based on renal function in a population database.. <i>Endocrinología Diabetes Y Nutrición (English Ed)</i> , 2022 , 69, 83-91	0.1	
1	The Cost-Effectiveness of Oral Semaglutide in Spain: A Long-Term Health Economic Analysis Based on the PIONEER Clinical Trials.. <i>Advances in Therapy</i> , 2022 , 1	4.1	