

Josep Franch-Nadal

List of Publications by Year in Descending Order

Source: <https://exaly.com/author-pdf/4873187/josep-franch-nadal-publications-by-year.pdf>

Version: 2024-04-23

This document has been generated based on the publications and citations recorded by exaly.com. For the latest version of this publication list, visit the link given above.

The third column is the impact factor (IF) of the journal, and the fourth column is the number of citations of the article.

137
papers

3,221
citations

27
h-index

53
g-index

172
ext. papers

4,285
ext. citations

3.8
avg, IF

4.79
L-index

#	Paper	IF	Citations
137	Compliance with the DPP-4 inhibitors dose adjustment recommendations based on renal function in a population database.. <i>Endocrinología Diabetes Y Nutrición (English Ed)</i> , 2022 , 69, 83-91	0.1	0
136	Heterogeneity in the association between prediabetes categories and reduction on glomerular filtration rate in a 5-year follow-up.. <i>Scientific Reports</i> , 2022 , 12, 7373	4.9	0
135	The Cost-Effectiveness of Oral Semaglutide in Spain: A Long-Term Health Economic Analysis Based on the PIONEER Clinical Trials.. <i>Advances in Therapy</i> , 2022 , 1	4.1	0
134	Associations Between Diabetic Retinopathy and Parkinson's Disease: Results From the Catalanian Primary Care Cohort Study.. <i>Frontiers in Medicine</i> , 2021 , 8, 800973	4.9	0
133	Trends in the Degree of Control and Treatment of Cardiovascular Risk Factors in People With Type 2 Diabetes in a Primary Care Setting in Catalonia During 2007-2018.. <i>Frontiers in Endocrinology</i> , 2021 , 12, 810757	5.7	1
132	Circulating CD5L is associated with cardiovascular events and all-cause mortality in individuals with chronic kidney disease. <i>Aging</i> , 2021 , 13, 22690-22709	5.6	1
131	Atherogenic dyslipidemia, but not hyperglycemia, is an independent factor associated with liver fibrosis in subjects with type 2 diabetes and NAFLD: a population-based study. <i>European Journal of Endocrinology</i> , 2021 , 184, 587-596	6.5	2
130	Perfil lipoproteico avanzado en individuos con metabolismo glucémico normal y alterado. <i>Revista Espanola De Cardiologia</i> , 2021 , 75, 22-22	1.5	2
129	The heterogeneity of reversion to normoglycemia according to prediabetes type is not explained by lifestyle factors. <i>Scientific Reports</i> , 2021 , 11, 9667	4.9	0
128	Compliance with the DPP-4 inhibitors dose adjustment recommendations based on renal function in a population database. <i>Endocrinología, Diabetes Y Nutrición</i> , 2021 , 69, 83-83	1.3	0
127	Evaluation of clinical and antidiabetic treatment characteristics of different sub-groups of patients with type 2 diabetes: Data from a Mediterranean population database. <i>Primary Care Diabetes</i> , 2021 , 15, 588-595	2.4	4
126	Cardiovascular outcomes with sodium-glucose cotransporter-2 inhibitors vs other glucose-lowering drugs in 13 countries across three continents: analysis of CVD-REAL data. <i>Cardiovascular Diabetology</i> , 2021 , 20, 159	8.7	3
125	Risk factors for severe outcomes in people with diabetes hospitalised for COVID-19: a cross-sectional database study. <i>BMJ Open</i> , 2021 , 11, e051237	3	0
124	Analysis of the Adherence and Safety of Second Oral Glucose-Lowering Therapy in Routine Practice From the Mediterranean Area: A Retrospective Cohort Study. <i>Frontiers in Endocrinology</i> , 2021 , 12, 708372	5.7	2
123	Iodine Deficiency and Mortality in Spanish Adults: Di@bet.es Study. <i>Thyroid</i> , 2021 , 31, 106-114	6.2	0
122	Analysis of the effectiveness of second oral glucose-lowering therapy in routine clinical practice from the mediterranean area: A retrospective cohort study. <i>Diabetes Research and Clinical Practice</i> , 2021 , 171, 108616	7.4	1
121	Advanced lipoprotein profile in individuals with normal and impaired glucose metabolism. <i>Revista Espanola De Cardiologia (English Ed)</i> , 2021 , 75, 22-22	0.7	1

120	Classifying and communicating risks in prediabetes according to fasting glucose and/or glycated hemoglobin: PREDAPS cohort study. <i>Scandinavian Journal of Primary Health Care</i> , 2021 , 39, 355-363	2.7	
119	Cardiovascular and mortality benefits of sodium-glucose co-transporter-2 inhibitors in patients with type 2 diabetes mellitus: CVD-Real Catalonia. <i>Cardiovascular Diabetology</i> , 2021 , 20, 139	8.7	2
118	Fatty liver index as a predictor for type 2 diabetes in subjects with normoglycemia in a nationwide cohort study. <i>Scientific Reports</i> , 2021 , 11, 16453	4.9	0
117	Incidence and characteristics of diabetic foot ulcers in subjects with type 2 diabetes in Catalanian primary care centres: An observational multicentre study. <i>Primary Care Diabetes</i> , 2021 , 15, 1033-1039	2.4	0
116	Diabetes and tuberculosis: a syndemic complicated by COVID-19. <i>Medicina Clínica</i> , 2021 , 157, 288-293	1	4
115	Diabetes and tuberculosis: A syndemic complicated by COVID-19. <i>Medicina Clínica (English Edition)</i> , 2021 , 157, 288-293	0.3	0
114	Mediterranean diet and diabetes risk in a cohort study of individuals with prediabetes: propensity score analyses.. <i>Diabetic Medicine</i> , 2021 , e14768	3.5	0
113	The Circulating Fatty Acid Transporter Soluble CD36 Is Not Associated with Carotid Atherosclerosis in Subjects with Type 1 and Type 2 Diabetes Mellitus. <i>Journal of Clinical Medicine</i> , 2020 , 9,	5.1	2
112	Risk of cardiovascular events and death associated with initiation of SGLT2 inhibitors compared with DPP-4 inhibitors: an analysis from the CVD-REAL 2 multinational cohort study. <i>Lancet Diabetes and Endocrinology</i> , 2020 , 8, 606-615	18.1	37
111	Prevalence of pre-diabetes and undiagnosed diabetes in the Mollerussa prospective observational cohort study in a semi-rural area of Catalonia. <i>BMJ Open</i> , 2020 , 10, e033332	3	5
110	The Association Between Poor Glycemic Control and Health Care Costs in People With Diabetes: A Population-Based Study. <i>Diabetes Care</i> , 2020 , 43, 751-758	14.6	18
109	Spanish People with Type 2 Diabetes Show an Improved Adherence to the Mediterranean Diet. <i>Nutrients</i> , 2020 , 12,	6.7	3
108	Half of patients with type 2 diabetes mellitus are at very high cardiovascular risk according to the ESC/EASD: data from a large Mediterranean population. <i>European Journal of Preventive Cardiology</i> , 2020 ,	3.9	5
107	Incidence and regression of metabolic syndrome in a representative sample of the Spanish population: results of the cohort di@bet.es study. <i>BMJ Open Diabetes Research and Care</i> , 2020 , 8,	4.5	1
106	Therapeutic Inertia: Still a Long Way to Go That Cannot Be Postponed. <i>Diabetes Spectrum</i> , 2020 , 33, 50-57.9		2
105	Incidence of diabetes mellitus in Spain as results of the nation-wide cohort di@bet.es study. <i>Scientific Reports</i> , 2020 , 10, 2765	4.9	24
104	Analysis of the location of retinal lesions in central retinographies of patients with Type 2 diabetes. <i>Acta Ophthalmologica</i> , 2020 , 98, e13-e21	3.7	4
103	miR-10b and miR-223-3p in serum microvesicles signal progression from prediabetes to type 2 diabetes. <i>Journal of Endocrinological Investigation</i> , 2020 , 43, 451-459	5.2	21

102	Prediabetes Is Independently Associated with Subclinical Carotid Atherosclerosis: An Observational Study in a Non-Urban Mediterranean Population. <i>Journal of Clinical Medicine</i> , 2020 , <i>9</i> ,	5.1	4
101	Improving management of glycaemic control in people with T2DM in primary care: estimation of the impact on the clinical complications and associated costs. <i>BMC Health Services Research</i> , 2020 , <i>20</i> , 803	2.9	3
100	Incidence and regression of metabolic syndrome in a representative sample of the Spanish population: results of the cohort di@bet.es study 2020 , <i>8</i> , e001715		1
99	Glucagon-Like Peptide-1 Receptor Agonists in Patients with Type 2 Diabetes: Prescription According to Reimbursement Constraints and Guideline Recommendations in Catalonia. <i>Journal of Clinical Medicine</i> , 2019 , <i>8</i> ,	5.1	4
98	High levels of fasting glucose and glycosylated hemoglobin values are associated with hyperfiltration in a Spanish prediabetes cohort. The PREDAPS Study. <i>PLoS ONE</i> , 2019 , <i>14</i> , e0222848	3.7	3
97	Dairy Product Consumption and Metabolic Diseases in the Di@bet.es Study. <i>Nutrients</i> , 2019 , <i>11</i> ,	6.7	4
96	Cross-sectional study about the use of telemedicine for type 2 diabetes mellitus management in Spain: patient's perspective. The EnREDA2 Study. <i>BMJ Open</i> , 2019 , <i>9</i> , e028467	3	8
95	Lifestyle factors do not explain the difference on diabetes progression according to type of prediabetes: Results from a Spanish prospective cohort of prediabetic patients. <i>Diabetes Research and Clinical Practice</i> , 2019 , <i>153</i> , 66-75	7.4	4
94	Glucagon-like peptide-1 receptor agonists in patients with type 2 diabetes: real-world evidence from a Mediterranean area. <i>Current Medical Research and Opinion</i> , 2019 , <i>35</i> , 1735-1744	2.5	7
93	Ultrasound Tissue Characterization of Carotid Plaques Differs Between Patients with Type 1 Diabetes and Subjects without Diabetes. <i>Journal of Clinical Medicine</i> , 2019 , <i>8</i> ,	5.1	3
92	Ambient temperature and prevalence of diabetes and insulin resistance in the Spanish population: Di@bet.es study. <i>European Journal of Endocrinology</i> , 2019 , <i>180</i> , 273-280	6.5	7
91	Glycaemic control after treatment intensification in patients with type 2 diabetes uncontrolled on two or more non-insulin antidiabetic drugs in a real-world setting. <i>Diabetes, Obesity and Metabolism</i> , 2019 , <i>21</i> , 1373-1380	6.7	9
90	Prevalence and coprevalence of chronic comorbid conditions in patients with type 2 diabetes in Catalonia: a population-based cross-sectional study. <i>BMJ Open</i> , 2019 , <i>9</i> , e031281	3	22
89	How Many Patients with Type 2 Diabetes Meet the Inclusion Criteria of the Cardiovascular Outcome Trials with SGLT2 Inhibitors? Estimations from a Population Database in a Mediterranean Area. <i>Journal of Diabetes Research</i> , 2019 , <i>2019</i> , 2018374	3.9	9
88	Physicians and Pharmacists' Clinical Considerations for Elderly Patients with Type 2 Diabetes Mellitus: The IMPLICA2 Study. <i>Clinical Drug Investigation</i> , 2019 , <i>39</i> , 73-84	3.2	0
87	Low-grade Inflammatory Marker Profile May Help to Differentiate Patients With LADA, Classic Adult-Onset Type 1 Diabetes, and Type 2 Diabetes. <i>Diabetes Care</i> , 2018 , <i>41</i> , 862-868	14.6	22
86	Iron deficiency is associated with Hypothyroxinemia and Hypotriiodothyroninemia in the Spanish general adult population: Di@bet.es study. <i>Scientific Reports</i> , 2018 , <i>8</i> , 6571	4.9	11
85	Cardiovascular Events Associated With SGLT-2 Inhibitors Versus Other Glucose-Lowering Drugs: The CVD-REAL 2 Study. <i>Journal of the American College of Cardiology</i> , 2018 , <i>71</i> , 2628-2639	15.1	263

84	Decreased iron stores are associated with cardiovascular disease in patients with type 2 diabetes both cross-sectionally and longitudinally. <i>Atherosclerosis</i> , 2018 , 272, 193-199	3.1	6
83	Glycaemic control and treatment of type 2 diabetes in adults aged 75 years or older. <i>International Journal of Clinical Practice</i> , 2018 , 72, e13075	2.9	5
82	Therapeutic inertia in patients treated with two or more antidiabetics in primary care: Factors predicting intensification of treatment. <i>Diabetes, Obesity and Metabolism</i> , 2018 , 20, 103-112	6.7	38
81	Epidemiological multicentre study on the education provided to patients with type 2 diabetes mellitus in the Spanish Health Care System. The Forma2 study. <i>Semergen</i> , 2018 , 44, 5-12	1.9	1
80	Adiposity and Diabetes Risk in Adults with Prediabetes: Heterogeneity of Findings Depending on Age and Anthropometric Measure. <i>Obesity</i> , 2018 , 26, 1481-1490	8	3
79	Diabetic retinopathy is associated with the presence and burden of subclinical carotid atherosclerosis in type 1 diabetes. <i>Cardiovascular Diabetology</i> , 2018 , 17, 66	8.7	25
78	Vitamin D deficiency is associated with poorer satisfaction with diabetes-related treatment and quality of life in patients with type 2 diabetes: a cross-sectional study. <i>Health and Quality of Life Outcomes</i> , 2018 , 16, 44	3	12
77	Fatty liver index is a predictor of incident diabetes in patients with prediabetes: The PREDAPS study. <i>PLoS ONE</i> , 2018 , 13, e0198327	3.7	23
76	Therapeutic inertia in patients treated with two or more antidiabetics in primary care: Factors predicting intensification of treatment 2018 , 20, 103		1
75	Diabetes Treatment in Patients with Renal Disease How Many Are Receiving Contraindicated Drugs?. <i>Diabetes</i> , 2018 , 67, 1647-P	0.9	
74	Multiple Chronic Comorbidities in a T2DM Mediterranean Population. <i>Diabetes</i> , 2018 , 67, 1646-P	0.9	
73	Short-term foot complications in Charcot neuroarthropathy: A retrospective study in tertiary care centres in Spain. <i>Endocrinología, Diabetes Y Nutrición</i> , 2018 , 65, 479-485	1.3	2
72	Knowledge management through two virtual communities of practice (Endobloc and Pneumobloc). <i>Health Informatics Journal</i> , 2017 , 23, 170-180	3	4
71	Reference values for TSH may be inadequate to define hypothyroidism in persons with morbid obesity: Di@bet.es study. <i>Obesity</i> , 2017 , 25, 788-793	8	21
70	Prevalence, clinical features and risk assessment of pre-diabetes in Spain: the prospective Mollerussa cohort study. <i>BMJ Open</i> , 2017 , 7, e015158	3	14
69	Prevalence and control of type 2 diabetes mellitus among primary care physicians in Spain. PRISMA Study. <i>Endocrinología, Diabetes Y Nutrición</i> , 2017 , 64, 265-271	1.3	2
68	Inadequate Glycaemic Control and Therapeutic Management of Adults over 65 Years Old with Type 2 Diabetes Mellitus in Spain. <i>Journal of Nutrition, Health and Aging</i> , 2017 , 21, 1365-1370	5.2	6
67	Prevalence and control of type 2 diabetes mellitus among primary care physicians in Spain. PRISMA Study. <i>Endocrinología Diabetes Y Nutrición (English Ed)</i> , 2017 , 64, 265-271	0.1	

66	Clinical characteristics of type 2 diabetic patients on basal insulin therapy with adequate fasting glucose control who do not achieve HbA1c targets. <i>Journal of Diabetes</i> , 2017 , 9, 34-44	3.8	14
65	Glycaemic control and antidiabetic treatment trends in primary care centres in patients with type 2 diabetes mellitus during 2007-2013 in Catalonia: a population-based study. <i>BMJ Open</i> , 2016 , 6, e012463 ³		49
64	Prevalencia, diagnóstico, tratamiento y control de la hipertensión arterial en España. Resultados del estudio Di@bet.es. <i>Revista Espanola De Cardiologia</i> , 2016 , 69, 572-578	1.5	47
63	Direct medical costs attributable to type 2 diabetes mellitus: a population-based study in Catalonia, Spain. <i>European Journal of Health Economics</i> , 2016 , 17, 1001-1010	3.6	58
62	Chronic Kidney Disease and Diabetic Retinopathy in Patients with Type 2 Diabetes. <i>PLoS ONE</i> , 2016 , 11, e0149448	3.7	39
61	Understanding the physical, social, and emotional experiences of people with uncontrolled Type 2 diabetes: a qualitative study. <i>Patient Preference and Adherence</i> , 2016 , 10, 2323-2332	2.4	9
60	Prevalence of Diabetic Foot Disease in Patients with Diabetes Mellitus under Renal Replacement Therapy in Lleida, Spain. <i>BioMed Research International</i> , 2016 , 2016, 7217586	3	9
59	Noninsulin Antidiabetic Drugs for Patients with Type 2 Diabetes Mellitus: Are We Respecting Their Contraindications?. <i>Journal of Diabetes Research</i> , 2016 , 2016, 7502489	3.9	13
58	Low Physical Activity and Its Association with Diabetes and Other Cardiovascular Risk Factors: A Nationwide, Population-Based Study. <i>PLoS ONE</i> , 2016 , 11, e0160959	3.7	30
57	Subclinical Carotid Atherosclerosis in Asymptomatic Subjects With Type 2 Diabetes Mellitus. <i>Journal of Cardiovascular Nursing</i> , 2016 , 31, E1-7	2.1	10
56	Saxagliptin and risk of heart failure hospitalization: Concern or miscalculation?. <i>International Journal of Cardiology</i> , 2016 , 220, 573-4	3.2	7
55	Análisis de coste-efectividad de saxagliptina como tratamiento triple oral (con metformina y una sulfonilurea) en el manejo de la diabetes tipo 2 en España. <i>Pharmacoeconomics - Spanish Research Articles</i> , 2016 , 13, 25-35		
54	Association of low oleic acid intake with diabetic retinopathy in type 2 diabetic patients: a case-control study. <i>Nutrition and Metabolism</i> , 2016 , 13, 40	4.6	10
53	Is diabetes mellitus correctly registered and classified in primary care? A population-based study in Catalonia, Spain. <i>Endocrinología Y Nutrición: Organo De La Sociedad Espanola De Endocrinología Y Nutrición</i> , 2016 , 63, 440-448		12
52	Risk of Cause-Specific Death in Individuals With Diabetes: A Competing Risks Analysis. <i>Diabetes Care</i> , 2016 , 39, 1987-1995	14.6	165
51	Estimación del riesgo cardiovascular en España según la guía europea sobre prevención de la enfermedad cardiovascular en la práctica clínica. <i>Revista Espanola De Cardiologia</i> , 2015 , 68, 417-425	1.5	11
50	Modifiable risk factors associated with prediabetes in men and women: a cross-sectional analysis of the cohort study in primary health care on the evolution of patients with prediabetes (PREDAAPS-Study). <i>BMC Family Practice</i> , 2015 , 16, 5	2.6	33
49	The impact of a programme to improve quality of care for people with type 2 diabetes on hard to reach groups: The GEDAPS study. <i>Primary Care Diabetes</i> , 2015 , 9, 211-8	2.4	12

48	Metabolic control and cardiovascular risk factors in type 2 diabetes mellitus patients according to diabetes duration. <i>Family Practice</i> , 2015 , 32, 27-34	1.9	15
47	Older type 2 diabetic patients are more likely to achieve glycaemic and cardiovascular risk factors targets than younger patients: analysis of a primary care database. <i>International Journal of Clinical Practice</i> , 2015 , 69, 1486-95	2.9	38
46	Comparison of different vascular risk engines in the identification of type 2 diabetes patients with high cardiovascular risk. <i>BMC Cardiovascular Disorders</i> , 2015 , 15, 121	2.3	4
45	Cardiometabolic Risk Profiles in Patients With Impaired Fasting Glucose and/or Hemoglobin A1c 5.7% to 6.4%: Evidence for a Gradient According to Diagnostic Criteria: The PREDAPS Study. <i>Medicine (United States)</i> , 2015 , 94, e1935	1.8	22
44	Patient-reported outcomes in type 2 diabetes mellitus: patients and primary care physicians' perspectives in the Spanish health care system. <i>Patient Preference and Adherence</i> , 2015 , 9, 1413-22	2.4	6
43	Prevalence of diabetic retinopathy in individuals with type 2 diabetes who had recorded diabetic retinopathy from retinal photographs in Catalonia (Spain). <i>British Journal of Ophthalmology</i> , 2015 , 99, 1628-33	5.5	25
42	Non-compliance and inertia in hypertensive Spaniards at high cardiovascular risk: CUMPLE study. <i>Current Medical Research and Opinion</i> , 2014 , 30, 11-7	2.5	9
41	Variable patterns of obesity and cardiometabolic phenotypes and their association with lifestyle factors in the Di@bet.es study. <i>Nutrition, Metabolism and Cardiovascular Diseases</i> , 2014 , 24, 947-55	4.5	24
40	Treatment of Hyperglycaemia in Type 2 Diabetic Patients in a Primary Care Population Database in a Mediterranean Area (Catalonia, Spain). <i>Journal of Diabetes & Metabolism</i> , 2014 , 05,	0	1
39	Ambient temperature and prevalence of obesity in the Spanish population: The Di@bet.es study. <i>Obesity</i> , 2014 , 22, 2328-32	8	23
38	Differences in the Cardiometabolic Control in Type 2 Diabetes according to Gender and the Presence of Cardiovascular Disease: Results from the eControl Study. <i>International Journal of Endocrinology</i> , 2014 , 2014, 131709	2.7	22
37	Impact of chronic kidney disease on the prevalence of cardiovascular disease in patients with type 2 diabetes in Spain: PERCEDIME2 study. <i>BMC Nephrology</i> , 2014 , 15, 150	2.7	12
36	Prevalence of plasma lipid abnormalities and its association with glucose metabolism in Spain: the di@bet.es study. <i>Clinica E Investigaci3n En Arteriosclerosis</i> , 2014 , 26, 107-14	1.4	7
35	Prevalence of chronic kidney disease in patients with type 2 diabetes in Spain: PERCEDIME2 study. <i>BMC Nephrology</i> , 2013 , 14, 46	2.7	60
34	Prevalence of the metabolic syndrome in Spain using regional cutoff points for waist circumference: the di@bet.es study. <i>Acta Diabetologica</i> , 2013 , 50, 615-23	3.9	30
33	Investigar en diabetes desde una base de datos de atenci3n primaria: la experiencia del Sistema de Informaci3n para el Desarrollo de la Investigaci3n en Atenci3n Primaria (SIDIAP). <i>Avances En Diabetolog3a</i> , 2013 , 29, 169-174		5
32	Clinical inertia in the treatment of hyperglycemia in type 2 diabetes patients in primary care. <i>Current Medical Research and Opinion</i> , 2013 , 29, 1495-502	2.5	42
31	Olive oil has a beneficial effect on impaired glucose regulation and other cardiometabolic risk factors. Di@bet.es study. <i>European Journal of Clinical Nutrition</i> , 2013 , 67, 911-6	5.2	27

30	Use of drugs related to the treatment of diabetes mellitus and other cardiovascular risk factors in the Spanish population. The Di@bet.es study. <i>Revista Espanola De Cardiologia (English Ed)</i> , 2013 , 66, 854-63	0.7	0
29	Consumo de fármacos relacionados con el tratamiento de la diabetes mellitus y otros factores de riesgo cardiovascular en la población española. Estudio Di@bet.es. <i>Revista Espanola De Cardiologia</i> , 2013 , 66, 854-863	1.5	6
28	Mediterranean diet adherence in individuals with prediabetes and unknown diabetes: the Di@bet.es Study. <i>Annals of Nutrition and Metabolism</i> , 2013 , 62, 339-46	4.5	17
27	The diabetic immigrant: cardiovascular risk factors and control. Contributions of the IDIME study. <i>Revista Espanola De Cardiologia</i> , 2013 , 66, 39-46	1.5	9
26	Factors determining high-sensitivity C-reactive protein values in the Spanish population. Di@bet.es study. <i>European Journal of Clinical Investigation</i> , 2013 , 43, 1-10	4.6	11
25	Factors affecting levels of urinary albumin excretion in the general population of Spain: the Di@bet.es study. <i>Clinical Science</i> , 2013 , 124, 269-77	6.5	8
24	Barriers associated with poor control in Spanish diabetic patients. A consensus study. <i>International Journal of Clinical Practice</i> , 2013 , 67, 888-94	2.9	17
23	[Cohort Study in Primary Health Care on the Evolution of Patients with Prediabetes (PREDAPS): basis and methodology]. <i>Revista Espanola De Salud Publica</i> , 2013 , 87, 121-35	0.4	17
22	Prevalence of diabetes mellitus and impaired glucose regulation in Spain: the Di@bet.es Study. <i>Diabetologia</i> , 2012 , 55, 88-93	10.3	586
21	Do Patients With High Vascular Risk Take Antihypertensive Medication Correctly? Cumple-MEMS Study. <i>Revista Espanola De Cardiologia (English Ed)</i> , 2012 , 65, 544-550	0.7	1
20	Inercia e incumplimiento a partes iguales. <i>Hipertension Y Riesgo Vascular</i> , 2012 , 29, 1-3	0.5	1
19	Do Patients With High Vascular Risk Take Antihypertensive Medication Correctly? Cumple-MEMS Study. <i>Revista Espanola De Cardiologia</i> , 2012 , 65, 544-50	1.5	20
18	Chronic kidney disease in the type 2 diabetic patients: prevalence and associated variables in a random sample of 2642 patients of a Mediterranean area. <i>BMC Nephrology</i> , 2012 , 13, 87	2.7	39
17	Trends in chronic complications of type 2 diabetic patients from Spanish primary health care centres (GEDAPS study): ten year-implementation of St. Vincent recommendations. <i>Primary Care Diabetes</i> , 2012 , 6, 11-8	2.4	20
16	Iodine intake in the adult population. Di@bet.es study. <i>Clinical Nutrition</i> , 2012 , 31, 882-8	5.9	45
15	Fifteen years of continuous improvement of quality care of type 2 diabetes mellitus in primary care in Catalonia, Spain. <i>International Journal of Clinical Practice</i> , 2012 , 66, 289-98	2.9	47
14	Evaluation of Health-Related Quality of Life according to Carbohydrate Metabolism Status: A Spanish Population-Based Study (Di@bet.es Study). <i>International Journal of Endocrinology</i> , 2012 , 2012, 872305	2.7	14
13	Control of glycemia and cardiovascular risk factors in patients with type 2 diabetes in primary care in Catalonia (Spain). <i>Diabetes Care</i> , 2012 , 35, 774-9	14.6	151

12	Validity of self-reported diabetes in health interview surveys for measuring social inequalities in the prevalence of diabetes. <i>Journal of Epidemiology and Community Health</i> , 2012 , 66, e15	5.1	27
11	Traditional risk factors alone could not explain the excess mortality in patients with diabetes: a national cohort study of older Spanish adults. <i>Diabetes Care</i> , 2012 , 35, 2503-9	14.6	17
10	Estudio observacional transversal en consultas de atención primaria y atención hospitalaria del perfil clínico del paciente con diabetes mellitus tipo 2 no controlado en fase inicial de tratamiento (estudio EUPHORIA). <i>Avances En Diabetología</i> , 2011 , 27, 69-77		3
9	La historia clínica electrónica y los registros sobre diabetes en España. <i>Avances En Diabetología</i> , 2011 , 27, 128-136		5
8	Efectos hipoglucemiantes de las resinas de intercambio. Del secuestro de los ácidos biliares a la activación del metabolismo de la glucosa. <i>Clínica E Investigación En Arteriosclerosis</i> , 2011 , 23, 3-8	1.4	
7	Long-term cardiovascular risk in type 2 diabetic compared with nondiabetic first acute myocardial infarction patients: a population-based cohort study in southern Europe. <i>Diabetes Care</i> , 2010 , 33, 2004-9	14.6	38
6	Adding questions about cardiovascular risk factors improve the ability of the ADA questionnaire to identify unknown diabetic patients in Spain. <i>Avances En Diabetología</i> , 2010 , 26, 347-352		0
5	El control metabólico en pacientes diabéticos inmigrantes. <i>Avances En Diabetología</i> , 2010 , 26, 107-111		2
4	Anthropologic study of immigrant patients with T2DM from Morocco to Spain. <i>European Diabetes Nursing</i> , 2010 , 7, 24-28		2
3	Eficacia de los fármacos hipolipemiantes: un metaanálisis. <i>FMC Formacion Medica Continuada En Atencion Primaria</i> , 2006 , 13, 650	0	
2	Impact of the American Diabetes Association diagnosis criteria on high-risk Spanish population. IGT Research Group. Impaired glucose tolerance. <i>Diabetes Research and Clinical Practice</i> , 1999 , 46, 75-81	7.4	4
1	Risk factors associated with alterations in carotid intima-media thickness in hypertension: baseline data from the European Lacidipine Study on Atherosclerosis. <i>Journal of Hypertension</i> , 1998 , 16, 949-61	1.9	214