Leonardo Bolognese

List of Publications by Year in Descending Order

Source: https://exaly.com/author-pdf/4885005/leonardo-bolognese-publications-by-year.pdf

Version: 2024-04-20

This document has been generated based on the publications and citations recorded by exaly.com. For the latest version of this publication list, visit the link given above.

The third column is the impact factor (IF) of the journal, and the fourth column is the number of citations of the article.

174	13,355	50	114
papers	citations	h-index	g-index
183	16,402 ext. citations	6.2	5.25
ext. papers		avg, IF	L-index

#	Paper	IF	Citations
174	Ticagrelor Monotherapy or Dual Antiplatelet Therapy After Drug-Eluting Stent Implantation: Per-Protocol Analysis of the GLOBAL LEADERS Trial <i>Journal of the American Heart Association</i> , 2022 , e024291	6	O
173	Safety of the oral factor XIa inhibitor asundexian compared with apixaban in patients with atrial fibrillation (PACIFIC-AF): a multicentre, randomised, double-blind, double-dummy, dose-finding phase 2 study <i>Lancet, The</i> , 2022 , 399, 1383-1390	40	9
172	Cytosorb treatment in severe COVID-19 cardiac and pulmonary disease. <i>European Heart Journal - Case Reports</i> , 2021 , 5, ytab123	0.9	O
171	Post-discharge antithrombotic management and clinical outcomes of patients with new-onset or pre-existing atrial fibrillation and acute coronary syndromes undergoing coronary stenting: Follow-up data of the MATADOR-PCI study. <i>European Journal of Internal Medicine</i> , 2021 , 88, 28-34	3.9	4
170	Combinations of antithrombotic therapies prescribed after percutaneous coronary intervention in patients with acute coronary syndromes and atrial fibrillation: data from the nationwide MATADOR-PCI registry. <i>European Heart Journal - Cardiovascular Pharmacotherapy</i> , 2021 , 7, e45-e47	6.4	2
169	Comparison of Investigator-Reported and Clinical Event Committee-Adjudicated Outcome Events in GLASSY. <i>Circulation: Cardiovascular Quality and Outcomes</i> , 2021 , 14, e006581	5.8	6
168	Potential resistance to SARS-CoV-2 infection in lysosomal storage disorders. <i>CKJ: Clinical Kidney Journal</i> , 2021 , 14, 1488-1490	4.5	1
167	Is percutaneous coronary intervention safe during uninterrupted direct oral anticoagulant therapy in patients with atrial fibrillation and acute coronary syndromes?. <i>Open Heart</i> , 2021 , 8,	3	1
166	Long-term mortality in patients undergoing lower-limb revascularization with Paclitaxel eluting devices. <i>International Journal of Cardiology</i> , 2021 , 339, 150-157	3.2	
165	Multimodality Imaging Assessment and Follow-Up in a Case of Rickettsial Myocarditis: Echocardiographic and Cardiac Magnetic Resonance Features. <i>Circulation: Cardiovascular Imaging</i> , 2021 , CIRCIMAGING121013481	3.9	
164	An autoantibody profile detects Brugada syndrome and identifies abnormally expressed myocardial proteins. <i>European Heart Journal</i> , 2020 , 41, 2878-2890	9.5	23
163	Ticagrelor versus clopidogrel in patients undergoing implantation of paclitaxel-eluting stent in the femoropopliteal district: A randomized pilot study using frequency-domain optical coherence tomography. <i>International Journal of Cardiology</i> , 2020 , 304, 192-197	3.2	3
162	Antithrombotic management of patients with acute coronary syndrome and atrial fibrillation undergoing coronary stenting: a prospective, observational, nationwide study. <i>BMJ Open</i> , 2020 , 10, e04	17044	5
161	Reduction of hospitalizations for myocardial infarction in Italy in the COVID-19 era. <i>European Heart Journal</i> , 2020 , 41, 2083-2088	9.5	437
160	Electroanatomic mapping-guided endomyocardial biopsy in patients with apparently idiopathic ventricular arrhythmias. <i>PACE - Pacing and Clinical Electrophysiology</i> , 2020 , 43, 1028-1038	1.6	2
159	Prevalence, causes and predictors of cardiovascular hospitalization in patients with hypertrophic cardiomyopathy. <i>International Journal of Cardiology</i> , 2020 , 318, 94-100	3.2	2
158	PRECISE-DAPT score for bleeding risk prediction in patients on dual or single antiplatelet regimens: insights from the GLOBAL LEADERS and GLASSY. <i>European Heart Journal - Cardiovascular Pharmacotherapy</i> , 2020 ,	6.4	13

(2018-2020)

157	Impact of preprocedural TIMI flow on clinical outcome in low-risk patients with ST-elevation myocardial infarction: Results from the ATLANTIC study. <i>Catheterization and Cardiovascular Interventions</i> , 2020 , 95, 494-500	2.7	3
156	Myocardial ischemia and coronary disease in heart failure. <i>Heart Failure Reviews</i> , 2020 , 25, 53-65	5	45
155	Reply: Myocardial Inflammation in Brugada Syndrome. <i>Journal of the American College of Cardiology</i> , 2019 , 73, 1370	15.1	
154	Rationale and design of a prospective substudy of clinical endpoint adjudication processes within an investigator-reported randomised controlled trial in patients with coronary artery disease: the GLOBAL LEADERS Adjudication Sub-StudY (GLASSY). <i>BMJ Open</i> , 2019 , 9, e026053	3	15
153	Interval From Initiation of Prasugrel to Coronary Angiography in Patients With Non-ST-Segment Elevation Myocardial Infarction. <i>Journal of the American College of Cardiology</i> , 2019 , 73, 906-914	15.1	9
152	Impact of long-term ticagrelor monotherapy following 1-month dual antiplatelet therapy in patients who underwent complex percutaneous coronary intervention: insights from the Global Leaders trial. <i>European Heart Journal</i> , 2019 , 40, 2595-2604	9.5	51
151	Drug-Eluting Balloon Versus Drug-Eluting Stent for Complex Femoropopliteal Arterial Lesions: The DRASTICO Study. <i>Journal of the American College of Cardiology</i> , 2019 , 74, 205-215	15.1	19
150	Ticagrelor Alone Versus Dual Antiplatelet Therapy From 1 Month After Drug-Eluting Coronary Stenting. <i>Journal of the American College of Cardiology</i> , 2019 , 74, 2223-2234	15.1	51
149	Trends in management and outcome of patients with non-ST elevation acute coronary syndromes and peripheral arterial disease. <i>European Journal of Internal Medicine</i> , 2019 , 59, 70-76	3.9	8
148	Time Course of Ischemic and Bleeding Burden in Elderly Patients With Acute Coronary Syndromes Randomized to Low-Dose Prasugrel or Clopidogrel. <i>Journal of the American Heart Association</i> , 2019 , 8, e010956	6	17
147	Morphine and Ticagrelor Interaction in Primary Percutaneous Coronary Intervention in ST-Segment Elevation Myocardial Infarction: ATLANTIC-Morphine. <i>American Journal of Cardiovascular Drugs</i> , 2019 , 19, 173-183	4	15
146	Pre-hospital administration of ticagrelor in diabetic patients with ST-elevation myocardial infarction undergoing primary angioplasty: A sub-analysis of the ATLANTIC trial. <i>Catheterization and Cardiovascular Interventions</i> , 2019 , 93, E369-E377	2.7	3
145	Clinical impact and predictors of complete ST segment resolution after primary percutaneous coronary intervention: A subanalysis of the ATLANTIC Trial. <i>European Heart Journal: Acute Cardiovascular Care</i> , 2019 , 8, 208-217	4.3	11
144	Comparison of Reduced-Dose Prasugrel and Standard-Dose Clopidogrel in Elderly Patients With Acute Coronary Syndromes Undergoing Early Percutaneous Revascularization. <i>Circulation</i> , 2018 , 137, 2435-2445	16.7	72
143	European Society of Cardiology: Cardiovascular Disease Statistics 2017. <i>European Heart Journal</i> , 2018 , 39, 508-579	9.5	395
142	Long-term prognostic value of delirium in elderly patients with acute cardiac diseases admitted to two cardiac intensive care units: a prospective study (DELIRIUM CORDIS). <i>European Heart Journal: Acute Cardiovascular Care</i> , 2018 , 7, 661-670	4.3	25
141	2017 ESC Guidelines for the management of acute myocardial infarction in patients presenting with ST-segment elevation: The Task Force for the management of acute myocardial infarction in patients presenting with ST-segment elevation of the European Society of Cardiology (ESC).	9.5	4237
140	Ticagrelor plus aspirin for 1 month, followed by ticagrelor monotherapy for 23 months vs aspirin plus clopidogrel or ticagrelor for 12 months, followed by aspirin monotherapy for 12 months after implantation of a drug-eluting stent: a multicentre, open-label, randomised superiority trial. <i>Lancet</i> ,	40	356

139	Duration of ischemia and treatment effects of pre- versus in-hospital ticagrelor in patients with ST-segment elevation myocardial infarction: Insights from the ATLANTIC study. <i>American Heart Journal</i> , 2018 , 196, 56-64	4.9	6
138	Electroanatomic and Pathologic Right Ventricular Outflow Tract Abnormalities in Patients With Brugada Syndrome. <i>Journal of the American College of Cardiology</i> , 2018 , 72, 2747-2757	15.1	38
137	Antithrombotic therapy before, during and after emergency angioplasty for ST elevation myocardial infarction. <i>European Heart Journal: Acute Cardiovascular Care</i> , 2017 , 6, 173-190	4.3	11
136	All-cause and cardiovascular mortality in a consecutive series of patients with diabetic foot osteomyelitis. <i>Diabetes Research and Clinical Practice</i> , 2017 , 131, 12-17	7.4	9
135	Electrical treatment of atrial arrhythmias in heart failure patients implanted with a dual defibrillator CRT device. Results from the TRADE-HF study. <i>International Journal of Cardiology</i> , 2017 , 236, 181-186	3.2	1
134	Comparative assessment of "plaque/media" change on three modalities of IVUS immediately after implantation of either everolimus-eluting bioresorbable vascular scaffold or everolimus-eluting metallic stent in Absorb II study. <i>International Journal of Cardiovascular Imaging</i> , 2017 , 33, 441-449	2.5	2
133	Italian contributions to cardiovascular research during the last 20 years. <i>European Heart Journal</i> , 2017 , 38, 130-135	9.5	
132	The Italian Federation of Cardiology. European Heart Journal, 2017, 38, 135-136	9.5	
131	Impact of presentation and transfer delays on complete ST-segment resolution before primary percutaneous coronary intervention: insights from the ATLANTIC trial. <i>EuroIntervention</i> , 2017 , 13, 69-77	, 3.1	13
130	Optimal timing of initiation of oral P2Y12-receptor antagonist therapy in patients with non-ST elevation acute coronary syndromes. Lessons learnt from the ACCOAST-trial. <i>European Heart Journal: Acute Cardiovascular Care</i> , 2016 , 5, 282-8	4.3	3
129	Effect of Pre-Hospital Ticagrelor During the First 24 h After Primary Percutaneous Coronary Intervention in Patients With ST-Segment Elevation Myocardial Infarction: The ATLANTIC-HID Analysis. <i>JACC: Cardiovascular Interventions</i> , 2016 , 9, 646-56	5	25
128	Impact of Access Site on Bleeding and Ischemic Events in Patients With Non-ST-Segment Elevation Myocardial Infarction Treated With Prasugrel: The ACCOAST Access Substudy. <i>JACC: Cardiovascular Interventions</i> , 2016 , 9, 897-907	5	20
127	Radial access in patients with acute coronary syndrome without persistent ST-segment elevation: Systematic review, collaborative meta-analysis, and meta-regression. <i>International Journal of Cardiology</i> , 2016 , 222, 1031-1039	3.2	5
126	Progression of electroanatomic substrate and electric storm recurrence in a patient with Brugada syndrome. <i>Circulation</i> , 2015 , 131, 838-41	16.7	10
125	Feasibility of combined unipolar and bipolar voltage maps to improve sensitivity of endomyocardial biopsy. <i>Circulation: Arrhythmia and Electrophysiology</i> , 2015 , 8, 625-32	6.4	46
124	Renal dysfunction, coronary revascularization and mortality among elderly patients with non ST elevation acute coronary syndrome. <i>European Heart Journal: Acute Cardiovascular Care</i> , 2015 , 4, 453-60	4.3	10
123	Ventricular arrhythmias induced by long-term use of ephedrine in two competitive athletes. <i>Heart and Vessels</i> , 2015 , 30, 280-3	2.1	14
122	Optical coherence tomography assessment and quantification of intracoronary thrombus: Status and perspectives. <i>Cardiovascular Revascularization Medicine</i> , 2015 , 16, 172-8	1.6	14

121	Prehospital ticagrelor in ST-segment elevation myocardial infarction. <i>New England Journal of Medicine</i> , 2014 , 371, 1016-27	59.2	409
120	Letter by Porto et al regarding article, "MRI-induced stent dislodgment soon after left main coronary artery stenting". <i>Circulation: Cardiovascular Interventions</i> , 2014 , 7, 127	6	1
119	One-year mortality in elderly adults with non-ST-elevation acute coronary syndrome: effect of diabetic status and admission hyperglycemia. <i>Journal of the American Geriatrics Society</i> , 2014 , 62, 1297-	363	23
118	Renal function estimation and one-year mortality in elderly patients with non-ST-segment elevation acute coronary syndromes. <i>International Journal of Cardiology</i> , 2014 , 174, 127-8	3.2	9
117	A decade of changes in clinical characteristics and management of elderly patients with non-ST elevation myocardial infarction admitted in Italian cardiac care units. <i>Open Heart</i> , 2014 , 1, e000148	3	42
116	Meta-analysis of bioabsorbable versus durable polymer drug-eluting stents in 20,005 patients with coronary artery disease: an update. <i>Catheterization and Cardiovascular Interventions</i> , 2014 , 83, E193-206	5 ^{2.} 7	25
115	Effect of prasugrel pre-treatment strategy in patients undergoing percutaneous coronary intervention for NSTEMI: the ACCOAST-PCI study. <i>Journal of the American College of Cardiology</i> , 2014 , 64, 2563-2571	15.1	53
114	Post-procedural renal microvascular perfusion measured using the Quantitative Blush Evaluator (QuBE) predicts improvement in renal function in patients undergoing percutaneous renal artery stenting. <i>International Journal of Cardiology</i> , 2014 , 172, e127-9	3.2	1
113	Outcome of patients on oral anticoagulation undergoing coronary artery stenting: data from discharge to 12 months in the Warfarin and Coronary Stenting (WAR-STENT) Registry. <i>Journal of Invasive Cardiology</i> , 2014 , 26, 563-9	0.7	23
112	Early and long-term outlook of percutaneous coronary intervention for bifurcation lesions in young patients. <i>International Journal of Cardiology</i> , 2013 , 167, 2995-9	3.2	4
111	Endothelial progenitor cells, microvascular obstruction, and left ventricular remodeling in patients with ST elevation myocardial infarction undergoing primary percutaneous coronary intervention. American Journal of Cardiology, 2013, 112, 782-91	3	13
110	Drug eluting balloon versus drug eluting stent in percutaneous coronary interventions: insights from a meta-analysis of 1462 patients. <i>International Journal of Cardiology</i> , 2013 , 168, 4608-16	3.2	16
109	Drug-eluting balloon in peripheral intervention for the superficial femoral artery: the DEBATE-SFA randomized trial (drug eluting balloon in peripheral intervention for the superficial femoral artery). JACC: Cardiovascular Interventions, 2013, 6, 1295-302	5	126
108	Pretreatment with prasugrel in non-ST-segment elevation acute coronary syndromes. <i>New England Journal of Medicine</i> , 2013 , 369, 999-1010	59.2	424
107	Comparison of pre-hospital 600 mg or 900 mg vs. peri-interventional 300 mg clopidogrel in patients with ST-elevation myocardial infarction undergoing primary coronary angioplasty. The Load&Go randomized trial. <i>International Journal of Cardiology</i> , 2013 , 168, 4814-6	3.2	31
106	Bilateral intralobar pulmonary sequestrations receiving separate arterial supply from the right and circumflex coronary arteries: a case report. <i>International Journal of Cardiology</i> , 2013 , 166, e12-3	3.2	
105	Three-year follow-up of the MULTIcentre evaluation of Single high-dose Bolus TiRofiban versus Abciximab with Sirolimus-eluting STEnt or Bare-Metal Stent in Acute Myocardial Infarction StudY (MULTISTRATEGY). <i>International Journal of Cardiology</i> , 2013 , 165, 134-41	3.2	24
104	Elutax paclitaxel-eluting balloon followed by bare-metal stent compared with Xience V drug-eluting stent in the treatment of de novo coronary stenosis: a randomized trial. <i>American Heart Journal</i> , 2013 , 166, 920-6	4.9	25

103	Stress cardiomyopathy (tako-tsubo) triggered by nervous system diseases: a systematic review of the reported cases. <i>International Journal of Cardiology</i> , 2013 , 167, 2441-8	3.2	58
102	Causes of death in patients II5 years of age with non-ST-segment elevation acute coronary syndrome. <i>American Journal of Cardiology</i> , 2013 , 112, 1-7	3	29
101	Progression of Fabry cardiomyopathy despite enzyme replacement therapy. <i>Circulation</i> , 2013 , 128, 16	8 7±8 .7	13
100	Impact of critical limb ischemia on long-term cardiac mortality in diabetic patients undergoing percutaneous coronary revascularization. <i>Diabetes Care</i> , 2013 , 36, 1495-500	14.6	9
99	Long-term morphofunctional remodeling of internal thoracic artery grafts: a frequency-domain optical coherence tomography study. <i>Circulation: Cardiovascular Interventions</i> , 2013 , 6, 269-76	6	11
98	Drug-eluting balloon in peripheral intervention for below the knee angioplasty evaluation (DEBATE-BTK): a randomized trial in diabetic patients with critical limb ischemia. <i>Circulation</i> , 2013 , 128, 615-21	16.7	235
97	Mechanisms, prevention and treatment of distal embolization. <i>Current Pharmaceutical Design</i> , 2013 , 19, 4576-85	3.3	5
96	Sex-specific benefits of sirolimus-eluting stent on long-term outcomes in patients with ST-elevation myocardial infarction undergoing primary percutaneous coronary intervention: insights from the Multicenter Evaluation of Single High-Dose Bolus Tirofiban Versus Abciximab With	4.9	12
95	Timing of events in STEMI patients treated with immediate PCI or standard medical therapy: implications on optimisation of timing of treatment from the CARESS-in-AMI trial. <i>International Journal of Cardiology</i> , 2012 , 154, 275-81	3.2	9
94	Impact of iso-osmolar versus low-osmolar contrast agents on contrast-induced nephropathy and tissue reperfusion in unselected patients with ST-segment elevation myocardial infarction undergoing primary percutaneous coronary intervention (from the Contrast Media and	3	54
93	Amiodarone-induced pulmonary toxicity mimicking acute pulmonary edema. <i>Journal of Cardiovascular Medicine</i> , 2011 , 12, 361-5	1.9	10
92	Persistent coronary no flow after wire insertion is an early and readily available mortality risk factor despite successful mechanical intervention in acute myocardial infarction: a pooled analysis from the STRATEGY (Single High-Dose Bolus Tirofiban and Sirolimus-Eluting Stent Versus Abciximab and	5	22
91	Prediction of 1-year clinical outcomes using the SYNTAX score in patients with acute ST-segment elevation myocardial infarction undergoing primary percutaneous coronary intervention: a substudy of the STRATEGY (Single High-Dose Bolus Tirofiban and Sirolinus-Eluting Stent Versus	5	111
90	Upstream use of small-molecule glycoprotein iib/iiia inhibitors in patients with non-ST-segment elevation acute coronary syndromes: a systematic overview of randomized clinical trials. <i>Circulation: Cardiovascular Quality and Outcomes</i> , 2011 , 4, 448-58	5.8	25
89	Impact of drug-eluting stents and diabetes mellitus in patients with coronary bifurcation lesions: a survey from the Italian Society of Invasive Cardiology. <i>Circulation: Cardiovascular Interventions</i> , 2011 , 4, 72-9	6	4
88	Tirofiban as adjunctive therapy for acute coronary syndromes and percutaneous coronary intervention: a meta-analysis of randomized trials. <i>European Heart Journal</i> , 2010 , 31, 35-49	9.5	83
87	Comparison of sirolimus-eluting and bare metal stent for treatment of patients with total coronary occlusions: results of the GISSOC II-GISE multicentre randomized trial. <i>European Heart Journal</i> , 2010 , 31, 2014-20	9.5	48
86	Long-term effectiveness and safety of sirolimus stent implantation for coronary in-stent restenosis results of the TRUE (Tuscany Registry of sirolimus for unselected in-stent restenosis) registry at 4	15.1	19

85	Congestive heart failure during imatinib mesylate treatment. <i>International Journal of Cardiology</i> , 2010 , 145, 148-50	3.2	16
84	Changing patterns of ST elevation myocardial infarction epidemiology. <i>American Heart Journal</i> , 2010 , 160, S1-3	4.9	11
83	The contrast media and nephrotoxicity following coronary revascularization by primary angioplasty for acute myocardial infarction study: design and rationale of the CONTRAST-AMI study. <i>Journal of Cardiovascular Medicine</i> , 2010 , 11, 199-206	1.9	3
82	Impact of acute coronary syndromes on two-year clinical outcomes in patients with unprotected left main coronary artery stenosis treated with drug-eluting stents. <i>American Journal of Cardiology</i> , 2010 , 105, 174-8	3	10
81	Randomized study of the crush technique versus provisional side-branch stenting in true coronary bifurcations: the CACTUS (Coronary Bifurcations: Application of the Crushing Technique Using Sirolimus-Eluting Stents) Study. <i>Circulation</i> , 2009 , 119, 71-8	16.7	401
80	Are drug-eluting stents superior to bare-metal stents in patients with unprotected non-bifurcational left main disease? Insights from a multicentre registry. <i>European Heart Journal</i> , 2009 , 30, 1171-9	9.5	42
79	Ostial and midshaft lesions vs. bifurcation lesions in 1111 patients with unprotected left main coronary artery stenosis treated with drug-eluting stents: results of the survey from the Italian Society of Invasive Cardiology. <i>European Heart Journal</i> , 2009 , 30, 2087-94	9.5	97
78	Comparison of drug-eluting stents and bare-metal stents for the treatment of unprotected left main coronary artery disease in acute coronary syndromes. <i>American Journal of Cardiology</i> , 2009 , 103, 187-93	3	31
77	Long-term clinical benefit of drug-eluting stents over bare-metal stents in diabetic patients with de novo left main coronary artery disease: results from a real-world multicenter registry. <i>Catheterization and Cardiovascular Interventions</i> , 2009 , 73, 310-6	2.7	4
76	Temporal pattern of ischemic events in relation to dual antiplatelet therapy in patients with unprotected left main coronary artery stenosis undergoing percutaneous coronary intervention. <i>Journal of the American College of Cardiology</i> , 2009 , 53, 1176-81	15.1	13
75	Incidence and management of restenosis after treatment of unprotected left main disease with drug-eluting stents 70 restenotic cases from a cohort of 718 patients: FAILS (Failure in Left Main Study). <i>Journal of the American College of Cardiology</i> , 2009 , 54, 1131-6	15.1	44
74	Impact of thrombus aspiration on myocardial tissue reperfusion and left ventricular functional recovery and remodeling after primary angioplasty. <i>Circulation: Cardiovascular Interventions</i> , 2009 , 2, 376-83	6	65
73	A prospective multicentre observational study on the management of patients on oral anticoagulation undergoing coronary artery stenting: rationale and design of the ongoing warfarin and coronary stenting (WAR-STENT) registry. <i>Journal of Cardiovascular Medicine</i> , 2009 , 10, 200-3	1.9	7
72	Two-year clinical outcome with drug-eluting stents versus bare-metal stents in a real-world registry of unprotected left main coronary artery stenosis from the Italian Society of Invasive Cardiology. <i>American Journal of Cardiology</i> , 2008 , 102, 1463-8	3	50
71	Immediate angioplasty versus standard therapy with rescue angioplasty after thrombolysis in the Combined Abciximab REteplase Stent Study in Acute Myocardial Infarction (CARESS-in-AMI): an open, prospective, randomised, multicentre trial. <i>Lancet, The</i> , 2008 , 371, 559-68	40	289
70	Transfer for primary angioplasty in elderly patients with acute myocardial infarction. <i>Acute Cardiac Care</i> , 2008 , 10, 152-8		O
69	Comparison of angioplasty with infusion of tirofiban or abciximab and with implantation of sirolimus-eluting or uncoated stents for acute myocardial infarction: the MULTISTRATEGY randomized trial. <i>JAMA - Journal of the American Medical Association</i> , 2008 , 299, 1788-99	27.4	214
68	Modalities of treatment and 30-day outcomes of unselected patients older than 75 years with acute ST-elevation myocardial infarction: data from the BLITZ study. <i>Journal of Cardiovascular Medicine</i> , 2008 , 9, 1045-51	1.9	5

67	Shifting from pharmacological to systematic mechanical reperfusion therapy for acute myocardial infarction via a cooperating network: impact on reperfusion rate and in-hospital mortality. <i>Journal of Cardiovascular Medicine</i> , 2008 , 9, 245-50	1.9	1
66	Early aggressive vs. initially conservative treatment in elderly patients with non-ST-elevation acute coronary syndrome: the Italian Elderly ACS study. <i>Journal of Cardiovascular Medicine</i> , 2008 , 9, 217-26	1.9	10
65	The emerging role of platelet glycoprotein IIb/IIIa inhibitors in managing high-risk patients with non-ST segment elevation acute coronary syndromes. <i>Current Medical Research and Opinion</i> , 2007 , 23, 1217-26	2.5	2
64	Antiplatelet therapy in patients undergoing coronary stent implantation: Italian Society of Interventional Cardiology consensus document. <i>Journal of Cardiovascular Medicine</i> , 2007 , 8, 782-91	1.9	4
63	Two-by-two factorial comparison of high-bolus-dose tirofiban followed by standard infusion versus abciximab and sirolimus-eluting versus bare-metal stent implantation in patients with acute myocardial infarction: design and rationale for the MULTI-STRATEGY trial. <i>American Heart Journal</i> ,	4.9	14
62	2007 , 154, 39-45 Epidemiology of non-ST elevation acute coronary syndromes in the Italian cardiology network: the BLITZ-2 study. <i>European Heart Journal</i> , 2006 , 27, 393-405	9.5	46
61	Carotid artery stenting: first consensus document of the ICCS-SPREAD Joint Committee. <i>Stroke</i> , 2006 , 37, 2400-9	6.7	83
60	Randomized comparison of upstream tirofiban versus downstream high bolus dose tirofiban or abciximab on tissue-level perfusion and troponin release in high-risk acute coronary syndromes treated with percutaneous coronary interventions: the EVEREST trial. <i>Journal of the American</i>	15.1	79
59	Effectiveness and safety of sirolimus stent implantation for coronary in-stent restenosis: the TRUE (Tuscany Registry of Sirolimus for Unselected In-Stent Restenosis) Registry. <i>Journal of the American College of Cardiology</i> , 2006 , 48, 270-5	15.1	29
58	Late acute thrombosis after implantation of sirolimus-eluting stent to treat in-stent restenosis. <i>Acute Cardiac Care</i> , 2006 , 8, 116-8		4
57	Adding tirofiban to standard pretreatment does not alter outcomes for people with non ST-elevation ACS undergoing early invasive therapy. <i>Evidence-based Cardiovascular Medicine</i> , 2006 , 10, 38-40		1
56	Elevations in troponin I after percutaneous coronary interventions are associated with abnormal tissue-level perfusion in high-risk patients with non-ST-segment-elevation acute coronary syndromes. <i>Circulation</i> , 2004 , 110, 1592-7	16.7	37
55	Impact of microvascular dysfunction on left ventricular remodeling and long-term clinical outcome after primary coronary angioplasty for acute myocardial infarction. <i>Circulation</i> , 2004 , 109, 1121-6	16.7	339
54	Combined Abciximab REteplase Stent Study in acute myocardial infarction (CARESS in AMI). <i>American Heart Journal</i> , 2004 , 148, 378-85	4.9	30
53	Epidemiology of acute myocardial infarction in the Italian CCU network: the BLITZ study. <i>European Heart Journal</i> , 2003 , 24, 1616-29	9.5	93
52	Drug-Eluting Stents. <i>Cardiology</i> , 2003 , 3, 203-213		12
51	Angiographic findings, time course of regional and global left ventricular function, and clinical outcome in diabetic patients with acute myocardial infarction treated with primary percutaneous transluminal coronary angioplasty. <i>American Journal of Cardiology</i> , 2003 , 91, 544-9	3	26
50	Comparison of impact of emergency percutaneous revascularization on outcome of patients > or =75 to those American Journal of Cardiology, 2003 , 91, 1458-61, A6	3	33

(2000-2003)

49	by cardiogenic shock who undergo percutaneous coronary intervention?. <i>Catheterization and Cardiovascular Interventions</i> , 2003 , 59, 423-8	2.7	14
48	A randomized trial comparing primary infarct artery stenting with or without abciximab in acute myocardial infarction. <i>Journal of the American College of Cardiology</i> , 2003 , 42, 1879-85	15.1	214
47	Tolerability of percutaneous coronary interventions in patients receiving nadroparin calcium for unstable angina or non-Q-wave myocardial infarction: the Angiofrax study. <i>Current Medical Research and Opinion</i> , 2003 , 19, 107-13	2.5	
46	Relation between preintervention angiographic evidence of coronary collateral circulation and clinical and angiographic outcomes after primary angioplasty or stenting for acute myocardial infarction. <i>American Journal of Cardiology</i> , 2002 , 89, 121-5	3	98
45	Relation of time to treatment and mortality in patients with acute myocardial infarction undergoing primary coronary angioplasty. <i>American Journal of Cardiology</i> , 2002 , 89, 1248-52	3	126
44	Abciximab therapy improves survival in patients with acute myocardial infarction complicated by early cardiogenic shock undergoing coronary artery stent implantation. <i>American Journal of Cardiology</i> , 2002 , 90, 353-7	3	59
43	Left ventricular remodeling after primary coronary angioplasty: patterns of left ventricular dilation and long-term prognostic implications. <i>Circulation</i> , 2002 , 106, 2351-7	16.7	438
42	Abciximab therapy improves 1-month survival rate in unselected patients with acute myocardial infarction undergoing routine infarct artery stent implantation. <i>American Heart Journal</i> , 2002 , 144, 315	- 22 9	8
41	Comparison of dobutamine echocardiography and 99mTc-sestamibi tomography for prediction of left ventricular ejection fraction outcome after acute myocardial infarction treated with successful primary coronary angioplasty. <i>Journal of Nuclear Medicine</i> , 2002 , 43, 8-14	8.9	8
40	Clinical and angiographic outcomes following elective implantation of the Carbostent in patients at high risk of restenosis and target vessel failure. <i>Catheterization and Cardiovascular Interventions</i> , 2001 , 54, 420-6	2.7	23
39	Sex-based differences in clinical and angiographic outcomes after primary angioplasty or stenting for acute myocardial infarction. <i>American Journal of Cardiology</i> , 2001 , 87, 289-93	3	73
38	Prognostic implications of restrictive left ventricular filling in reperfused anterior acute myocardial infarction. <i>Journal of the American College of Cardiology</i> , 2001 , 37, 793-9	15.1	82
37	Coronary stenting and platelet glycoprotein IIb/IIIa receptor blockade in acute myocardial infarction. <i>American Heart Journal</i> , 2001 , 141, S26-35	4.9	4
36	Direct infarct artery stenting without predilation and no-reflow in patients with acute myocardial infarction. <i>American Heart Journal</i> , 2001 , 142, 684-90	4.9	66
35	Stenting for in-stent restenosis. <i>Catheterization and Cardiovascular Interventions</i> , 2000 , 49, 376-81	2.7	18
34	Primary stenting in nonselected patients with acute myocardial infarction: the Multilink Duet in Acute Myocardial Infarction (MIAMI) trial. <i>Catheterization and Cardiovascular Interventions</i> , 2000 , 51, 27	3 ² 9 ⁷	7
33	Clinical and angiographic outcome after coronary arterial stenting with the carbostent. <i>American Journal of Cardiology</i> , 2000 , 85, 821-5	3	79
32	Cost-effective analysis of primary infarct-artery stenting versus optimal primary angioplasty (the Florence Randomized Elective Stenting in Acute Coronary Occlusions [FRESCO] trial). <i>American Journal of Cardiology</i> , 2000 , 85, 1247-9	3	5

31	Early dobutamine echocardiography predicts improvement in regional and global left ventricular function after reperfused acute myocardial infarction without residual stenosis of the infarct-related artery. <i>American Heart Journal</i> , 2000 , 139, 153-63	4.9	12
30	A randomized study of intravenous magnesium in acute myocardial infarction treated with direct coronary angioplasty. <i>American Heart Journal</i> , 2000 , 140, 891-7	4.9	19
29	Prognostic implications of Tc-99m sestamibi viability imaging and subsequent therapeutic strategy in patients with chronic coronary artery disease and left ventricular dysfunction. <i>Journal of the American College of Cardiology</i> , 2000 , 36, 739-45	15.1	59
28	Doppler-derived mitral deceleration time: an early strong predictor of left ventricular remodeling after reperfused anterior acute myocardial infarction. <i>Circulation</i> , 1999 , 99, 230-6	16.7	111
27	Primary coronary infarct artery stenting in acute myocardial infarction. <i>American Journal of Cardiology</i> , 1999 , 84, 505-10	3	15
26	Systematic primary angioplasty in octogenarian and older patients. <i>American Heart Journal</i> , 1999 , 138, 670-4	4.9	30
25	Early predictors of left ventricular remodeling after acute myocardial infarction. <i>American Heart Journal</i> , 1999 , 138, S79-83	4.9	34
24	Relation between ST-segment changes and myocardial perfusion evaluated by myocardial contrast echocardiography in patients with acute myocardial infarction treated with direct angioplasty. American Journal of Cardiology, 1998, 82, 932-7	3	188
23	Preliminary experience with stent-supported coronary angioplasty in long narrowings using the long Freedom Force stent: acute and six-month clinical and angiographic results in a series of 27 consecutive patients. <i>Catheterization and Cardiovascular Diagnosis</i> , 1998 , 43, 163-7		7
22	A clinical trial comparing primary stenting of the infarct-related artery with optimal primary angioplasty for acute myocardial infarction: results from the Florence Randomized Elective Stenting in Acute Coronary Occlusions (FRESCO) trial. <i>Journal of the American College of Cardiology</i> ,	15.1	244
21	Restenosis after coronary stenting in current clinical practice. <i>American Heart Journal</i> , 1998 , 135, 510-8	4.9	74
20	Rapid reduction of ST-segment elevation after successful direct angioplasty in acute myocardial infarction. <i>American Journal of Cardiology</i> , 1997 , 80, 685-9	3	51
19	Bailout coronary stenting without anticoagulation or intravascular ultrasound guidance: acute and six-month angiographic results in a series of 120 consecutive patients. <i>Catheterization and Cardiovascular Diagnosis</i> , 1997 , 41, 14-9		3
18	Influence of infarct-zone viability on left ventricular remodeling after acute myocardial infarction. <i>Circulation</i> , 1997 , 96, 3353-9	16.7	94
17	Myocardial contrast echocardiography versus dobutamine echocardiography for predicting functional recovery after acute myocardial infarction treated with primary coronary angioplasty. Journal of the American College of Cardiology, 1996, 28, 1677-83	15.1	124
16	Myocardial contrast versus dobutamine echocardiography as predictors of late functional recovery in acute myocardial infarction treated with primary PTCA. <i>Journal of the American College of Cardiology</i> , 1996 , 27, 22-23	15.1	17
15	Risk stratification after myocardial infarction: targets and tools. <i>Echocardiography</i> , 1995 , 12, 311-6	1.5	
14	Stress echocardiographic results predict risk of reinfarction early after uncomplicated acute myocardial infarction: large-scale multicenter study. Echo Persantine International Cooperative (EPIC) Study Group. <i>Journal of the American College of Cardiology</i> , 1995 , 26, 908-13	15.1	44

LIST OF PUBLICATIONS

13	Bailout Palmaz-Schatz coronary stenting in 39 patients with occlusive dissection complicating conventional angioplasty. <i>Catheterization and Cardiovascular Diagnosis</i> , 1995 , 35, 204-9		10
12	Prognostic value of left ventricular mass in uncomplicated acute myocardial infarction and one-vessel coronary artery disease. <i>American Journal of Cardiology</i> , 1994 , 73, 1-5	3	82
11	Prognostic value of dipyridamole echocardiography early after uncomplicated myocardial infarction: a large-scale, multicenter trial. The EPIC Study Group. <i>American Journal of Medicine</i> , 1993 , 95, 608-18	2.4	148
10	Angiographic Correlates of Different Mechanical Effects During the Dipyridamole Echocardiography Test Shortly After Acute Myocardial Infarction. <i>Echocardiography</i> , 1993 , 10, 99-106	1.5	
9	Prognostic value of dipyridamole echocardiography early after myocardial infarction in elderly patients. Echo Persantine Italian Cooperative (EPIC) Study Group. <i>Journal of the American College of Cardiology</i> , 1993 , 22, 1809-15	15.1	59
8	Oral vs intravenous dipyridamole echocardiography for detecting coronary artery disease. <i>Chest</i> , 1992 , 102, 1189-92	5.3	4
7	Silent versus symptomatic dipyridamole-induced ischemia after myocardial infarction: clinical and prognostic significance. <i>Journal of the American College of Cardiology</i> , 1992 , 19, 953-9	15.1	57
6	Safety of intravenous high-dose dipyridamole echocardiography. The Echo-Persantine International Cooperative Study Group. <i>American Journal of Cardiology</i> , 1992 , 70, 252-8	3	139
5	Comparison of intravenous urokinase plus heparin versus heparin alone in acute myocardial infarction. Urochinasi per via Sistemica nell@nfarto Miocardico (USIM) Collaborative Group. <i>American Journal of Cardiology</i> , 1991 , 68, 585-92	3	53
4	Early assessment of coronary artery bypass graft patency by high-dose dipyridamole echocardiography. <i>American Journal of Cardiology</i> , 1991 , 67, 133-6	3	15
3	Lack of influence of atrioventricular delay on stroke volume at rest in patients with complete atrioventricular block and dual chamber pacing. <i>PACE - Pacing and Clinical Electrophysiology</i> , 1990 , 13, 916-26	1.6	14
2	Clinical Experience with Orthocor II Antitachycardia Pacing System for Recurrent Tachyarrhythmia Termination. <i>Journal of Electrophysiology</i> , 1989 , 3, 289-300		4
1	High dose dipyridamole echocardiography early after uncomplicated acute myocardial infarction: correlation with exercise testing and coronary angiography. <i>Journal of the American College of Cardiology</i> , 1989 , 14, 357-63	15.1	103