

Anna Cho

List of Publications by Year in descending order

Source: <https://exaly.com/author-pdf/4890611/publications.pdf>

Version: 2024-02-01

22
papers

304
citations

1040056

9
h-index

888059

17
g-index

23
all docs

23
docs citations

23
times ranked

512
citing authors

#	ARTICLE	IF	CITATIONS
1	Cerebral Venous Sinus Thrombosis in Closed Head Injury: Systematic Review and Meta-analysis. <i>Journal of Clinical Neuroscience</i> , 2022, 98, 254-260.	1.5	4
2	How to dose-stage large or high-risk brain metastases: an alternative two-fraction radiosurgical treatment approach. <i>Journal of Neurosurgery</i> , 2022, 137, 1666-1675.	1.6	4
3	SURG-02. The site of origin of medulloblastoma: Does the neurosurgical perspective support the current concept from molecular data?. <i>Neuro-Oncology</i> , 2022, 24, i142-i142.	1.2	0
4	How can we optimize the long-term outcome in children with intracranial cavernous malformations? A single-center experience of 61 cases. <i>Neurosurgical Review</i> , 2022, 45, 3299-3313.	2.4	2
5	Increased concentrations of bioactive adrenomedullin subsequently to angiotensinâ€receptor/neprilysinâ€inhibitor treatment in chronic systolic heart failure. <i>British Journal of Clinical Pharmacology</i> , 2021, 87, 916-924.	2.4	13
6	Pre-radiosurgery leucocyte ratios and modified glasgow prognostic score predict survival in non-small cell lung cancer brain metastases patients. <i>Journal of Neuro-Oncology</i> , 2021, 151, 257-265.	2.9	9
7	Neprilysin inhibition does not alter dynamic of proenkephalinâ€A 119â€159 and proâ€substance P in heart failure. <i>ESC Heart Failure</i> , 2021, 8, 2016-2024.	3.1	3
8	The clinical relevance of laboratory prognostic scores for patients with radiosurgically treated brain metastases of non-pulmonary primary tumor. <i>Journal of Neuro-Oncology</i> , 2021, 153, 497-505.	2.9	4
9	Neutrophil-to-Lymphocyte Ratio Is Superior to Other Leukocyte-Based Ratios as a Prognostic Predictor in Nonâ€Small Cell Lung Cancer Patients with Radiosurgically Treated Brain Metastases Under Immunotherapy or Targeted Therapy. <i>World Neurosurgery</i> , 2021, 151, e324-e331.	1.3	9
10	Inflammation-Based Scores as a Common Tool for Prognostic Assessment in Heart Failure or Cancer. <i>Frontiers in Cardiovascular Medicine</i> , 2021, 8, 725903.	2.4	12
11	Second Window Indocyanine Green (SWIG) Near Infrared Fluorescent Transventricular Biopsy of Pineal Tumor. <i>World Neurosurgery</i> , 2020, 134, 196-200.	1.3	8
12	Gamma Knife Radiosurgery for Brain Metastases in Non-Small Cell Lung Cancer Patients Treated with Immunotherapy or Targeted Therapy. <i>Cancers</i> , 2020, 12, 3668.	3.7	13
13	Peri-interventional Behavior of the Neutrophil to Lymphocyte Ratio in Patients with Intracranial Aneurysms. <i>World Neurosurgery</i> , 2020, 141, e223-e230.	1.3	6
14	The inflammationâ€based modified Glasgow prognostic score is associated with survival in stable heart failure patients. <i>ESC Heart Failure</i> , 2020, 7, 654-662.	3.1	23
15	Toxicity and efficacy of Gamma Knife radiosurgery for brain metastases in melanoma patients treated with immunotherapy or targeted therapyâ€A retrospective cohort study. <i>Cancer Medicine</i> , 2020, 9, 4026-4036.	2.8	27
16	GDFâ€15 in solid vs nonâ€solid treatmentâ€naÃve malignancies. <i>European Journal of Clinical Investigation</i> , 2019, 49, e13168.	3.4	10
17	The circulating form of neprilysin is not a general biomarker for overall survival in treatment-naÃve cancer patients. <i>Scientific Reports</i> , 2019, 9, 2554.	3.3	18
18	Natural History of Functionalâ€Tricuspidâ€Regurgitation. <i>JACC: Cardiovascular Imaging</i> , 2019, 12, 389-397.	5.3	102

#	ARTICLE	IF	CITATIONS
19	N-terminal B-type natriuretic peptide (NT-proBNP) is associated with disease severity in multiple myeloma. <i>European Journal of Clinical Investigation</i> , 2018, 48, e12905.	3.4	8
20	Epidemiological Profile of Intracerebral Hemorrhage during a 10-Year Period in a Southern Brazilian Region. <i>Brazilian Neurosurgery</i> , 2018, 37, 7-12.	0.1	1
21	Non-occlusive mesenteric ischaemia in out of hospital cardiac arrest survivors. <i>European Heart Journal: Acute Cardiovascular Care</i> , 2018, 7, 450-458.	1.0	13
22	Subclinical involvement of the liver is associated with prognosis in treatment naïve cancer patients. <i>Oncotarget</i> , 2017, 8, 81250-81260.	1.8	15