

# Hui Wang

## List of Publications by Year in descending order

Source: <https://exaly.com/author-pdf/4910570/publications.pdf>

Version: 2024-02-01

22  
papers

552  
citations

567281

15  
h-index

677142

22  
g-index

22  
all docs

22  
docs citations

22  
times ranked

845  
citing authors

#	ARTICLE	IF	CITATIONS
1	A mitochondria-targeted probe containing multi-rotors for visualizing the viscosity change in living cells. <i>Australian Journal of Chemistry</i> , 2022, 75, 381-386.	0.9	2
2	A red-emitting fluorescent probe for visualizing mitochondrial microviscosity by cell imaging. <i>Chemical Papers</i> , 2021, 75, 2517-2523.	2.2	3
3	A polarity-sensitive fluorescent probe based on a difluoroboron derivative for monitoring the variation of lipid droplets. <i>New Journal of Chemistry</i> , 2021, 45, 21553-21556.	2.8	3
4	Small-molecule fluorescent dyes based on benzothiazole derivatives for targeting endoplasmic reticulum and tissue imaging. <i>Tetrahedron Letters</i> , 2020, 61, 151703.	1.4	7
5	Multiphoton Absorption Iridium(III)–Organotin(IV) Dimetal Complex with AIE Behavior for Both Sensitive Detection of Tyrosine and Antibacterial Activity. <i>ACS Applied Bio Materials</i> , 2020, 3, 8105-8112.	4.6	14
6	A fluorescent probe based on Ir(III) solvent complex for specific recognition of histidine in aqueous solution and the application in cell imaging. <i>Inorganica Chimica Acta</i> , 2020, 511, 119799.	2.4	4
7	A red-emissive mitochondrial probe for imaging of the viscosity in living cells. <i>New Journal of Chemistry</i> , 2019, 43, 8811-8815.	2.8	23
8	A series of two-photon absorption organotin (IV) cyano carboxylate derivatives for targeting nuclear and visualization of anticancer activities. <i>Journal of Inorganic Biochemistry</i> , 2019, 192, 1-6.	3.5	22
9	A reversible and highly selective two-photon fluorescent $\text{Cu}^{2+}$ probe for biological $\text{Cu}^{2+}$ detection. <i>Organic and Biomolecular Chemistry</i> , 2018, 16, 2264-2268.	2.8	21
10	A benzoic acid terpyridine-based cyclometalated iridium(III) complex as a two-photon fluorescence probe for imaging nuclear histidine. <i>Chemical Communications</i> , 2018, 54, 3771-3774.	4.1	32
11	Mitochondria-targeted iridium (III) complexes as two-photon fluorogenic probes of cysteine/homocysteine. <i>Sensors and Actuators B: Chemical</i> , 2018, 255, 408-415.	7.8	22
12	Two-Photon-Active Organotin(IV) Complexes for Antibacterial Function and Superresolution Bacteria Imaging. <i>Inorganic Chemistry</i> , 2018, 57, 6340-6348.	4.0	43
13	A reversible two-photon fluorescence probe for $\text{Cu}(\text{II})$ based on Schiff-base in HEPES buffer and in vivo imaging. <i>Sensors and Actuators B: Chemical</i> , 2017, 251, 993-1000.	7.8	36
14	Two-Photon Active Organotin(IV) Carboxylate Complexes for Visualization of Anticancer Action. <i>ACS Biomaterials Science and Engineering</i> , 2017, 3, 836-842.	5.2	40
15	Highly Hydrophilic, Two-photon Fluorescent Terpyridine Derivatives Containing Quaternary Ammonium for Specific Recognizing Ribosome RNA in Living Cells. <i>ACS Applied Materials &amp; Interfaces</i> , 2017, 9, 31424-31432.	8.0	31
16	A small-molecule with large two-photon action cross-section serves as the membrane-permeable probe for live cells imaging and bacteria viability. <i>Sensors and Actuators B: Chemical</i> , 2017, 241, 1082-1089.	7.8	16
17	A two-photon fluorescent RNA probe screened from a series of oxime-functionalized 2,2',6',6'-terpyridine $\text{ZnX}_2$ ( $\text{X} = \text{Cl}, \text{Br}, \text{I}$ ) complexes. <i>Journal of Materials Chemistry B</i> , 2016, 4, 4818-4825.	6.4	25
18	NIR-region two-photon fluorescent probes for $\text{Fe}^{3+}/\text{Cu}^{2+}$ ions based on pyrimidine derivatives with different flexible chain. <i>Sensors and Actuators B: Chemical</i> , 2016, 222, 574-578.	7.8	17

#	ARTICLE	IF	CITATIONS
19	A mitochondrial targeted two-photon iridium(III) phosphorescent probe for selective detection of hypochlorite in live cells and in vivo. <i>Biomaterials</i> , 2015, 53, 285-295.	11.4	117
20	Design of turn-on fluorescent probe for effective detection of Hg <sup>2+</sup> by combination of AIEE-active fluorophore and binding site. <i>Sensors and Actuators B: Chemical</i> , 2015, 221, 730-739.	7.8	36
21	Tunable two-photon absorption near-infrared materials containing different electron-donors and a Ir-bridge center with applications in bioimaging in live cells. <i>Journal of Materials Chemistry C</i> , 2015, 3, 5580-5588.	5.5	19
22	Synthesis, crystal structures and two-photon absorption properties of triphenylamine cyanoacetic acid derivative and its organooxotin complexes. <i>Dalton Transactions</i> , 2015, 44, 701-709.	3.3	19