

Takao Kobayashi

List of Publications by Year in descending order

Source: <https://exaly.com/author-pdf/4911930/publications.pdf>

Version: 2024-02-01

28
papers

728
citations

567281

15
h-index

526287

27
g-index

28
all docs

28
docs citations

28
times ranked

797
citing authors

#	ARTICLE	IF	CITATIONS
1	Venous Thromboembolism Deep Vein Thrombosis With Pulmonary Embolism, Deep Vein Thrombosis Alone, and Pulmonary Embolism Alone. <i>Circulation Journal</i> , 2009, 73, 305-309.	1.6	105
2	Preeclampsia as Chronic Disseminated Intravascular Coagulation. <i>Gynecologic and Obstetric Investigation</i> , 1987, 24, 170-178.	1.6	77
3	Antithrombin Therapy for Severe Preeclampsia. <i>Thrombosis and Haemostasis</i> , 2000, 84, 583-590.	3.4	66
4	Diagnosis and Management of Acute Obstetrical DIC. <i>Seminars in Thrombosis and Hemostasis</i> , 2001, 27, 161-168.	2.7	59
5	COVID-19 coagulopathy in pregnancy: Critical review, preliminary recommendations, and ISTH registry Communication from the ISTH SSC for Women's Health. <i>Journal of Thrombosis and Haemostasis</i> , 2020, 18, 3086-3098.	3.8	54
6	Obstetrical disseminated intravascular coagulation score. <i>Journal of Obstetrics and Gynaecology Research</i> , 2014, 40, 1500-1506.	1.3	42
7	Pulmonary Thromboembolism in Obstetrics and Gynecology Increased by 6.5-Fold Over the Past Decade in Japan. <i>Circulation Journal</i> , 2008, 72, 753-756.	1.6	41
8	Coagulation/Fibrinolysis Disorder in Patients with Severe Preeclampsia. <i>Seminars in Thrombosis and Hemostasis</i> , 1999, 25, 451-454.	2.7	36
9	Recombinant activated factor VII (rFVIIa/NovoSeven®) in the management of severe postpartum haemorrhage: initial report of a multicentre case series in Japan. <i>International Journal of Hematology</i> , 2012, 95, 57-63.	1.6	29
10	Unusual Pulmonary Embolism Septic Pulmonary Embolism and Amniotic Fluid Embolism. <i>Circulation Journal</i> , 2007, 71, 772-775.	1.6	28
11	Risks of thromboembolism associated with hormonal contraceptives related to body mass index and aging in Japanese women. <i>Thrombosis Research</i> , 2016, 137, 11-16.	1.7	25
12	National survey of fibrinogen concentrate usage for postpartum hemorrhage in Japan: Investigated by the Perinatology Committee, Japan Society of Obstetrics and Gynecology. <i>Journal of Obstetrics and Gynaecology Research</i> , 2015, 41, 1155-1160.	1.3	24
13	Incidence, Characteristics and Management of Venous Thromboembolism in Japan During 2011. <i>Circulation Journal</i> , 2018, 82, 555-560.	1.6	23
14	Pulmonary Embolism is an Important Cause of Death in Young Adults. <i>Circulation Journal</i> , 2007, 71, 1765-1770.	1.6	21
15	Experience with recombinant activated factor VII for severe postpartum hemorrhage in Japan, investigated by Perinatology Committee, Japan Society of Obstetrics and Gynecology. <i>Journal of Obstetrics and Gynaecology Research</i> , 2015, 41, 1161-1168.	1.3	20
16	Thromboembolism as the adverse event of combined oral contraceptives in Japan. <i>Thrombosis Research</i> , 2015, 136, 1110-1115.	1.7	13
17	Activated Protein C Is Effective for Disseminated Intravascular Coagulation Associated with Placental Abruptio. <i>Thrombosis and Haemostasis</i> , 1999, 82, 1363-1363.	3.4	9
18	Surveillance of the safety and efficacy of recombinant human soluble thrombomodulin in patients with obstetrical disseminated intravascular coagulation. <i>Thrombosis Research</i> , 2017, 159, 109-115.	1.7	9

#	ARTICLE	IF	CITATIONS
19	Risks of thromboembolism associated with hormone contraceptives in Japanese compared with Western women. <i>Journal of Obstetrics and Gynaecology Research</i> , 2017, 43, 789-797.	1.3	8
20	The incidence and prognosis of thromboembolism associated with oral contraceptives: Age-dependent difference in Japanese population. <i>Journal of Obstetrics and Gynaecology Research</i> , 2018, 44, 1766-1772.	1.3	7
21	Effect of hypofibrinogenemia on obstetrical disseminated intravascular coagulation in Japan in 2018: a multicenter retrospective cohort study. <i>International Journal of Hematology</i> , 2021, 114, 18-34.	1.6	7
22	A retrospective cohort study using a national surveillance questionnaire to investigate the characteristics of maternal venous thromboembolism in Japan in 2018. <i>BMC Pregnancy and Childbirth</i> , 2021, 21, 514.	2.4	7
23	The epidemiological characteristics of thromboembolism related to oral contraceptives in Japan: Results of a national survey. <i>Journal of Obstetrics and Gynaecology Research</i> , 2021, 47, 198-207.	1.3	6
24	Clinical guidance for peripartum management of patients with hereditary thrombophilia. <i>Journal of Obstetrics and Gynaecology Research</i> , 2021, 47, 3008-3033.	1.3	4
25	Capacity of Japanese institutions to manage obstetrical disseminated intravascular coagulation in 2018: A national surveillance questionnaire and retrospective cohort study. <i>Journal of Obstetrics and Gynaecology Research</i> , 2021, 47, 3159-3170.	1.3	4
26	Peripartum management of hereditary thrombophilia: results of primary surveillance in Japan. <i>International Journal of Hematology</i> , 2022, 116, 364-371.	1.6	3
27	National survey of confirmed thromboembolism related to female hormones in Japan. <i>Journal of Obstetrics and Gynaecology Research</i> , 2020, 46, 1173-1182.	1.3	1
28	Differences in the prevention and incidence of maternal venous thromboembolism according to the type of institution in Japan in 2018: A sub-analysis of national questionnaire surveillance. <i>Journal of Obstetrics and Gynaecology Research</i> , 2022, 48, 663-672.	1.3	0