
Antonio Rosal-Vela

List of Publications by Year in descending order

Source: https://exaly.com/author-pdf/4912268/publications.pdf Version: 2024-02-01



#	Article	IF	CITATIONS
1	IFN \hat{I}^3 induces epigenetic programming of human T-bethi B cells and promotes TLR7/8 and IL-21 induced differentiation. ELife, 2019, 8, .	6.0	116
2	Increased CD38 expression in T cells and circulating anti-CD38 IgG autoantibodies differentially correlate with distinct cytokine profiles and disease activity in systemic lupus erythematosus patients. Cytokine, 2013, 62, 232-243.	3.2	37
3	Mice Deficient in CD38 Develop an Attenuated Form of Collagen Type II-Induced Arthritis. PLoS ONE, 2012, 7, e33534.	2.5	36
4	Increased expression of microRNAâ€155 in peripheral blood mononuclear cells from psoriasis patients is related to disease activity. Journal of the European Academy of Dermatology and Venereology, 2017, 31, 312-322.	2.4	25
5	CD38 promotes pristane-induced chronic inflammation and increases susceptibility to experimental lupus by an apoptosis-driven and TRPM2-dependent mechanism. Scientific Reports, 2018, 8, 3357.	3.3	25
6	Increased expression and phosphorylation of the two S100A9 isoforms in mononuclear cells from patients with systemic lupus erythematosus: A proteomic signature for circulating low-density granulocytes. Journal of Proteomics, 2012, 75, 1778-1791.	2.4	21
7	Molecular signatures of atherosclerotic plaques: An up-dated panel of protein related markers. Journal of Proteomics, 2020, 221, 103757.	2.4	12
8	Atherosclerotic Pre-Conditioning Affects the Paracrine Role of Circulating Angiogenic Cells Ex-Vivo. International Journal of Molecular Sciences, 2020, 21, 5256.	4.1	11
9	Identification of the initial molecular changes in response to circulating angiogenic cells-mediated therapy in critical limb ischemia. Stem Cell Research and Therapy, 2020, 11, 106.	5.5	11
10	Identification of multiple transferrin species in the spleen and serum from mice with collagen-induced arthritis which may reflect changes in transferrin glycosylation associated with disease activity: The role of CD38. Journal of Proteomics, 2016, 134, 127-137.	2.4	10
11	Human canonical CD157/Bst1 is an alternatively spliced isoform masking a previously unidentified primate-specific exon included in a novel transcript. Scientific Reports, 2017, 7, 15923.	3.3	10
12	Distinct serum proteome profiles associated with collagenâ€induced arthritis and complete Freund's adjuvantâ€induced inflammation in <i>CD38^{â~'/â~'}</i> mice: The discriminative power of protein species or proteoforms. Proteomics, 2015, 15, 3382-3393.	2.2	6
13	REX-001, a BM-MNC Enriched Solution, Induces Revascularization of Ischemic Tissues in a Murine Model of Chronic Limb-Threatening Ischemia. Frontiers in Cell and Developmental Biology, 2020, 8, 602837.	3.7	4
14	Long Term Response to Circulating Angiogenic Cells, Unstimulated or Atherosclerotic Pre-Conditioned, in Critical Limb Ischemic Mice. Biomedicines, 2021, 9, 1147.	3.2	3
15	Supporting data for the MS identification of distinct transferrin glycopeptide glycoforms and citrullinated peptides associated with inflammation or autoimmunity. Data in Brief, 2016, 6, 587-602.	1.0	1
16	Assessment of endothelial colony forming cells delivery routes in a murine model of critical limb threatening ischemia using an optimized cell tracking approach. Stem Cell Research and Therapy, 2022, 13, .	5.5	1