

Ruirui Lu

List of Publications by Year in descending order

Source: <https://exaly.com/author-pdf/4914099/publications.pdf>

Version: 2024-02-01

13
papers

360
citations

933447

10
h-index

1199594

12
g-index

13
all docs

13
docs citations

13
times ranked

273
citing authors

#	ARTICLE	IF	CITATIONS
1	A molecular mechanism of UDCA engagement with GPBAR and subsequent G protein interaction revealed by scattered alanine scanning. <i>Biochemical and Biophysical Research Communications</i> , 2022, 600, 14-21.	2.1	0
2	JAK/STAT pathway promotes the progression of diabetic kidney disease via autophagy in podocytes. <i>European Journal of Pharmacology</i> , 2021, 902, 174121.	3.5	35
3	Zhen-Wu-Tang Protects IgA Nephropathy in Rats by Regulating Exosomes to Inhibit NF- κ B/NLRP3 Pathway. <i>Frontiers in Pharmacology</i> , 2020, 11, 1080.	3.5	28
4	Protective role of Astragaloside <sc>IV</sc> in chronic glomerulonephritis by activating autophagy through <sc>PI3K</sc>/<sc>AKT</sc>/<sc>AS160</sc> pathway. <i>Phytotherapy Research</i> , 2020, 34, 3236-3248.	5.8	13
5	Integrated Fecal Microbiome and Serum Metabolomics Analysis Reveals Abnormal Changes in Rats with Immunoglobulin A Nephropathy and the Intervention Effect of Zhen Wu Tang. <i>Frontiers in Pharmacology</i> , 2020, 11, 606689.	3.5	13
6	Salvianolic acid B attenuates epithelial-mesenchymal transition in renal fibrosis rats through activating Sirt1-mediated autophagy. <i>Biomedicine and Pharmacotherapy</i> , 2020, 128, 110241.	5.6	36
7	Renoprotective effects of artemisinin and hydroxychloroquine combination therapy on IgA nephropathy via suppressing NF- κ B signaling and NLRP3 inflammasome activation by exosomes in rats. <i>Biochemical Pharmacology</i> , 2019, 169, 113619.	4.4	47
8	Paeoniflorin Inhibits Mesangial Cell Proliferation and Inflammatory Response in Rats With Mesangial Proliferative Glomerulonephritis Through PI3K/AKT/GSK-3 β Pathway. <i>Frontiers in Pharmacology</i> , 2019, 10, 978.	3.5	40
9	Immunosuppressive effect of artemisinin and hydroxychloroquine combination therapy on IgA nephropathy via regulating the differentiation of CD4+ T cell subsets in rats. <i>International Immunopharmacology</i> , 2019, 70, 313-323.	3.8	26
10	Zhen-wu-tang ameliorates membranous nephropathy rats through inhibiting NF- κ B pathway and NLRP3 inflammasome. <i>Phytomedicine</i> , 2019, 59, 152913.	5.3	39
11	Immunosuppressive effects of hydroxychloroquine and artemisinin combination therapy via the nuclear factor- κ B signaling pathway in lupus nephritis mice. <i>Experimental and Therapeutic Medicine</i> , 2018, 15, 2436-2442.	1.8	24
12	Zhen-wu-tang protects against podocyte injury in rats with IgA nephropathy via PPAR γ /NF- κ B pathway. <i>Biomedicine and Pharmacotherapy</i> , 2018, 101, 635-647.	5.6	24
13	Paeoniflorin ameliorates Adriamycin-induced nephrotic syndrome through the PPAR γ /ANGPTL4 pathway in vivo and vitro. <i>Biomedicine and Pharmacotherapy</i> , 2017, 96, 137-147.	5.6	35